

THE INDIANA PREVENTION RESOURCE CENTER
2006 PREVALENCE STATISTICS
MAIN FINDINGS

**ALCOHOL, TOBACCO AND OTHER DRUG USE
BY INDIANA CHILDREN AND ADOLESCENTS**



THE INDIANA PREVENTION RESOURCE CENTER
2006 PREVALENCE STATISTICS
MAIN FINDINGS

ALCOHOL, TOBACCO AND OTHER DRUG USE BY INDIANA CHILDREN AND ADOLESCENTS

Survey Conducted March & April 2006
Report Dated August 2006

Conducted by:
Indiana Prevention Resource Center
Published by:
Institute for Drug Abuse Prevention

Department of Applied Health Science
School of Health, Physical Education, and Recreation
Indiana University

Contact:
Indiana Prevention Resource Center
2735 East 10th street
Creative Arts Building, Room 110
Bloomington, IN 47408

<http://www.drugs.indiana.edu>
drugprc@indiana.edu
Toll Free: 1-800-346-3077
Fax: 812-855-4940

Funded, in part by a contract with the
Indiana Family and Social Services Administration
Division of Mental Health and Addiction, and the
School of Health, Physical Education, and Recreation

Source Citation:

Gassman, R., Jun, M. K., Samuel, S., Martin, E. V., McCarthy-Jean, J. A., Lee, J., Kim, N., Konchada, S., Kondapuram, S. P., Morrison, A., Nautiyal, V., Pardue, N., Rayaprolu, S., Roby, R., Wang, T., and Zhou, B. (2006). Alcohol, Tobacco, and Other Drug Use by Indiana Children and Adolescents: The Indiana Prevention Resource Center Survey – 2006 (IDAP Monograph No. 06-01). Bloomington, IN: Indiana Prevention Resource Center.

Table of Contents

I. Report	1
Introduction	2
Results	5
Methodology	19
How to Interpret the Results	30
References	35
II. Prevalence Rates and Frequencies	37
 1. Statewide Data	39
Prevalence Rates by Grade: Lifetime, Annual, Monthly, Daily and Special Use	40
Grade 6	40
Grade 7	44
Grade 8	48
Grade 9	52
Grade 10	56
Grade 11	60
Grade 12	64
Frequencies by Gender, Grades 6-12	68
Lifetime, Annual, and Monthly ATOD Prevalence: All, Male, Female	68
Binge Drinking in the Past Two Weeks: All, Male, Female	104
First-Time Use of ATOD, Age Distribution: All, Male, Female	105
Prevalence Graphs, Selected Drugs	124
Selected Drugs and Binge Drinking, Monthly Prevalence, Grade 12, Indiana and USA	124
Alcohol	125
Alcohol Use, Grades 6-12	
Alcohol Use, Monthly Prevalence, by Grade and Gender	
Binge Drinking by Grade and Gender	
Tobacco	128
Cigarette Use, Grades 6-12	
Cigarette Use, Monthly Prevalence, by Grade and Gender	
Smokeless Tobacco Use, Grades 6-12	
Smokeless Tobacco, Monthly Prevalence, by Grade and Gender	
Marijuana	132
Marijuana Use, Grades 6-12	
Marijuana Use, Monthly Prevalence, by Grade and Gender	
Inhalants	134
Inhalant Use, Grades 6-12	
Trends Graphs – Selected Drugs	136
Alcohol	136
Annual / Monthly Alcohol Use, Grades 6-12	
Binge Drinking, Grades 6-12	
Binge Drinking, Grade 12, Indiana and USA Comparison	

Tobacco	140
Annual / Monthly Cigarette Use	
Monthly Prevalence, Grade 12, Indiana and USA Comparison	
Annual / Monthly Smokeless Tobacco Use	
Marijuana	145
Annual / Monthly Marijuana Use	
Monthly Prevalence, Grade 12, Indiana and USA Comparison	
Inhalants	148
Annual / Monthly Inhalant Use	
2. Regional Data	151
Prevalence Rates by Grade: Lifetime, Annual, Monthly, Daily and Special Use	152
Grade 6	152
Grade 7	156
Grade 8	160
Grade 9	164
Grade 10	168
Grade 11	172
Grade 12	176
III. Risk and Protective Factors	181
Perceptions, Grades 6-12	182
Perceived Risk of Selected Substance Use Behavior	182
Perceived Peer Approval of Using Selected Substances	185
After-School and Extra-Curricular Activities, Grades 6-12	188
Number of Days per School Year Spent in Various After-School Activities	188
Participation in Some Form of Camp	197
Participation in Organized Family Events	198
Consequences of Use, Grades 6-12	200
Annual Prevalence of ATOD Use-Related Incidents	
Safety, Grades 6-12	206
Perceived Safety in Neighborhood and in School	206
Weapon Use and Violent Behavior, Annual Prevalence	209
Weapon Use and Violent Behavior, Monthly Prevalence	215
IV. Gambling, Grades 6-12	221
Gambling during the Last Twelve Months	222
Appendix: Instrument	225

CORRECTION NOTICE

This printed report contains revisions and corrections made since the electronic version was released online on July 31, 2006, at <http://www.drugs.indiana.edu/survey/atod/2006/>. Corrections are noted on respective pages.

I. REPORT

• Introduction	2
• Results	5
• Methodology	19
• How to Interpret the Results	30
• References	35

INTRODUCTION

Overview

In Spring, 2006, the Indiana Prevention Resource Center (IPRC) conducted the *Sixteenth Annual Survey of Alcohol, Tobacco, and Other Drug Use by Indiana Children and Adolescents*, referred to hereafter as the *Indiana Survey*. The IPCR has conducted the *Indiana Survey*, and managed and reported the resulting data since 1991. This project is administered through a contract with the Division of Mental Health and Addiction of the Indiana Family and Social Services Administration (FSSA). The purpose of this project is to provide data for state and local planning with respect to the use of alcohol, tobacco, and other drugs (ATOD), gambling behaviors, and risk and protective factors.

In March and April, 2006, local school officials administered surveys to students in grades 6 through 12 in 383 schools throughout Indiana. A total of 131,017 youth from both public and nonpublic schools completed a written survey that asked about their use of various drugs, the age of onset of first use of various drugs, and risk and protective factors. This resulted in 120,914 usable surveys, the primary source of the data reported here. The Methodology section describes this process in more detail. The instrument is in the appendix.

This report summarizes the findings of the 2006 *Indiana Survey*. These findings include statistics and other information about the prevalence of ATOD use, gambling behaviors, and risk and protective factors. Statewide prevalence-of-use data are shown for past years to help distinguish trends and facilitate comparisons among years. National prevalence estimates, reported by *Monitoring the Future* (MTF) (Johnston, O'Malley, Bachman, & Schulenberg, 2005), are provided to facilitate comparison with the United States as a whole. For a summary of trends and statistically significant findings, see the Results section.

Other surveys, such as the *Youth Risk Behavior Surveillance System* (YRBSS) and MTF collect representative state-level data for Indiana. However, the results of these data are not reported for substate regions, nor are local results available to local participants. The strength of the *Indiana Survey* is its ability to describe reported ATOD use at the local level. This reporting facilitates local needs assessment, planning, and evaluation of drug abuse prevention activities. The IPCR provides a report of local results to each participating school corporation. Data from an

individual school corporation is the property of that corporation. Local data may neither be released nor obtained without the permission of the corporation where the survey was conducted.

National Outcome Measures (NOMs)

The Government Performance and Results Act of 1993 requires government agencies to report the results of their activities. In response, the United States Department of Health and Human Services (HHS), Substance Abuse and Mental Health Services Administration (SAMHSA) designated a set of measures, called the National Outcome Measures (NOMs), for substance abuse prevention. SAMHSA collects the NOMs through its National Survey on Drug Use and Health (NSDH) for all states, and it reports these data for the states and substate regions. The *Indiana Survey* collects data on the majority of the NOMs, and the IPRC also reports that data for Indiana as a whole and for the substate regions.

Increasingly, federal and state agencies require the NOMs for needs assessments and evaluation of local prevention programs that they fund. To compete for grants and to measure outcomes, communities will need to be able to capture, track, and analyze the NOMs at the local level.

A unique benefit of participating in the *Indiana Survey* is that participants receive NOMs data on their own schools and communities. The *Indiana Survey* collects NOMs data on alcohol, tobacco, marijuana and a variety of other illicit drugs for the following measures:

- thirty day prevalence of use.
- perceived risk of harm.
- age of first use.
- perception of peer disapproval.

Regional Reporting

This year, for the first time, this report presents prevalence data broken out at the level of FSSA's sub-state planning regions. These eight regions are aggregates of the State's 92 counties, as indicated in Table 1. Prevalence data for substate planning regions provides more targeted information than statewide data. Substate regional results are detailed in the Results section.

Table 1. Family and Social Service Administrations Planning Regions

Sub-state Regions	Counties
Central	Boone, Hamilton, Hancock, Hendricks, Johnson, Marion, Morgan, Shelby
East	Blackford, Delaware, Fayette, Grant, Henry, Jay, Madison, Randolph, Rush, Union, Wayne
North Central	Cass, Elkhart, Fulton, Howard, Kosciusko, La Porte, Marshall, Miami, St. Joseph, Tipton, Wabash
Northeast	Adams, Allen, De Kalb, Huntington, LaGrange, Noble, Steuben, Wells, Whitley
Northwest	Jasper, Lake, Newton, Porter, Pulaski, Starke
West	Benton, Carroll, Clay, Clinton, Fountain, Monroe, Montgomery, Owen, Parke, Putnam, Sullivan, Tippecanoe, Vermillion, Vigo, Warren, White
Southeast	Brown, Bartholomew, Decatur, Franklin, Lawrence, Jackson, Jennings, Ripley, Dearborn, Orange, Washington, Scott, Jefferson, Ohio, Switzerland, Crawford, Clark,
Southwest	Greene, Knox, Daviess, Martin, Gibson, Pike, Dubois, Posey, Vanderburgh, Warrick, Spencer, Perry

RESULTS

This section summarizes:

- statistically significant changes in statewide prevalence measures from 2005 to 2006.
- regional findings, which are reported for the first time.
- important patterns relating risk and protective factors to substance use.

Statewide Prevalence Rates

For many measures and grades, the *Indiana Survey* results yielded no statistically significant difference between the two years. Because this section focuses on changes, prevalence rates for certain grades and drugs may not be mentioned in this section. Complete results from the *Indiana Survey* appear in subsequent sections of this report.

Gateway Drugs

Research has shown that the majority of individuals' illicit drug use occurs only after they use tobacco, alcohol, or marijuana; these three are known as the "gateway drugs" (Bailey, 1992; Donovan & Jessor, 1983; Fleming, Leventhal, Glynn, & Ershler, 1989; Golub & Johnson, 2001; Kandel & Yamaguchi, 1993).

Gateway Drug Use in Grades 6-8

Statewide, the reported prevalence of cigarette, alcohol and marijuana use held steady or declined in grades 6, 7, and 8. Cigar use also decreased in these grades. These are favorable results because less gateway drug use indicates a decreased likelihood of other illicit drug use.

Gateway Drug Use in Grades 9-12

Tobacco use in grades 9-12.

Reported use of cigarettes in grades 9-12 remained unchanged from the previous year, with the exception of reported lifetime prevalence among 10th graders, which decreased. Reported daily smokeless tobacco use, on the other hand, increased

among students in grades 9-12. In addition, the reported monthly prevalence of smoking tobacco in a pipe increased among youth in grades 10-12. This is noteworthy because it suggests a shift may be occurring in the manner in which tobacco is used.

Alcohol use in grades 9-12.

The reported annual and lifetime prevalence of alcohol use among students in grades 10 and 12 declined. However, the reported prevalence of binge drinking increased among 12th graders, as did alcohol use in the last month among 9th grade students. These findings suggest that, compared to last year, a greater proportion of students in grades 10 and 12 have never started drinking. In addition, these findings suggest that the proportion of students in grades 9 and 12 who drink regularly has increased.

Marijuana use in grades 9-12.

As with the younger students, the reported prevalence of marijuana use for older students declined or held steady, as compared with 2005. The reported daily prevalence of marijuana use among youth in grades 10 and 11 decreased, and monthly prevalence decreased for students in grade 10. In addition, annual and lifetime prevalence of marijuana use decreased for students in grades 11 and 12.

Methamphetamine Use

This is the second year that the *Indiana Survey* included questions about methamphetamine.

Compared to last year, the reported prevalence of methamphetamine either held steady or declined across grade levels (see Table 2). Reported monthly and annual use of methamphetamine decreased among students in grades 8 through 10. Annual and lifetime use decreased for students in grade 6.

Table 2. Methamphetamine Use by Youth in Grades 6-12

Grade	2005	2006	Change*	Nation**
6				
Lifetime	0.7	0.5	-0.2	
Annual	0.5	0.3	-0.2	
Monthly	0.3	0.2		
7				
Lifetime	1.2	1.2		
Annual	0.9	0.8		
Monthly	0.6	0.6		
8				
Lifetime	2.4	2.0	-0.4	3.1
Annual	1.7	1.4	-0.3	1.8
Monthly	1.1	0.9	-0.2	0.7
9				
Lifetime	3.5	3.0	-0.5	
Annual	2.5	2.2	-0.3	
Monthly	1.5	1.2	-0.3	
10				
Lifetime	4.5	3.5	-1.0	4.1
Annual	3.1	2.4	-0.7	2.9
Monthly	1.6	1.2	-0.4	1.1
11				
Lifetime	4.6	4.2		
Annual	3.0	2.7		
Monthly	1.5	1.5		
12				
Lifetime	5.5	5.0		4.5
Annual	3.3	3.1		2.5
Monthly	1.7	1.5		0.9

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<0.05$).

** Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2005). Teen drug use down but progress halts among youngest teens. Retrieved December 19, 2005, from <http://www.monitoringthefuture.org>.

Other Drugs

Statistically significant changes appeared in the prevalence rates for six other drugs or drug categories (cocaine, psychedelics, heroin, methylenedioxymethamphetamine, the drug commonly known as Ecstasy or MDMA, rohypnol, and gamma hydroxyl butyrate (GHB)). The changes can be characterized as prevalence-rate changes for multiple drugs across particular grades, rather than for particular drugs across multiple grades. The broadest shifts occurred in grades 6, 8, 11, and 12. For all but one drug, rohypnol (the monthly prevalence rate for which increased among seventh graders), prevalence rates declined for youth in grades 6, 8, and 9. Meanwhile, the

statistically significant changes for eleventh and twelfth graders were all increases.

Lifetime, annual, and monthly prevalence of cocaine and MDMA use, as well as annual prevalence of non-LSD psychedelics, declined for youth in grade 6. For eighth graders, lifetime, annual, and monthly prevalence of cocaine, psychedelics overall, and LSD in particular, also declined. Ninth graders saw a decline for lifetime prevalence in psychedelics overall and for LSD in particular.

Indiana Survey data for eleventh graders indicated an increase in monthly prevalence of cocaine use, annual prevalence of heroin use, and both annual and monthly prevalence of MDMA use. The monthly prevalence of use of non-LSD psychedelics, heroin, and MDMA by twelfth graders all rose. The annual prevalence of the use of rohypnol also rose for these youth.

Needle Injection of Drugs

Youth in grade 12 indicated an increase and all time high in monthly and lifetime practice of injecting drugs (see Table 3). This finding is most alarming as injection drug use (IDU) is strongly correlated with the transmission of secondary infections such as human immunodeficiency virus (HIV), the virus that causes acquired immunodeficiency syndrome (AIDS). Injection drug use is also strongly associated with the spread of both hepatitis B and C, viral infections that lead to chronic disease and death. According to the Indiana State Department of Health, 11% of all HIV infections in 2005 were directly associated with IDU.

Table 3. *Injected Drug Use by Indiana 12th Grade Students: 2001-2006 (percentages)*

	2001	2002	2003	2004	2005	2006	Change
Lifetime	1.8	1.4	1.7	1.8	1.7	2.2	0.5 *
Annual	1.1	0.8	0.9	1.2	1.0	1.2	0.2
Monthly	0.6	0.6	0.6	0.7	0.6	0.9	0.3 *

*Statistically significant changes between 2005 and 2006 prevalence rates (p<.05).

The *Indiana Survey* does not ask directly which drugs were injected. However, a sub-analysis of responses indicating any use of a needle to inject a drug suggests that the drugs most commonly injected were heroin, methamphetamine and steroids. Differences among the prevalence rates for injection of these three drugs were not statistically significant.

Regional Prevalence Rates

This is the first year that this report presents the *Indiana Survey* results for Indiana's Family and Social Service Administration's substate planning regions (see Figure 1). Regional data provides more targeted information for local analyses. For the drugs studied, use prevalence rates varied considerably among regions and between any given region and the state. This summary does not attempt to capture the extent of this variation; however, a few examples are provided.

Compared to the state as a whole, in the...

- Southeast Region, all youth were more likely to report daily use of smokeless tobacco.
- Central Region, 7th and 9th graders were less likely to report daily use of cigarettes.
- East and Southeast Regions, 7th graders were more likely to report monthly use of tobacco, alcohol, or marijuana.

The reasons for the regional variation are not addressed in this report. Communities may gain a better understanding of their own youth drug use by comparing their prevalence rates (in their local report) to the regional rates presented in this report.




Figure 1. Indiana's Family and Social Service Administration's substate planning regions.

Risk and Protective Factors

Age of First Use

Research has shown that the younger a person is when she or he begins using alcohol, the more likely the person is to experience alcohol dependence and abuse (Grant, Stinson, & Harford, 2001; Warner & White, 2003). A recent study found that compared to persons who began drinking at age 21 or older, those who began drinking before age 14 were more likely to experience alcohol dependence later in life (Hingson, Heeren, & Winter, 2006). Communities can use age of first use to determine when to implement prevention programs. Primary prevention programs focus on intercepting drug use before it occurs and secondary prevention efforts aim to intervene early or while use is infrequent.

In the Indiana data, age of first use of gateway drugs was examined by plotting youths' age of first use of cigarettes, alcohol and marijuana since 1993 (see Figure 2). Age of first use was measured by asking, "If you have ever used these drugs, at what age did you first use them?" Over the years of *Indiana Survey* data, the average age of first use of gateway drugs has ranged consistently from 12 to 14. Reported initiation of the gateway drugs exhibits a typical pattern: cigarettes first, then alcohol, followed by marijuana. Reported age of first use did not decrease from 2005 for any of the gateway drugs.




Figure 2. Age of first-time use of a gateway drug.

Perceived Risk of Harm

Prior research has demonstrated that perceived risk of harm in using drugs is negatively related to prevalence of use (Millstein & Halpern-Felsher, 2002; Wild, Hinson, Cunningham, & Bacchichiochi, 2001).

Table 4. *Glossary of Prevalence of Gateway Drug Use for All Correlation Analyses*

Annual Use	
Cigarettes	How often in the past year have you smoked cigarettes?
Alcohol	How many times in the last year have you used alcohol (beer, wine, wine coolers, liquor)?
Marijuana	How many times in the last year have you used marijuana (hashish or hash oil)?
Monthly Use	
Cigarettes	How often in the past month have you smoked cigarettes?
Alcohol	How many times in the last month (30 days) have you used alcohol (beer, wine, wine coolers, liquor)?
Marijuana	How many times in the last month (30 days) have you used marijuana (hashish or hash oil)?

To examine this relationship in the Indiana data, respondents' perceived harm in using gateway drugs was correlated with reported monthly and annual use of gateway drugs. See Table 4 for definitions of annual and monthly use of gateway drugs. Perceived risk of harm was measured for the three gateway substances using this question: "How much do you think people risk harming themselves (physically or in other ways) if they... (a) smoke one or more packs of cigarettes per day, (b) smoke marijuana regularly, (c) have five or more drinks once or twice each weekend."

The results show that the higher the perceived risk of harm, the lower the reported annual and monthly levels of smoking cigarettes, smoking marijuana, or drinking alcohol (see Figures 3 and 4).




Figure 3. Perceived risk of harm and monthly prevalence of gateway drugs.




Figure 4. Perceived risk of harm and annual prevalence of gateway drugs.

Perceptions of Peer Disapproval

Research has demonstrated that peer perceptions of disapproval exert an influence that lowers drug use (Butters, 2004; Chassin, Presson, & Sherman, 1984). To examine this relationship in the Indiana data, respondents' perceptions of peer approval were correlated with reported use of gateway drugs. See Table 4 for definitions of annual and monthly use of gateway drugs.

Perceived peer approval was measured for the three gateway drugs using the survey items: "How do you think your close friends feel (or would feel) about you doing each of the following things... (a) smoke one or more packs of cigarettes per day; (b) smoke marijuana regularly, (c) have five or more drinks once or twice each weekend."

The results illustrate that when reported perception of peer disapproval for a particular behavior is stronger, the reported level of the behavior is lower (see Figures 5 and 6).




Figure 5. Perceived peer approval and annual prevalence of gateway drugs.




Figure 6. Perceived peer approval and monthly prevalence of gateway drugs.

Consequences of Use

Research has shown that prevention and/or reduction of alcohol and drug use among student populations leads to a decrease in related problems (Hingson et al., 1996; Wodarski, 1988). Communities are better positioned to address substance use if they understand the relationship of that use to its consequences; these include, for example, poor performance on tests, missing school, and being hung over.

The percentage of respondents who reported specific consequences of alcohol or other drug use is shown for each grade level in Table 5. Youth in grades 6 and 7 most commonly reported having trouble with parents, getting into a fight, and riding with a driver who had been drinking alcohol. About twice as many 11th and 12th grade students reported these same consequences. Considerably higher percentages of students in the 11th and 12th grade reported having had a hangover (35.5-39.6 percent) and nausea or vomiting (28.9-33.1 percent). Alarmingly, 26 percent of 12th graders—one in four—reported having ridden with a driver who had been drinking alcohol.

Table 5. *Alcohol and Other Drug Use-Related Consequences (percentages)*

Consequences of Use	Grade						
	6	7	8	9	10	11	12
Had a hangover	5.1	10.2	18.7	26.0	31.4	35.5	39.6
Performed poorly on a test or project	7.9	10.0	13.0	13.7	13.6	12.2	12.2
Been in trouble with the police	4.4	5.7	8.5	8.9	8.7	8.2	9.1
Been in trouble with their parents	10.2	12.1	16.8	18.9	20.2	20.4	19.6
Damaged property, pulled fire alarms, etc.	2.9	4.7	7.2	7.7	7.5	5.9	5.8
Got into a fight or argument	10.8	12.9	18.2	19.8	20.8	20.4	20.0
Got nauseated or vomited	6.3	9.8	15.2	20.6	25.6	28.9	33.1
Driven a car while under the influence	2.1	2.8	4.4	5.5	9.9	15.6	20.2
Ridden in a car driven by someone who had been drinking alcohol	12.2	14.3	19.3	21.0	23.6	23.9	26.1
Missed school	8.0	8.5	10.7	10.7	11.5	11.1	11.4
Had a memory loss	4.5	7.2	11.1	15.4	18.7	20.4	21.7

After-School Activities

Research has demonstrated that supervised after-school activities protect adolescents from using alcohol and other drugs (Grossman et al., 2002; Riggs & Greenberg, 2004). To determine if this relationship exists in the Indiana data, prevalence of

monthly and annual use of the gateway drugs (see Table 4) was correlated with an index score measuring student participation in after-school activities. These activities included playing on a school sports team; participating in arts, music, or drama programs; participating in supervised activities at youth centers; and classes such as music, art, computer, dance, or sports lessons. The results illustrate that the more frequent the participation in after-school activities, the lower the prevalence of reported gateway drug use (see Figures 7 and 8). These findings support the value of after-school programming as a strategy to prevent or reduce illegal substance use among adolescent populations.




Figure 7. After-school activities and annual prevalence of gateway drugs.




Figure 8. After-school activities and monthly prevalence of gateway drugs.

Personal Safety

Research suggests that violent interactions and weapon carrying are positively related to prevalence of drug use (Ash & Kellermann, 1996; Dukarm, Byrd, Auinger, & Weitzman, 1996). To investigate whether these relationships exist in the Indiana data, prevalence of annual use of gateway drugs (see Table 4) was correlated with an index score measuring violent interactions. Violent interactions were measured by asking youth “In the past year, how often have you...? (a) been in a physical fight, (b) been in a serious argument involving shouting.”

The findings indicate that the higher the frequency of involvement in violent interactions, the higher the reported prevalence of annual alcohol, cigarettes and marijuana use (see Figure 9). It is important to mention that correlation does not mean causation, but rather a relationship or association.

In addition, the Indiana prevalence data on annual use of gateway drugs (see Table 4) was correlated with an index score measuring weapon carrying to school. Weapon carrying was measured by asking youth: “In the past year, how often have you...? (a) carried a gun to school, (b) carried some other weapon to school.”

The results illustrate that the higher the frequency of weapon carrying, the higher the reported prevalence of annual gateway drug use (see Figure 10). It is important to note that all *Indiana Survey* data was collected at a single point in time. It therefore cannot distinguish the order in which the reported behaviors occurred. Therefore, the data cannot suggest a causal relationship between behaviors such as weapon carrying and drug use. However, research indicates that other deviant behaviors, such as truancy, vandalism, poor grades, and fighting, occur before drug use (Ensminger, Juon, & Fothergill, 2002; Poikolainen, 2002).




Figure 9. Violence and annual prevalence of gateway drugs.




Figure 10. Carrying weapons and annual prevalence of gateway drug use.

Gambling

As gambling options and venues in the state expand, monitoring the prevalence of gambling activities among children and adolescents is increasingly important for planning prevention efforts. This is the second year that the *Indiana Survey* included

gambling prevalence items. The *Indiana Survey* included questions about playing cards, betting on games, betting on sports, buying lottery tickets and gambling in a casino. Compared to last year, the reported annual prevalence of gambling decreased or remained the same across grade levels (see Table 6). Seventh graders were more likely to have reported gambling than 6th graders, and 8th graders were more likely to have reported gambling than seventh graders. Youth in grades 8-11 were equally likely to report having gambled. Twelfth graders were the most likely to report having gambled. Males reported significantly more gambling than females.

Table 6. Any Gambling Behavior by Indiana Adolescents (percentages)

	2005	2006	Change	
6 th	36.3	35.4	-0.9	
7 th	40.7	39.1	-1.6	*
8 th	47.4	45.3	-2.1	*
9 th	47.5	45.3	-2.2	*
10 th	48.0	45.2	-2.8	*
11 th	47.1	45.1	-2.0	*
12 th	55.1	55.0	-0.1	

*Statistically significant changes between 2005 and 2006 prevalence rates (p<.05).

Compared to 2005, a smaller proportion of students reported feeling bad about the amount of money they bet or about what happened when they bet money; the exception was grade 8 which stayed the same. In addition, a smaller proportion of students in grades 9 and 10 reported they would like to stop betting money but did not think they could.

METHODOLOGY

The methodological history of the *Indiana Survey* can be read about in previous reports. The 2005 version can be found at <http://www.drugs.indiana.edu/survey/atod/2005/index.html>.

Sample

The sampling frame of the *Indiana Survey* is the universe of all Indiana school corporations and dioceses. The IPRC obtained from the Indiana Department of Education a list of names and addresses of all 2,774 public and nonpublic schools. In November 2005, recruitment materials were sent to all school superintendents, principals, and Safe and Drug-Free School Coordinators. The recruitment packet consisted of:

- an invitation letter.
- a memorandum of understanding (MOU).
- a statement on parental consent.
- an application form.
- a list of frequently asked questions.
- a stamped addressed return envelope.
- a blank survey form.

The corporations and dioceses had a 7-8 week window to respond by sending back an application and a signed MOU to the IPRC. In order to participate, a school or corporation was required to:

- identify a designated coordinator at each public school corporation or multiple coordinators if schools are scattered in several locations.
- describe its policy on parental consent.
- identify its spring break period.
- identify desired survey administration date.
- tell the number of students scheduled to participate in each grade level.

Schools were responsible for obtaining parental consent for students to participate in the *Indiana Survey*. Information about whether a particular school or corporation

participated in any year is strictly confidential. The MOU designated the results of school and corporation-level surveys as the property of the participating school or corporation. These results are released to third parties *only* with the explicit written permission of the school or corporation's survey coordinator or other appropriate authority.

Participation

Since 1993, participation in the *Indiana Survey* has been open to all Indiana school corporations, public and nonpublic. The number of usable surveys, shown in Table 7, has fluctuated from year to year. While it is not possible to explain all of the fluctuation, the causes are likely to include:

- The exclusion from participation of 5th grade students since 1994.
- The participation of some schools on biennial and triennial bases.
- The arbitrary selection by some schools of the grade levels surveyed.
- Variation in the interpretation of parental consent requirements.
- The growing length of the *Indiana Survey* instrument.

Table 7. Number of Usable Surveys (1993-2006)

Year	Usable Surveys	Year	Usable Surveys
1993	90,586	2000	72,523
1994	81,732	2001	89,861
1995	63,631	2002	77,068
1996	36,586	2003	141,342
1997	72,571	2004	91,577
1998	44,232	2005	136,782
1999	81,685	2006	120,914

The rate of participation for school corporations, schools and students statewide and for each region is shown in Table 8. Rate of participation is the proportion of corporations, schools and students scheduled to participate in the survey out of the entire student population. There is wide variation in rates of participation across regions. The West Region had the highest rate of participation among corporations (64.1 percent), schools (25.3 percent) and students (30.9 percent) the North East Region the lowest rates, 23, 4.2 and 11.1 percent, respectively.

Table 8. Participation Rates for Students, Schools and Corporations by Region

	State	North West	North Central	North East	West	Central	East	South West	South East
Students									
Participants	131,017	16,336	13,332	6,979	17,474	34,435	14,006	10,364	18,091
Total Population	585,977	69,225	87,548	62,699	56,503	14,3383	56,905	43,198	66,516
Participation Rate ¹	22.4	23.6	15.2	11.1	30.9	24.0	24.6	24.0	27.2
Schools									
Participants	383	31	43	15	70	67	46	58	53
Total Population	2,774	300	344	357	277	600	254	320	322
Participation Rate ²	13.8	10.3	12.5	4.2	25.3	11.2	18.1	18.1	16.5
Corporations									
Participants	133	13	18	6	25	17	16	17	21
Total Population	305	34	43	26	39	43	38	36	46
Participation Rate ³	43.6	38.2	41.9	23.1	64.1	39.5	42.1	47.2	45.7

¹The percentage of student participants out of the total population of 6-12th grade students.²The percentage of schools that participated out of the total population of schools.³The percentage of school corporations that participated out of the total population of corporations.

Administration of the Survey

Two weeks before the designated survey date, the IPRC mailed to participant schools a packet containing:

- the survey forms.
- machine-readable answer sheets.
- written instructions for administering the survey.
- a twelve-minute survey administration training video produced by the IPRC.

The training video describes and demonstrates for school personnel each step of administering the survey to a group of students. The video and the written instructions were provided to improve the consistency of cross-site survey administration.

The written and video documentation included the following instructions for school personnel who administered the survey:

1. Inform students that:
 - their participation in the survey is completely voluntary.
 - the results are confidential.
 - there are no penalties for deciding not to participate or for not responding to specific items.
2. Students will need about 30 minutes to complete the survey.
3. Remain seated while students fill out the survey.
4. Instruct the students that, when they complete the survey, they should place their survey forms into the single envelope, which the IPRC has provided and which is labeled with a school and classroom code.

No individually identifying information is collected, except gender, grade level, ethnic background and school code. IPRC survey staff is available to provide support and answer questions through a toll-free 800 telephone number from Monday through Friday. Schools have a six-week window to administer the surveys between early March and mid-April and are requested to return them immediately to the IPRC.



Survey Instrument

Origin and Relevance of the Questions

Year-to-year changes in the *Indiana Survey* instrument are documented in the reports corresponding to the years in which they occur. In general, survey items have been derived from a variety of sources for purposes of comparability with national and other state survey data. All items use multiple choice response options.

Alcohol and drug use prevalence items were developed based on those used in the *MTF*(Johnston, Bachman, & O’Malley, 1989), conducted by the University of Michigan. The *Indiana Survey* asked respondents about their use of twenty different types of drugs or drug classifications (See *Indiana Survey* instrument in appendix). In some instances, examples and/or slang descriptions appeared in parentheses next to the name of the drug or drug classification. The purpose of this was to elicit responses on the use of a single substance in any of the following situations:

- The substance was a common ingredient in multiple products (i.e., alcohol in beer, wine, and liquor).
- The respondent was familiar with a slang name for the substance, but not with its proper name.
- The respondent was familiar with a slang name for the substance, but not with its proper name, and that substance had different names in different places or contexts.

For example, items related to MDMA included the terms “ecstasy,” “XTC” and “X”. Items for methamphetamines included the terms “meth,” “crank,” and “crystal.”

The *Indiana Survey* also asks about the use of a needle or syringe to inject a drug. This question is meaningful because drugs can have multiple routes of administration, and drug injection can introduce secondary infection and disease (Center for Disease Control and Prevention, 2002; Miller et al., 2004).

Prevalence of alcohol, tobacco, and a variety of other drug use is a primary focus of the *Indiana Survey*. Prevalence refers to the rate of total cases of a condition in a population. For the *Indiana Survey*, prevalence is the percentage of reported drug use among child and adolescent populations over specified time frames. Definitions for the various prevalence-of-use measures used in this report are listed below.

Lifetime prevalence: the percentage of respondents who report using a

particular drug at least once in their lifetime.

Annual prevalence: the percentage of respondents who report using a particular drug at least once during the year prior to the administration of the survey.

Current use or monthly prevalence: the percentage of respondents who report using a particular drug at least once in the 30 days prior to the administration of the survey.

Binge drinking prevalence: the percentage of respondents who report drinking at least five alcoholic drinks at a sitting in the two weeks prior to administration of the survey.

Daily prevalence: the percentage of respondents that report using a particular drug an average of once per day during the 30 day period prior to the administration of the survey. This rate is formulated based on responses of “20-40 times” or “more than 40 times” during the past month. This definition is also used by the *MTF Study*.

The perceived risk of harm items have origins in both *MTF* and the National Household Survey on Drug Abuse (NHSDA). The item set for age of first use is similar to that in the NHSDA. However, in the *Indiana Survey* these items are closed ended. The higher the perceived risk of harm the lower the reported level of drinking and other drug use.

The item sets about safety and security in the environment (for example, “How safe do you feel in school?”, “How safe do you feel in the neighborhood?”) are similar to items found in the National Longitudinal Study for Adolescent Health (Kingery, Coggeshall, & Alford, 1998). Items pertaining to personal safety such as having been in a physical fight and having been threatened by a weapon on school property are similar to those found in the YRBSS (Kingery, Coggeshall, & Alford, 1998). The questions that address carrying a gun to school or some other weapon have similarities to items used in the Teen Assessment Project (Atav & Spencer, 2002). To distinguish various purposes or motivations for carrying weapons, the *Indiana Survey* asks whether the respondent carried a gun or weapon for self defense, for hunting, or for some other reason.

The set of gambling items, added in 2005, originated from the Minnesota Student Survey (Stinchfield, Kushner, & Winters, 2005; Winters & Anderson, 2000).

Reliability

Reliability is an estimate of how consistently survey items measure the same concept each time they are administered under the same conditions and with the same subjects. Internal reliability coefficients describe the consistency of results across sets of items. Internal reliability coefficients (i.e. Cronbach's alpha) based on the average correlation among item sets are shown in Table 9. Alpha coefficients range in value from 0 to 1. The higher the score is, the more reliable the item sets. Nunnaly (1978) has indicated 0.7 to be an acceptable reliability coefficient, and many researchers require at least 0.8 for a "good scale". Based on these criteria, the internal reliability of the *Indiana Survey*'s item sets is good to excellent. This indicates that the item sets are consistent measures of the concepts in the *Indiana Survey*.

Table 9. *Internal Reliability of Item Sets*

Item Set	Question Wording	Number of Items	Cronbach's Alpha
Lifetime Use	Have you ever used...?	22	.901
Annual Use	How many times in the last year have you used...	22	.884
Use in the Past Month	How many times in the last month (30 days) have you used...	22	.838
Age of First Use	If you have ever used these drugs, at what age did you first use them?	22	.866
Consequences of Use	How many times have you experienced the following due to your drinking or drug use during the past year?	11	.903
Beliefs about Drugs	How much do you think people risk harming themselves (physically or in other ways) if they...	7	.884
Peer Perceptions	How do you think close friends feel (or would feel) about you doing each of the following things?	7	.936
Neighborhood Environment	In your neighborhood, how safe do you feel?	5	.848
School Environment	In your school, how safe do you feel?	3	.889
Personal Safety Issues	In the past year, how often have you?	12	.789
Personal Safety Issues	In the past month, how often have you?	12	.823
After School Hours	There are about 180 school days per year. On how many of those days this year will you spend most of the after school hours...	14	.775
Gambling	During the last 12 months, how often have you done these activities?	5	.812

Validity

The validity of a survey is the extent to which it measures what its designers intend it to measure. The *Indiana Survey* is intended to measure the reported prevalence of substance use, the consequences of use, peer perceptions of substance use and other concepts areas, including perceptions of personal safety and prevalence of gambling, among children and adolescents in grades 6 through 12 at a given point in time located in schools across Indiana. The *Indiana Survey* uses a nonrandom sampling procedure. Therefore, the greatest concern is external validity, or bias in the process of generalizing findings from the sample of participants to the population. A nonrandom procedure was used because a longstanding priority of the *Indiana Survey* has been to provide local level data to any and all schools or school corporations that request it. To the extent that school districts have obtained information from every student or everyone in a particular grade level, the local results represent those community populations. In this respect, participating school districts can determine their own student response rate and decide how representative it is of their local youth population. Schools can estimate their student response rate when they receive their local report by applying formula 1.0.

Formula 1.0 Response Rate = Number of Respondents / Total Number of Students x 100

Where number of respondents is the students who completed and turned the survey in, and where total number of students is the number of students intended to receive the survey.

If the response rate is 90% or greater, then the district level results (not shown in this report) are likely to be representative of the local student population.

At the regional level, the ability to generalize the data is far less because many school districts and therefore many children and adolescents in the regional population did not participate in the *Indiana Survey*. As shown in Table 8, only 11 to 31 percent of students participated in the *Indiana Survey* across regions. The best approach to address this limitation is to assess the extent to which the samples of student respondents compare to the entire population of students in each respective region. This comparison of regional respondents and the population is shown by gender, ethnicity and grade level in Table 10. Although this comparison does not provide a basis for generalizing results at the regional level, it provides guidance on where the data do and do not coincide proportionally with demographic subgroups in the regional populations.

Table 10. *Regional Comparison of Samples and Population by Demographics*

	Indiana		Northwest		Northcentral		Northeast		West	
	Pop.	Sample	Pop.	Sample	Pop.	Sample	Pop.	Sample	Pop.	Sample
Gender										
Male	51.2	49.1 *	50.9	48.6 *	51.2	49.4 *	51.1	48.9 *	51.6	49.3 *
Female	48.8	50.9 *	49.1	51.4 *	48.8	50.6 *	48.9	51.1 *	48.4	50.7 *
Ethnicity										
White	80.3	85.2 *	63.8	70.7 *	79.8	88.8 *	82.0	90.9 *	90.2	90.3
Black	11.4	6.3 *	21.1	12.7 *	9.1	1.7 *	9.8	2.1 *	2.6	2.0 *
Hispanics	4.8	3.9 *	11.7	11.0 *	7.2	6.0 *	4.7	2.8 *	3.5	2.3 *
Other [#]	3.6	4.7 *	3.4	5.5 *	4.0	3.6 *	3.5	4.3 *	3.8	5.4 *
Grade										
6	13.9	15.6 *	14.2	14.0	14.1	15.1 *	14.4	14.0	13.8	15.6 *
7	14.8	14.8	14.9	16.5 *	14.7	18.1 *	14.9	15.7 *	14.4	14.7
8	14.8	19.4 *	14.8	15.8 *	14.5	18.4 *	14.8	15.5 *	14.7	16.6 *
9	15.7	13.7 *	16.1	15.9	15.5	13.8 *	16.0	16.2 *	15.6	15.0 *
10	14.9	16.3 *	14.4	14.8	14.9	17.0 *	14.3	15.6 *	15.1	16.9 *
11	13.6	10.6 *	13.6	13.2	13.7	10.0 *	13.1	12.8 *	13.8	11.8 *
12	12.3	9.6 *	12.0	9.8 *	12.5	7.5 *	12.6	10.2 *	12.5	9.3 *
Indiana		Central		East		Southwest		Southeast		
Pop.	Sample	Pop.	Sample	Pop.	Sample	Pop.	Sample	Pop.	Sample	
Gender										
Male	51.2	49.1 *	51.1	49.2 *	51.3	48.6 *	51.2	49.4 *	51.3	49.0 *
Female	48.8	50.9 *	48.9	50.8 *	48.7	51.4 *	48.8	50.6 *	48.7	51.0 *
Ethnicity										
White	80.3	85.2 *	72.2	79.0 *	88.7	86.8 *	91.5	95.0 *	93.3	92.3 *
Black	11.4	6.3 *	19.5	11.8 *	6.5	6.6	4.7	0.8 *	2.5	2.7
Hispanics	4.8	3.9 *	4.0	3.4 *	1.5	2.1 *	1.2	1.5 *	1.4	1.6 *
Other [#]	3.6	4.7 *	4.3	5.8 *	3.2	4.5 *	2.6	2.8	2.8	3.4 *
Grade										
6	13.9	15.6 *	13.3	18.0 *	13.9	12.2 *	14.6	16.8 *	13.6	15.8 *
7	14.8	14.8 *	15.2	15.6	14.8	11.2 *	14.3	13.4 *	14.7	13.0 *
8	14.8	19.4 *	15.2	21.0 *	14.5	19.8 *	14.3	18.8 *	14.7	24.6 *
9	15.7	13.7 *	15.8	13.8 *	15.8	15.7	15.2	9.5 *	15.8	10.0 *
10	14.9	16.3 *	15.1	14.2 *	14.6	15.9 *	15.1	18.8 *	15.0	19.2 *
11	13.6	10.6 *	13.5	10.1 *	13.8	12.9 *	13.7	8.7 *	13.7	6.8 *
12	12.3	9.6 *	11.8	7.3 *	12.6	12.3	12.9	14.0 *	12.4	10.7 *

* statistically significant difference (p<.05)

Asian, American Indian, and multicultural students

As shown in Table 10, the majority of samples either under or over-represent the proportion of students by gender, ethnicity, or grade level in a region. For example, in the Northwest region the proportion of males are under represented in the survey

sample and the proportion of Whites and students of other ethnicity are over represented. This means that the sample has proportionally fewer males than what occurs in the Northwest region, and this bias should be taken into consideration when interpreting the results of the *Indiana Survey* data. Likewise, the sample has proportionally more Whites and students of other ethnicity. Thus, this is a known limitation to applying the survey results to Blacks and Hispanic groups, which are under represented in the sample.*

The strength of the *Indiana Survey* is the ability to collect and report valid and reliable data for local school corporations. At the state level, the data has the same degree of reliability. However, according to sampling theory, the state-level data is less valid (Levy & Lemeshow, 2003). This tradeoff reflects the state's current priorities for serving the substance use prevention needs of local communities. At the same time, statewide results from the *Indiana Survey* have a longstanding and high degree of concurrence with state-level surveys that use probability sampling, such as the YRBSS and the MTF survey. This is due to the large percentage of school corporations (43.6 percent) and students who participate in the *Indiana Survey*. Due to this level of agreement, it is reasonable to compare the statewide *Indiana Survey* data with national survey results. Furthermore, the results are consistent with findings from other prevention studies and this reinforces our confidence in the validity of *Indiana Survey* data at the state level.

Data

The IPRC enters the *Indiana Survey* data in-house using Pearson optical scanner equipment. The data are cleaned and screened for accuracy. An error check protocol was used to eliminate surveys that fell into three categories: (1) students who indicated he/she had not responded truthfully at all, (2) those with more than two inconsistent responses that suggested a random pattern (e.g. students who report “never used” a particular drug in their lifetime, but then report use of that same drug in the past year or month), and (3) pharmacologic implausibility (i.e. students who report such excessive use of multiple drugs as to create a cumulative effect that would be lethal).

Table 11 shows the reasons surveys were eliminated from the analysis. Failing the error check protocol described above accounted for the largest proportion (3.5 percent) of surveys eliminated. Refusal to participate or unusable surveys was the second most common reason for excluding respondents from the analysis.

* The original text stated, “Blacks are overrepresented in the Northwest region.” The correction reflects the fact that Blacks were indeed underrepresented in the Northwest region.

Table 11: Reasons for Excluding Respondents from the Analysis

	Frequency	Percent
No grade level reported	1,227	0.9
Refused to answer majority of questions	4,239	3.2
Rejected due to failing error check protocol	4,637	3.5
Total Number of Usable Surveys	120,914	92.3
Total	131,017	100.0

The final item of the *Indiana Survey* asked students, “How truthfully have you answered these questions,” with response options, (a) “not truthfully at all”, (b) “somewhat truthfully”, and (c) “completely truthfully”. The distribution of responses to this item is shown in Table 12. A significant degree of variation, from 1.1 to 1.9 percent (95% confidence interval), was found across regions for students who reported they did not answer truthfully at all. Further study is required to understand factors that contributed to this inter-regional variation. Those who responded “not truthfully at all” were eliminated from the analysis.

Table 12. Response to Item on Truthful Completion of the Indiana Survey by Region

Regions	No Answer		Not Truthfully at All		Somewhat Truthfully		Completely Truthfully		Total
	Number	%	Number	%	Number	%	Number	%	
Northwest	1,660	10.2	303	1.9	1,223	7.5	13,150	80.5	16,336
North Central	1,457	10.9	252	1.9	858	6.4	10,765	80.7	13,332
Northeast	553	7.9	103	1.5	413	5.9	5,910	84.7	6979
West	1,813	10.4	293	1.7	1,039	5.9	14,329	82.0	17,474
Central	4,251	12.3	594	1.7	2,293	6.7	27,297	79.3	34,435
East	1,535	11.0	221	1.6	1,003	7.2	11,247	80.3	14,006
Southwest	949	9.2	110	1.1	731	7.1	8,574	82.7	10,364
Southeast	1,659	9.2	259	1.4	1,259	7.0	14,914	82.4	18,091
Total	13,877	10.6	2,135	1.6	8,819	6.7	106,186	81.0	131,017

HOW TO INTERPRET THE RESULTS

Margin of Error

The *Indiana Survey* has a margin of error plus or minus less than 1 percent with a 95 percent confidence interval. This means that if we selected a group of students 100 times and asked a given question from this survey each time, then in 95 of those instances, the percentage of students giving a particular answer would be within one percentage point of the percentage who gave that same answer in this year's results of the *Indiana Survey*. Margin of error and confidence intervals, however, have limited applicability to the *Indiana Survey* because respondents were chosen nonrandomly.

'05-'06 Changes

For the statewide data tables, values shown in the 'change' column refer to statistically significant observed differences in prevalence of use between years 2005 and 2006. According to Daniel (1991), statistical significance describes a mathematical measure of difference between groups. The difference is said to be statistically significant if it is greater than what might be expected to happen by chance alone. "P" is an estimate of the probability that the result has occurred by statistical accident. In this report, the p-value is set at 0.05 which means the differences or changes indicated has a 95 percent chance of being true. The plus and minus sign next to the change value indicates whether the 2006 percentage is an increase or decrease, respectively, from the previous year.

For the regional data tables, the symbols ▲ or ▼ are used to indicate statistically significant regional differences in prevalence of use compared to the entire state. The ▲ symbol indicates that the regional rate is higher than the state rate, and the ▼ symbol indicates that the regional rate is lower than the state rate.

Generalizing Local Indiana Survey Data to Local Populations

The IPRC provides each participant school corporation with a local report of the *Indiana Survey* results from its district. This section describes the method for estimating how well local data applies to local populations. The IPRC strongly



encourages school corporations to share the results of this analysis with citizens and organizations engaged in local surveillance and planning.

The Response Rate

The response rate estimates how well the local results of the *Indiana Survey* describe all the youth in grade or grades, in a school or schools, or in an entire school corporation.

If the response rate is 90% or greater, then the results are likely to be representative of the population defined. If the response rate is lower than 90%, results should be interpreted cautiously. In this case, the results might not be representative of students who were absent the day the survey was administered or of those who did not turn in a survey.

In any case, results do not generalize to those whose responses the IPRC systematically excluded from tabulation. This occurred when a student's responses failed the error check protocol. A student's responses failed the error check protocol when the student:

- Did not provide answers to a majority of the questions
- Indicated that he or she did not respond truthfully at all.
- Provided a random pattern of responses.
- Provided a pattern of responses that was pharmacologically implausible.

The IPRC is unable to calculate the response rate for individual localities because one of the components of the response rate depends on information not collected along with the survey. This information, the Total Number of Students, is explained below.

To calculate the response rate, you need two pieces of information: the total number of students and the number of respondents.

The Total Number of Students

The total number of students can only be determined by each school district.

The total number of students is the number of students that the school district intended to receive the survey. For instance, if the intention was to survey all 9th through 10th grade students in all schools then the total enrollment number in grades

9 through 10 would constitute the total number of students.

Total Number of Respondents

Local reports of the *Indiana Survey* issued to participating school corporations include a table similar to Table 13. The asterisk (*) in Table 13 illustrates where to locate the total number of respondents.

Table 13. *Sample Table from Local Report*

	Frequency	Percent
No grade level reported		
Refused to answer majority of questions		
Rejected due to failing error check protocol		
Total Number of Usable Surveys		
Total	#,###,*	100.0

Calculating the Response Rate

To determine the response rate for a locality, divide the Total Number of Respondents by the Total Number of Students and multiply the result by 100.

$$\text{Response Rate} = \left(\frac{\text{Number of Respondents}}{\text{Total Number of Students}} \right) \times 100$$

Identifying Significant Differences between Local and State Prevalence Rates

In the local report, the first set of tables, titled “Significant Difference between Local and State Drug Use Rates, 2006,” show where a school corporation had statistically significant higher or lower rates compared to the state. Positive numbers indicate that a school corporation had rates higher than those of the state. Negative numbers indicate that a school corporation had rates lower than those of the state. Dash marks appear where there were no statistically significant differences between the



school corporation and the state.

If a school corporation had prevalence rates higher than those of the state on particular drugs, the community might want to consider targeting prevention efforts at those drugs.

Practical Significance

The next set of tables in a local report shows the percentage of student participants who reported that they used specific drugs daily, during the past month, during the past year and during their lifetime. These rates are shown for the school district over a five year period if such data are available and compared to the state and most recently available national rates.

Targeting prevention efforts at particular drugs might be appropriate even though local prevalence rates are *not* higher than those of the state. For instance, if 6th grade students in a corporation had a prevalence rate similar to that of the state for a specific drug, but the rate itself was high, prevention or treatment programs might be needed for 6th graders.

Key points for interpreting prevalence rate results of the *Indiana Survey*:

- Monthly prevalence is a general indicator of frequent or regular use of a drug.
- Annual prevalence is a general indicator of occasional or experimental use of a drug.
- “Prevalence” indicates any use of a drug within the specified time frame, regardless of its severity or frequency within that time frame (for example, any use of marijuana in the last year could be one time or two hundred times). Use monthly and annual prevalence rates in conjunction with detailed frequencies of use (for example, 1-5 times, 6-19 times, 20-40 times, more than 40 times). These are provided in tables in the back of a local report.

Key Points for Interpreting Local Prevalence Rates of Gateway Drug Use

- Cigarette smoking is the best predictor of future drug use.
- Binge drinking is a key indicator of future substance abuse.

- Alcohol use indicates a need for student assistance programs.
- Marijuana use is a key indicator of future drug use.
- Grades with the most rapid increases indicate appropriate timing of secondary prevention strategies.

If a school corporation has participated more than three times in the *Indiana Survey*, it has sufficient data to examine changes over time. If enrollment is relatively stable over time, trends and patterns may appear. A commonly observed pattern is the “cohort effect.” For certain groups of youth, prevalence rates are inconsistent with the average use rates. These differences often persist over time. For instance, 8th grade students this year may have used particular drugs 10% more than 8th grade students two years ago. This pattern may continue from year to year until the students graduate.

References

- Ash, P., & Kellermann, A. (1996). Gun acquisition and use by juvenile offenders. *JAMA*, 275(22), 1754-1758.
- Atav, S., & Spencer, G. A. (2002). Health risk behaviors among adolescents attending rural, suburban, and urban schools: A comparative study. *Family & Community Health* 25(2), 53-64.
- Bailey, S. L. (1992). Adolescents' multisubstance use patterns: The role of heavy alcohol and cigarette use. *American Journal of Public Health*, 82(9), 1220-1224.
- Butters, J. E. (2004). The impact of peers and social disapproval on high-risk cannabis use: Gender differences and implications for drug education. *Drugs Education Prevention Policy*, 11(5), 381-390.
- Center for Disease Control and Prevention. (2002). Viral hepatitis and injection drug users. Retrieved July 14, 2006, from http://www.cdc.gov/idi/hepatitis/viral_hep_drug_use.htm
- Chassin, L., Presson, C. C., & Sherman, S. J. (1984). Cognitive and social influence factors in adolescent smoking cessation. *Addictive Behaviors*, 9(4), 383-390.
- Daniel, W. W. (1991). *Biostatistics: A foundation for analysis in the health sciences* (5th ed.). New York, NY: Wiley.
- Donovan, J. E., & Jessor, R. (1983). Problem drinking and the dimension of involvement with drugs: A Guttman scalogram analysis of adolescent drug use. *American Journal of Public Health*, 73(5), 543-552.
- Dukarm, C. P., Byrd, R. S., Auinger, P., & Weitzman, M. (1996). Illicit substance use, gender, and the risk of violent behavior among adolescents. *Archives of Pediatrics & Adolescent Medicine*, 150(8), 797-801.
- Ensminger, M. E., Juon, H. S., & Fothergill, K. E. (2002). Childhood and adolescent antecedents of substance use in adulthood. *Addiction*, 97(7), 833-844.
- Fleming, R., Leventhal, H., Glynn, K., & Ershler, J. (1989). The role of cigarettes in the initiation and progression of early substance use. *Addictive Behaviors*, 14(3), 261-272.
- Golub, A., & Johnson, B. D. (2001). Variation in youthful risks of progression from alcohol and tobacco to marijuana and to hard drugs across generations. *American Journal of Public Health*, 91(2), 225-232.
- Grant, B. F., Stinson, F. S., & Harford, T. C. (2001). Age at onset of alcohol use and DSM-IV alcohol abuse and dependence: A 12-year follow-up. *Journal of Substance Abuse*, 13(4), 493-504.
- Grossman, J. B., Price, M. L., Fellerath, V., Jucovy, L. Z., Kotloff, L. J., Raley, R., et al. (2002). *Multiple choices after school: Findings from the extended-service schools initiative*. Philadelphia, PA : Public/Private Ventures.
- Hingson, R., McGovern, T., Howland, J., Heeren, T., Winter, M., & Zakocs, R. (1996). Reducing alcohol-impaired driving in Massachusetts: The saving lives program. *American Journal of Public Health*, 86(6), p791-791.
- Hingson, R. W., Heeren, T., & Winter, M. R. (2006). Age at drinking onset and

- alcohol dependence: Age at onset, duration, and severity. *Archives of Pediatrics & Adolescent Medicine, 160*(7), 739-746.
- Johnston, L. D., Bachman, J. G., & O'Malley, P. M. (1989). *Drug use, drinking, and smoking: National survey results from high school, college, and young adult populations, 1975-1989*. Rockville, MD: National Institute on Drug Abuse.
- Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2005). Teen drug use down but progress halts among youngest teens. Retrieved December 19, 2005, from <http://www.monitoringthefuture.org>
- Kandel, D., & Yamaguchi, K. (1993). From beer to crack: Developmental patterns of drug involvement. *American Journal of Public Health, 83*(6), 851-855.
- Kingery, P. M., Coggeshall, M. B., & Alford, A. A. (1998). Violence at school: Recent evidence from four national surveys. *Psychology in the Schools, 35*(3), 247-258.
- Levy, P. S., & Lemeshow, S. (2003). *Sampling of populations: Methods and applications* (3rd ed.). New York, NY: John Wiley & Sons.
- Miller, C. L., Wood, E., Spittal, P. M., Li, K., Frankish, J. C., Braitstein, P., et al. (2004). The future face of coinfection: Prevalence and incidence of HIV and hepatitis C virus coinfection among young injection drug users. *Journal of Acquired Immune Deficiency Syndromes, 36*(2), 743-749.
- Millstein, S. G., & Halpern-Felsher, B. L. (2002). Perceptions of risk and vulnerability. *The Journal of Adolescent Health, 31*(S), 10-27.
- Nunnally, J. C. (1978). *Psychometric theory* (2nd ed.). New York, NY: McGraw Hill.
- Poikolainen, K. (2002). Antecedents of substance use in adolescence. *Current Opinion in Psychiatry, 15*(3), 241-245.
- Riggs, N. R., & Greenberg, M. T. (2004). After-school youth development programs: A developmental-ecological model of current research. *Clinical Child & Family Psychology Review, 7*(3), 177-190.
- Stinchfield, R., Kushner, M. G., & Winters, K. C. (2005). Alcohol use and prior substance abuse treatment in relation to gambling problem severity and gambling treatment outcome. *Journal of Gambling Studies, 21*(3), 273-297.
- Warner, L. A., & White, H. R. (2003). Longitudinal effects of age at onset and first drinking situations on problem drinking. *Substance Use & Misuse, 38*(14), 1983-2016.
- Wild, T. C., Hinson, R., Cunningham, J., & Bacchiori, J. (2001). Perceived vulnerability to alcohol-related harm in young adults : Independent effects of risky alcohol use and drinking motives. *Experimental and Clinical Psychopharmacology, 9*(1), 117-125.
- Winters, K. C., & Anderson, N. (2000). Gambling involvement and drug use among adolescents. *Journal Of Gambling Studies 16*(2-3), 175-198.
- Wodarski, J. S. (1988). Teams-Games-Tournaments: Teaching adolescents about alcohol and driving. *Journal of Alcohol and Drug Education, 33*(3), 46-57.



II. PREVALENCE RATES AND FREQUENCIES

1. Statewide Data 39

- Prevalence Rates by Grade:
Lifetime, Annual, Monthly, Daily and Special Use
- Frequencies by Gender, Grades 6-12
- Prevalence Graphs – Selected Drugs
- Trends Graphs – Selected Drugs

2. Regional Data 151

- Prevalence Rates by Grade:
Lifetime, Annual, Monthly, Daily and Special Use

II. PREVALENCE RATES AND FREQUENCIES

1. STATEWIDE DATA

- Prevalence Rates by Grade:
Lifetime, Annual, Monthly, Daily and Special Use 40
- Frequencies by Gender, Grades 6-12
 - Lifetime, Annual, and Monthly ATOD Prevalence: All, Male, Female 68
 - Binge Drinking in the Past Two Weeks: All, Male, Female 104
 - First-Time Use of ATOD, Age Distribution: All, Male, Female 105
- Prevalence Graphs – Alcohol, Tobacco, Marijuana, and Inhalants 124
- Trends Graphs – Alcohol, Tobacco, Marijuana, and Inhalants 136



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*
Cigarettes	29.2	28.8	27.6	27.7	27.3	26.3	23.3	21.8	20.6	17.7	17.6	17.7	14.6	11.9	-2.7
Cigars	--	--	--	--	--	9.7	8.4	7.7	10.5	6.9	6.8	7.0	6.0	5.4	-0.6
Pipe	--	--	--	--	--	3.2	3.1	2.9	--	2.1	2.2	2.4	2.0	1.8	
Smokeless Tobacco	8.5	9.0	7.7	7.1	6.7	4.6	3.7	4.0	3.5	2.9	3.2	3.0	2.9	2.7	
Alcohol	37.8	36.1	32.4	29.3	32.3	25.7	26.9	25.8	29.6	29.4	29.4	29.7	26.1	25.2	
Marijuana	2.2	2.9	3.6	5.1	4.7	5.0	4.2	3.8	4.5	4.8	4.6	4.5	3.9	2.9	-1.0
Cocaine	0.1	0.5	0.6	0.8	0.8	0.9	0.7	0.7	0.8	0.7	0.8	0.7	0.8	0.6	-0.2
Crack	0.1	0.6	0.6	1.0	0.9	1.0	0.9	0.8	1.1	1.0	1.0	1.0	1.1	0.8	-0.3
Inhalants	7.8	8.4	8.1	8.1	10.1	7.9	7.3	6.4	6.9	6.1	7.2	8.1	7.3	6.5	-0.8
Amphetamines	1.5	1.6	1.3	1.5	1.5	1.3	0.9	0.9	1.0	0.8	0.7	0.7	0.6	0.5	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	0.7	0.5
Ritalin	--	--	--	--	--	1.4	1.0	1.2	1.3	1.1	1.1	1.1	1.1	0.9	-0.2
Tranquilizers	4.3	4.4	4.0	4.2	5.1	4.7	4.1	3.7	4.0	4.3	4.4	4.4	4.1	3.9	
Narcotics	1.7	1.4	1.2	1.3	1.5	1.4	1.0	0.9	1.1	1.0	1.0	1.2	0.9	0.8	
Psychedelics [#]	0.1	1.0	0.8	1.4	1.5	1.5	1.1	0.9	--	1.0	0.9	1.0	0.8	0.6	
LSD	--	--	--	--	--	--	--	--	--	0.8	0.7	0.7	0.5	0.5	
Others	--	--	--	--	--	--	--	--	--	0.5	0.5	0.6	0.5	0.4	
Heroin	0.1	0.5	0.4	0.8	0.7	0.9	0.6	0.6	0.8	0.7	0.7	0.7	0.6	0.5	
Steroids	1.2	1.5	1.0	1.4	1.4	1.3	1.4	1.2	1.1	0.9	1.0	1.1	0.9	0.8	
Injected Drugs	--	--	--	--	--	--	--	--	0.9	0.8	0.8	1.0	1.0	0.9	
MDMA	--	--	--	--	--	--	--	--	--	0.7	0.6	0.7	0.5	0.4	-0.1
Rohypnol	--	--	--	--	--	--	--	--	--	0.4	0.4	0.4	0.4	0.4	
GHB	--	--	--	--	--	--	--	--	--	0.3	0.3	0.3	0.3	0.3	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*	
Cigarettes	19.0	19.5	18.3	18.4	18.3	16.4	14.2	12.5	11.2	9.4	9.5	9.5	7.9	6.2	-1.7
Cigars	--	--	--	--	--	7.4	6.3	5.5	7.0	4.5	4.3	4.6	4.0	3.4	-0.6
Pipe	--	--	--	--	--	2.3	1.7	1.7	--	1.4	1.5	1.6	1.3	1.1	-0.2
Smokeless Tobacco	6.4	6.8	5.8	5.4	4.9	3.2	2.6	2.6	2.5	2.1	2.2	2.2	2.0	1.9	
Alcohol	27.7	27.3	23.2	21.7	23.7	19.3	21.0	20.3	21.4	20.4	20.7	20.7	17.6	17.3	
Marijuana	1.7	2.5	3.1	4.4	4.0	4.2	3.6	3.2	3.6	3.6	3.5	3.7	3.0	2.3	-0.7
Cocaine	0.1	0.4	0.4	0.8	0.7	0.7	0.5	0.6	0.7	0.5	0.6	0.6	0.5	0.4	-0.1
Crack	0.1	0.4	0.3	0.9	0.7	0.8	0.7	0.6	0.8	0.6	0.8	0.8	0.7	0.5	-0.2
Inhalants	6.1	6.2	5.8	6.4	7.9	6.0	5.4	4.5	5.2	4.1	5.2	5.9	5.1	4.5	-0.6
Amphetamines	1.2	1.3	1.0	1.4	1.3	1.1	0.7	0.7	0.7	0.6	0.6	0.6	0.4	0.4	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	0.5	0.3	-0.2
Ritalin	--	--	--	--	--	1.2	0.8	0.9	0.9	0.7	0.7	0.8	0.7	0.5	-0.2
Tranquilizers	3.5	3.0	3.2	4.1	3.8	3.2	2.8	3.0	3.0	3.1	3.2	3.0	2.7	-0.3	
Narcotics	1.3	1.0	0.9	1.0	1.3	1.0	0.6	0.7	0.9	0.6	0.7	0.8	0.6	0.5	
Psychedelics [#]	0.1	0.7	0.7	1.2	1.2	1.3	0.8	0.7	--	0.6	0.6	0.7	0.5	0.4	
LSD	--	--	--	--	--	--	--	--	--	0.5	0.5	0.5	0.3	0.3	
Others	--	--	--	--	--	--	--	--	--	0.3	0.3	0.4	0.3	0.2	-0.1
Heroin	0.1	0.4	0.2	0.6	0.5	0.7	0.4	0.4	0.6	0.4	0.5	0.5	0.4	0.3	
Steroids	0.1	1.0	0.6	1.0	1.0	1.1	0.8	0.8	0.8	0.6	0.7	0.8	0.6	0.5	
Injected Drugs	--	--	--	--	--	--	--	--	0.6	0.5	0.7	0.7	0.7	0.5	
MDMA	--	--	--	--	--	--	--	--	--	0.5	0.5	0.5	0.4	0.3	-0.1
Rohypnol	--	--	--	--	--	--	--	--	--	0.2	0.3	0.3	0.2	0.2	
GHB	--	--	--	--	--	--	--	--	--	0.2	0.2	0.2	0.2	0.1	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*
Cigarettes	8.5	10.0	9.3	9.7	9.5	8.2	7.0	5.7	6.0	4.8	5.1	5.4	4.2	3.3	-0.9
Cigars	--	--	--	--	--	3.9	3.3	2.9	3.8	2.4	2.5	2.5	2.2	1.8	-0.4
Pipe	--	--	--	--	--	1.5	1.0	0.8	--	0.8	0.9	0.8	0.7	0.7	
Smokeless Tobacco	3.5	4.0	3.3	3.1	3.1	1.9	1.4	1.4	1.4	1.2	1.4	1.3	1.1	1.1	
Alcohol	13.6	13.7	11.4	10.7	11.5	9.5	9.9	9.2	9.5	9.5	9.9	9.8	8.1	7.6	-0.5
Marijuana	1.2	1.7	1.9	3.0	2.7	2.2	2.2	1.8	2.4	2.2	2.3	2.5	1.8	1.4	-0.4
Cocaine	0.1	0.3	0.3	0.6	0.5	0.5	0.3	0.4	0.4	0.3	0.4	0.4	0.4	0.2	-0.2
Crack	0.1	0.3	0.3	0.6	0.5	0.6	0.3	0.4	0.4	0.4	0.4	0.5	0.5	0.3	-0.2
Inhalants	3.8	4.1	3.6	3.8	4.9	3.6	3.4	2.7	3.1	2.5	3.2	3.6	3.1	2.9	
Amphetamines	0.1	0.9	0.6	1.0	0.9	0.6	0.5	0.3	0.4	0.4	0.4	0.3	0.2	0.2	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	0.3	0.2
Ritalin	--	--	--	--	--	0.7	0.5	0.5	0.6	0.4	0.4	0.5	0.4	0.3	-0.1
Tranquilizers	2.0	2.2	1.6	1.9	2.4	2.0	1.8	1.6	1.7	1.7	1.8	1.9	1.7	1.5	-0.2
Narcotics	0.1	0.5	0.6	0.6	0.9	0.7	0.4	0.5	0.5	0.4	0.4	0.5	0.4	0.3	0.3
Psychedelics [#]	0.1	0.6	0.5	0.8	0.9	0.8	0.5	0.4	--	0.4	0.4	0.4	0.3	0.3	0.2
LSD	--	--	--	--	--	--	--	--	--	0.4	0.3	0.3	0.3	0.2	
Others	--	--	--	--	--	--	--	--	--	0.2	0.2	0.2	0.2	0.1	-0.1
Heroin	0.1	0.3	0.2	0.5	0.4	0.4	0.3	0.3	0.4	0.3	0.4	0.3	0.3	0.2	
Steroids	0.1	0.6	0.4	0.7	0.7	0.6	0.5	0.4	0.5	0.4	0.4	0.5	0.4	0.3	
Injected Drugs	--	--	--	--	--	--	--	--	0.4	0.3	0.4	0.4	0.5	0.3	-0.2
MDMA	--	--	--	--	--	--	--	--	--	0.4	0.3	0.3	0.3	0.1	-0.2
Rohypnol	--	--	--	--	--	--	--	--	--	0.2	0.2	0.1	0.2	0.1	
GHB	--	--	--	--	--	--	--	--	--	0.1	0.1	0.1	0.1	0.1	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*
CIGARETTES														
Daily Use	3.4	4.3	4.0	4.2	4.3	3.7	3.1	2.3	2.6	1.9	2.2	2.1	1.9	1.4
1/2+ Pack per Day	1.5	2.0	1.9	2.1	2.1	1.7	1.4	1.0	1.2	0.9	0.8	0.9	1.0	0.7
CIGARS	--	--	--	--	--	0.6	0.5	0.3	0.6	0.2	0.3	0.4	0.3	0.2
SMOKELESS TOBACCO														
0.1	0.6	0.4	0.5	0.5	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
ALCOHOL														
Daily Use	0.1	0.4	0.5	0.4	0.6	0.5	0.4	0.4	0.5	0.5	0.6	0.4	0.5	0.3
Binge Drinking	8.0	8.1	7.2	7.6	7.6	8.7	8.2	7.9	6.7	6.6	5.3	5.9	4.5	4.2
MARIJUANA	0.1	0.1	0.3	0.4	0.3	0.4	0.4	0.2	0.4	0.4	0.5	0.4	0.4	0.3

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 7th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*
Cigarettes	38.6	40.1	39.8	40.2	38.5	35.1	34.0	31.2	29.1	26.3	23.4	24.6	22.0	19.7	-2.3
Cigars	--	--	--	--	--	15.1	14.3	12.3	14.9	11.5	9.8	10.6	9.6	8.3	-1.3
Pipe	--	--	--	--	--	5.1	5.0	5.0	--	3.6	3.6	3.8	3.2	2.9	
Smokeless Tobacco	13.9	14.9	14.6	12.7	11.5	8.1	7.3	5.9	5.2	5.3	4.8	6.1	4.8	5.0	
Alcohol	48.7	47.4	44.4	44.0	42.4	36.2	37.5	36.5	39.5	39.4	37.7	38.3	34.0	33.3	
Marijuana	5.3	7.7	9.5	12.8	10.2	9.9	9.4	8.8	9.8	9.3	8.7	8.7	8.0	6.8	-1.2
Cocaine	1.0	1.2	1.3	2.1	1.6	1.6	1.6	1.2	1.2	1.5	1.6	1.6	1.3	1.5	
Crack	0.1	1.3	1.3	1.9	1.4	1.7	1.6	1.4	1.5	1.8	1.6	1.9	1.6	1.5	
Inhalants	9.8	12.0	12.3	13.5	12.9	10.6	8.8	7.9	7.8	7.6	8.8	10.3	8.9	9.0	
Amphetamines	3.9	4.5	4.6	5.7	3.5	3.0	2.5	2.4	2.1	2.4	2.1	2.0	1.5	1.4	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	1.2
Ritalin	--	--	--	--	--	2.5	2.1	2.4	1.9	2.0	1.9	1.9	1.9	1.8	
Tranquilizers	6.5	6.8	6.7	8.5	7.4	7.1	6.2	6.4	6.0	6.8	6.5	7.2	6.0	6.0	
Narcotics	3.0	2.9	2.8	3.6	3.4	2.9	2.4	2.3	2.1	2.4	2.2	2.2	2.0	2.0	
Psychedelics [#]	1.8	2.2	2.4	3.7	3.0	3.1	2.5	2.4	--	2.1	2.0	1.8	1.6	1.5	
LSD	--	--	--	--	--	--	--	--	--	1.7	1.6	1.5	1.2	1.1	
Others	--	--	--	--	--	--	--	--	--	0.9	0.9	0.8	0.8	0.8	
Heroin	1.0	1.2	1.1	1.2	1.3	1.1	1.0	0.9	1.0	1.1	1.2	1.0	1.1		
Steroids	2.2	2.4	1.7	2.0	1.8	1.6	2.1	1.5	1.3	1.6	1.5	1.6	1.3	1.2	
Injected Drugs	--	--	--	--	--	--	--	--	0.9	1.3	1.3	1.5	1.3	1.3	
MDMA	--	--	--	--	--	--	--	--	--	1.8	1.8	1.5	1.2	1.2	
Rohypnol	--	--	--	--	--	--	--	--	--	0.9	0.7	0.7	0.6	0.8	
GHB	--	--	--	--	--	--	--	--	--	0.5	0.5	0.4	0.4	0.5	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 7th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*
Cigarettes	28.4	29.7	29.8	31.4	29.0	25.2	22.7	20.0	18.1	15.7	13.9	15.2	13.0	11.7
Cigars	--	--	--	--	--	12.6	11.0	9.5	10.5	7.9	6.4	7.4	6.6	5.8
Pipe	--	--	--	--	--	3.7	3.2	2.9	--	2.6	2.7	2.8	2.1	2.1
Smokeless Tobacco	10.9	11.7	11.9	10.2	8.8	6.4	5.1	3.9	3.8	3.9	3.6	4.5	3.5	3.9
Alcohol	39.8	38.7	36.1	36.9	34.5	30.3	31.7	30.8	31.6	30.1	29.2	29.8	25.7	25.5
Marijuana	4.5	6.7	8.6	11.5	9.1	8.6	8.1	7.9	8.3	7.8	7.1	7.2	6.3	5.6
Cocaine	0.1	1.1	1.1	1.8	1.4	1.3	1.3	1.1	1.0	1.2	1.3	1.3	1.0	1.2
Crack	0.1	0.9	1.0	1.6	1.2	1.3	1.2	1.3	1.2	1.1	1.3	1.4	1.1	1.2
Inhalants	7.6	9.6	9.3	11.4	10.3	7.8	6.4	5.7	5.6	4.9	6.2	7.6	6.4	6.3
Amphetamines	3.3	3.9	4.0	5.1	3.2	2.4	2.0	2.0	1.8	1.8	1.5	1.5	1.1	1.1
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	0.9	0.8
Ritalin	--	--	--	--	--	2.0	1.6	2.0	1.5	1.4	1.4	1.3	1.3	1.3
Tranquilizers	5.2	5.6	5.4	7.2	6.2	5.9	5.0	5.2	5.0	5.1	4.9	5.5	4.5	4.6
Narcotics	2.4	2.2	2.2	3.1	2.7	2.2	1.9	1.8	1.7	1.7	1.5	1.7	1.4	1.4
Psychedelics [#]	1.5	1.7	2.2	3.2	2.6	2.6	2.1	2.1	--	1.5	1.5	1.4	1.2	1.1
LSD	--	--	--	--	--	--	--	--	--	1.3	1.2	1.1	0.9	0.8
Others	--	--	--	--	--	--	--	--	--	0.6	0.6	0.6	0.5	0.6
Heroin	0.1	0.9	0.9	1.3	1.0	0.9	0.8	0.7	0.8	0.9	0.9	1.0	0.7	0.8
Steroids	1.6	1.8	1.2	1.5	1.4	1.1	1.5	1.1	1.0	1.2	1.1	1.3	0.9	0.9
Injected Drugs	--	--	--	--	--	--	--	--	0.7	1.0	0.9	1.0	0.9	1.0
MDMA	--	--	--	--	--	--	--	--	--	1.4	1.4	1.1	0.9	0.9
Rohypnol	--	--	--	--	--	--	--	--	--	0.7	0.5	0.5	0.4	0.6
GHB	--	--	--	--	--	--	--	--	--	0.4	0.3	0.3	0.3	0.4

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 7th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*
Cigarettes	14.5	16.9	17.7	19.0	16.3	14.0	12.3	11.0	9.9	9.1	8.1	8.9	7.5	6.7	-0.8
Cigars	--	--	--	--	--	6.9	5.8	5.3	6.0	4.6	3.7	4.5	3.7	3.2	-0.5
Pipe	--	--	--	--	--	2.4	1.8	2.1	--	1.5	1.5	1.6	1.2	1.2	
Smokeless Tobacco	6.7	7.5	7.1	6.7	5.2	3.9	2.7	2.2	2.1	2.3	2.1	2.8	2.2	2.3	
Alcohol	21.8	21.5	19.6	21.3	18.4	16.8	16.5	16.1	15.2	14.7	15.3	12.7	12.9		
Marijuana	2.7	4.8	5.4	7.9	5.7	5.5	5.1	5.2	5.2	4.7	4.6	4.5	4.1	3.6	-0.5
Cocaine	0.1	0.8	0.7	1.3	1.0	0.9	0.7	0.7	0.6	0.8	0.9	0.8	0.7	0.8	
Crack	0.1	0.7	0.7	1.2	0.9	0.9	0.8	0.7	0.7	0.7	0.9	0.9	0.8	0.8	
Inhalants	4.7	6.1	5.6	7.0	6.6	4.8	3.7	3.3	3.1	3.0	3.7	4.8	3.7	3.6	
Amphetamines	2.2	2.7	2.4	3.3	2.3	1.7	1.2	1.2	1.1	1.1	1.0	1.1	0.7	0.8	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	0.6	0.6
Ritalin	--	--	--	--	--	1.4	1.0	1.2	0.8	0.8	0.9	0.8	0.8	0.8	
Tranquilizers	3.1	3.7	3.3	5.0	4.1	4.0	3.2	3.3	3.0	3.2	2.9	3.3	2.7	2.8	
Narcotics	1.4	1.4	1.3	2.2	1.7	1.6	1.2	1.2	1.0	1.1	1.0	1.1	0.9	0.9	
Psychedelics [#]	1.1	1.3	1.4	2.2	1.8	1.9	1.5	1.3	--	1.0	0.9	0.9	0.8	0.7	
LSD	--	--	--	--	--	--	--	--	--	0.8	0.8	0.8	0.7	0.5	
Others	--	--	--	--	--	--	--	--	--	0.4	0.4	0.4	0.3	0.4	
Heroin	0.1	0.8	0.6	0.9	0.8	0.5	0.5	0.5	0.6	0.6	0.6	0.5	0.5	0.5	
Steroids	0.1	1.3	0.9	1.0	0.8	0.6	1.0	0.7	0.7	0.8	0.7	0.8	0.6	0.6	
Injected Drugs	--	--	--	--	--	--	--	--	0.5	0.7	0.6	0.6	0.6	0.6	
MDMA	--	--	--	--	--	--	--	--	--	1.0	0.9	0.8	0.6	0.6	
Rohypnol	--	--	--	--	--	--	--	--	--	0.4	0.4	0.4	0.3	0.3	
GHB	--	--	--	--	--	--	--	--	--	0.2	0.2	0.2	0.2	0.3	0.1

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 7th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*
CIGARETTES														
Daily Use	7.3	8.6	8.8	10.4	8.3	6.9	6.0	5.6	4.4	4.2	3.6	4.2	3.3	3.0
1/2+ Pack per Day	3.9	4.4	4.5	5.7	4.3	3.7	2.9	2.8	1.9	2.2	1.7	2.1	1.5	1.4
CIGARS														
--	--	--	--	--	--	1.2	1.0	1.0	1.3	0.9	0.7	1.0	0.7	0.6
SMOKELESS TOBACCO														
	1.2	1.7	1.1	1.3	0.9	0.5	0.4	0.4	0.3	0.6	0.3	0.6	0.4	0.4
ALCOHOL														
Daily Use	1.2	1.4	1.0	1.4	1.1	1.2	1.0	1.1	0.9	1.0	1.0	1.2	0.9	0.8
Binge Drinking	12.7	13.0	12.3	13.5	12.0	12.3	12.4	12.4	9.7	10.0	8.0	8.9	6.8	7.0
MARIJUANA														
	0.1	0.7	0.9	1.4	1.0	1.1	1.0	1.0	1.0	0.9	0.9	1.0	0.9	0.7
														-0.2

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).



Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 1993-2006

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*	National##	2005
Cigarettes	50.0	51.3	52.4	53.7	52.6	49.5	45.7	43.7	39.9	37.4	35.1	34.7	32.1	30.2	-1.9	25.9
Cigars	--	--	--	--	--	24.4	22.5	21.3	22.1	17.4	16.4	16.1	15.8	14.6	-1.2	--
Pipe	--	--	--	--	--	8.0	8.6	7.8	--	5.5	5.5	5.6	5.2	4.9	--	--
Smokeless Tobacco	18.9	20.1	19.2	17.9	16.7	12.4	10.9	10.9	8.2	7.7	7.3	8.3	7.3	8.3	1.0	10.1
Alcohol	60.9	61.7	60.2	59.1	59.4	51.5	53.0	51.0	53.3	52.6	51.7	52.0	49.0	46.5	-2.5	41.0
Marijuana	12.0	16.0	20.2	24.3	23.0	22.1	19.7	19.0	19.5	20.0	19.1	18.6	17.6	15.6	-2.0	16.5
Cocaine	2.0	2.5	2.9	3.6	3.5	3.2	2.6	2.6	2.7	2.6	2.9	2.7	2.9	2.5	-0.4	3.7
Crack	1.5	2.0	2.4	3.0	2.9	2.9	2.4	2.4	2.5	2.3	2.7	2.4	2.6	2.2	-0.4	2.4
Inhalants	14.5	15.3	18.1	17.4	17.6	13.9	12.0	11.2	10.3	9.3	10.7	11.0	11.3	10.4	-0.9	17.1
Amphetamines	9.0	9.1	10.6	10.0	9.1	6.6	5.8	5.6	5.5	5.0	4.7	4.3	3.7	3.2	-0.5	7.4
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	2.0	-0.4
Ritalin	--	--	--	--	--	4.2	3.9	4.2	4.0	3.5	3.6	3.1	3.5	3.1	-0.4	--
Tranquilizers	10.3	10.7	11.4	11.9	12.5	11.5	10.1	10.3	10.2	10.2	10.7	10.2	9.8	8.9	-0.9	4.1
Narcotics	5.3	5.4	5.8	5.5	6.5	5.8	5.2	5.0	4.9	4.8	4.8	4.5	4.5	4.1	--	--
Psychedelics#	3.7	3.7	5.1	6.7	6.3	5.6	5.7	5.4	--	4.2	3.7	3.1	3.3	2.8	-0.5	3.8
LSD	--	--	--	--	--	--	--	--	--	3.8	3.2	2.6	2.7	2.3	-0.4	1.9
Others	--	--	--	--	--	--	--	--	--	1.4	1.4	1.4	1.5	1.3	--	3.3
Heroin	1.4	1.7	1.8	2.2	1.8	1.9	1.3	1.6	1.7	1.4	1.8	1.5	1.6	1.5	--	1.5
Steroids	2.6	2.7	2.6	2.1	2.3	2.5	2.1	2.5	2.2	2.1	2.2	2.2	1.8	1.5	-0.3	1.7
Injected Drugs	--	--	--	--	--	--	--	--	1.5	1.4	1.7	1.5	1.8	1.7	--	--
MDMA	--	--	--	--	--	--	--	--	--	3.4	3.3	2.7	2.5	2.3	--	2.8
Rohypnol	--	--	--	--	--	--	--	--	--	1.1	1.3	1.1	1.2	1.2	--	1.1
GHB	--	--	--	--	--	--	--	--	--	0.7	0.7	0.7	0.7	0.8	--	--

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

Monitoring the Future Study, the University of Michigan, 2005



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 8th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*	National##	2005
	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*	National##	2005
Cigarettes	37.9	39.6	41.4	42.6	41.5	37.3	33.1	30.6	26.3	24.1	22.6	22.2	20.6	19.1	-1.5	--	--
Cigars	--	--	--	--	--	20.0	18.1	16.9	16.6	11.9	11.3	11.6	10.3	10.3	-1.0	--	--
Pipe	--	--	--	--	--	5.8	5.9	5.1	--	3.8	4.0	4.2	3.8	3.5	-0.3	--	--
Smokeless Tobacco	14.9	16.5	15.4	14.2	12.8	9.5	8.3	8.0	6.2	5.8	5.5	6.3	5.6	6.4	0.8	--	--
Alcohol	52.3	52.9	51.6	51.4	51.2	45.1	47.0	45.1	45.4	43.8	42.4	42.1	39.0	38.3	-0.7	33.9	
Marijuana	10.4	14.3	18.4	22.4	21.0	19.8	17.7	17.1	17.1	16.8	16.0	15.4	14.5	12.7	-1.8	12.2	
Cocaine	1.6	2.1	2.5	2.9	3.0	2.7	2.1	2.2	2.2	2.0	2.3	2.1	2.3	1.9	-0.4	2.2	
Crack	1.2	1.6	2.0	2.3	2.4	2.4	1.9	1.9	2.0	1.7	2.0	1.7	1.8	1.7	1.4	1.4	
Inhalants	11.2	12.4	14.5	13.6	14.0	10.5	8.9	8.3	7.4	6.2	7.5	7.9	7.8	7.2	-0.6	9.5	
Amphetamines	7.8	7.8	9.3	8.8	7.9	5.6	4.7	4.7	4.5	3.8	3.7	3.4	2.9	2.5	-0.4	4.9	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	1.7	1.4	-0.3
Ritalin	--	--	--	--	--	3.6	3.2	3.5	3.1	2.5	2.6	2.2	2.6	2.3	-0.3	2.4	
Tranquilizers	8.6	9.1	9.5	10.3	10.8	9.7	8.5	8.5	7.9	8.3	8.1	7.5	6.8	-0.7	2.8		
Narcotics	4.3	4.0	4.8	4.6	5.5	4.9	4.2	4.1	4.1	3.6	3.5	3.3	3.2	3.0	-0.2	--	--
Psychedelics#	3.1	3.1	4.5	6.0	5.6	5.0	4.8	4.5	--	2.9	2.7	2.4	2.4	2.1	-0.3	2.4	
LSD	--	--	--	--	--	--	--	--	--	2.7	2.3	1.9	2.0	1.7	-0.3	1.2	
Others	--	--	--	--	--	--	--	--	--	1.0	1.0	1.1	1.0	1.1	1.2	2.0	
Heroin	1.1	1.5	1.8	1.6	1.5	1.1	1.2	1.3	1.0	1.4	1.2	1.2	1.3	1.2	1.0	0.8	
Steroids	2.0	2.0	1.8	1.6	1.7	2.0	1.5	1.8	1.7	1.6	1.6	1.6	1.4	1.1	-0.3	1.1	
Injected Drugs	--	--	--	--	--	--	--	--	1.1	1.0	1.2	1.2	1.3	1.2	--	--	
MDMA	--	--	--	--	--	--	--	--	--	2.7	2.6	2.0	1.8	1.7	1.7		
Rohypnol	--	--	--	--	--	--	--	--	--	0.8	1.0	0.9	0.9	0.8	0.7		
GHB	--	--	--	--	--	--	--	--	--	0.4	0.5	0.6	0.5	0.5	0.5	0.5	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

Monitoring the Future Study, the University of Michigan, 2005



Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 1993-2006

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*	National##	2005
Cigarettes	21.1	24.2	26.3	27.1	25.8	24.1	19.7	18.2	16.1	14.7	14.0	13.4	12.5	11.6	-0.9	9.3
Cigars	--	--	--	--	--	12.0	9.6	9.7	10.0	6.6	6.8	6.7	6.5	5.9	-0.6	--
Pipe	--	--	--	--	--	3.5	3.5	3.1	--	2.2	2.2	2.3	2.1	2.0	--	--
Smokeless Tobacco	9.0	10.6	9.3	8.6	7.2	5.6	4.6	4.5	3.5	3.4	3.2	3.8	3.3	3.8	0.5	3.3
Alcohol	30.9	32.0	31.3	31.3	30.3	27.2	27.7	26.4	24.7	24.9	24.3	23.5	21.1	20.9	--	17.1
Marijuana	6.5	9.9	13.1	15.2	13.8	12.8	11.3	11.3	11.1	10.6	9.8	9.3	8.2	-1.1	6.6	
Cocaine	0.1	1.3	1.4	1.7	1.9	1.5	1.1	1.1	1.3	1.2	1.3	1.2	1.4	1.1	-0.3	1.0
Crack	0.1	1.0	1.1	1.4	1.4	1.3	1.1	1.0	1.2	0.9	1.2	1.0	1.2	1.0	-0.2	0.6
Inhalants	6.5	7.5	8.4	8.0	8.2	5.7	4.8	4.4	4.0	3.4	4.2	4.7	4.3	4.1	--	4.2
Amphetamines	5.1	5.2	5.9	5.6	5.3	3.3	2.8	2.8	2.8	2.4	2.4	2.1	1.9	1.4	-0.5	2.3
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	0.7
Ritalin	--	--	--	--	--	2.0	1.7	2.0	1.7	1.3	1.4	1.3	1.5	1.3	-0.2	--
Tranquilizers	5.6	5.7	5.8	6.0	6.5	5.7	5.2	5.1	5.2	4.8	5.1	4.9	4.7	4.1	-0.6	1.3
Narcotics	2.6	2.5	2.6	2.7	3.2	2.8	2.4	2.4	2.4	2.0	2.1	1.9	1.9	1.8	--	--
Psychedelics#	2.0	2.1	3.1	3.6	3.6	3.0	2.6	2.7	--	1.6	1.5	1.4	1.5	1.2	-0.3	1.1
LSD	--	--	--	--	--	--	--	--	--	1.5	1.3	1.2	1.2	0.9	-0.3	0.5
Others	--	--	--	--	--	--	--	--	--	0.6	0.7	0.7	0.6	0.6	--	0.9
Heroin	0.1	1.0	0.9	1.1	1.1	1.0	0.6	0.7	0.8	0.7	0.9	0.8	0.8	0.7	--	0.5
Steroids	1.2	1.3	1.0	1.0	1.1	0.9	0.9	1.0	0.9	1.0	0.9	1.0	0.9	0.8	0.7	0.5
Injected Drugs	--	--	--	--	--	--	--	--	0.7	0.6	0.8	0.8	0.8	0.8	--	--
MDMA	--	--	--	--	--	--	--	--	--	1.6	1.5	1.3	1.1	1.0	--	0.6
Rohypnol	--	--	--	--	--	--	--	--	--	0.5	0.7	0.6	0.6	0.5	--	0.2
GHB	--	--	--	--	--	--	--	--	--	0.3	0.4	0.4	0.4	0.4	--	--

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

Monitoring the Future Study, the University of Michigan, 2005



**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 8th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*	National##	2005
	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*	National##	2005
CIGARETTES																
Daily Use	11.8	13.8	14.9	15.4	14.2	13.7	10.6	10.0	8.8	7.5	7.2	6.9	6.7	5.8	-0.9	4.0
1/2+ Pack per Day	6.7	7.2	8.4	8.8	8.4	7.5	5.6	5.3	4.9	4.3	3.7	3.5	3.3	2.9	-0.4	1.7
CIGARS	--	--	--	--	--	2.5	1.7	1.9	3.0	1.4	1.4	1.3	1.3	1.3	--	--
SMOKELESS TOBACCO																
	2.3	2.6	2.1	2.2	1.6	1.4	0.9	0.9	0.9	0.8	0.7	0.9	0.8	1.0	0.7	0.7
ALCOHOL																
Daily Use	1.8	2.3	2.1	2.2	2.4	2.1	2.4	2.2	1.9	2.1	1.9	1.8	1.6	1.6	0.5	0.5
Binge Drinking	18.5	18.6	18.4	19.0	18.9	19.7	19.5	18.6	14.8	15.5	13.4	13.4	11.6	11.0	10.5	10.5
MARIJUANA	0.9	1.9	2.1	3.1	2.7	2.7	2.7	3.1	2.7	2.6	2.3	2.4	2.0	-0.4	1.0	1.0

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).

Monitoring the Future Study, the University of Michigan, 2005



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 9th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*
Cigarettes	56.5	57.4	58.7	61.4	57.3	55.8	53.7	50.6	47.2	43.6	39.8	40.7	36.8	36.3	
Cigars	--	--	--	--	--	31.0	28.7	26.8	29.8	23.1	20.7	21.3	20.6	20.4	
Pipe	--	--	--	--	--	10.9	10.1	9.7	--	7.6	7.1	7.3	6.5	6.9	
Smokeless Tobacco	24.3	25.8	26.2	24.2	21.3	18.7	15.0	14.2	11.9	12.4	10.5	12.2	10.7	11.6	0.9
Alcohol	70.2	69.9	68.7	69.7	65.9	61.8	61.4	59.8	62.6	59.9	58.4	58.4	55.2	55.5	
Marijuana	18.4	23.9	26.1	35.0	28.9	30.7	27.8	26.7	28.4	28.1	25.2	24.0	22.9	22.5	
Cocaine	2.5	3.7	3.7	4.9	4.8	5.3	4.2	3.9	3.8	4.3	4.3	4.7	4.2	4.4	
Crack	1.8	2.5	2.7	4.0	3.5	4.2	3.3	3.0	3.0	3.1	3.2	3.3	3.0	3.0	
Inhalants	13.3	17.4	17.3	18.3	16.7	15.7	12.3	11.6	11.1	9.9	9.8	11.3	10.9	11.4	
Amphetamines	13.5	14.8	14.3	16.7	13.3	12.0	9.9	10.1	10.0	9.7	8.6	8.6	7.0	6.8	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	3.5	3.0
Ritalin	--	--	--	--	--	--	--	--	--	--	--	--	--	3.5	-0.5
Tranquilizers	11.1	13.8	13.4	15.3	15.0	14.9	13.7	14.0	14.2	14.4	13.2	13.6	12.3	12.4	
Narcotics	6.9	8.5	7.9	9.9	9.1	9.2	8.1	8.1	8.5	9.1	7.9	8.0	7.0	7.4	
Psychedelics [#]	5.6	6.4	7.8	12.1	8.9	9.9	9.2	8.0	--	6.9	5.2	5.2	4.7	4.2	-0.5
LSD	--	--	--	--	--	--	--	--	--	6.3	4.5	4.3	3.9	3.4	-0.5
Others	--	--	--	--	--	--	--	--	--	2.4	2.3	2.4	2.3	2.2	
Heroin	1.5	2.0	1.9	2.4	2.1	2.1	1.7	1.6	1.8	1.8	2.0	2.0	2.1	1.9	
Steroids	2.8	3.0	2.6	2.5	2.3	2.5	2.4	2.8	2.5	2.4	2.6	2.7	2.3	1.8	-0.5
Injected Drugs	--	--	--	--	--	--	--	--	1.3	1.6	2.0	2.0	2.0	1.8	
MDMA	--	--	--	--	--	--	--	--	--	5.4	4.2	4.1	3.6	3.7	
Rohypnol	--	--	--	--	--	--	--	--	--	1.6	1.6	1.8	1.8	1.8	
GHB	--	--	--	--	--	--	--	--	--	1.1	1.1	1.1	1.1	1.0	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 9th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*
Cigarettes	43.6	45.5	46.2	50.2	46.4	43.4	39.7	37.1	33.3	29.1	27.0	27.6	25.4	25.0
Cigars	--	--	--	--	--	26.3	23.8	21.8	23.2	16.2	14.6	15.4	15.1	15.3
Pipe	--	--	--	--	--	7.9	6.9	6.6	--	5.1	5.1	5.5	4.9	5.1
Smokeless Tobacco	19.2	20.8	20.8	19.6	16.7	14.1	11.5	11.2	9.0	8.8	8.1	9.3	8.3	9.6
Alcohol	62.5	61.6	60.5	62.2	58.1	56.0	56.0	54.6	54.7	50.4	50.1	49.9	46.3	47.3
Marijuana	16.1	21.7	23.7	31.8	25.8	27.3	24.6	24.0	24.6	23.4	20.9	19.6	18.0	18.4
Cocaine	2.1	2.9	3.1	4.1	4.0	4.5	3.2	3.3	3.1	3.2	3.2	3.5	3.1	3.3
Crack	1.4	2.0	2.2	3.2	2.9	3.3	2.4	2.4	2.3	2.1	2.2	2.4	2.2	2.2
Inhalants	9.8	13.5	13.2	13.8	12.1	11.5	8.8	8.1	7.2	6.1	6.3	7.5	6.9	7.5
Amphetamines	11.6	12.9	12.2	14.4	11.4	9.8	8.2	8.0	8.5	7.2	6.7	6.6	5.2	5.1
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	-0.3
Ritalin	--	--	--	--	--	5.7	3.9	4.8	4.9	3.7	3.6	3.4	3.6	3.8
Tranquilizers	9.2	11.9	10.8	13.0	12.7	12.5	11.7	12.2	12.2	10.8	10.2	10.7	8.9	9.6
Narcotics	5.5	6.9	6.3	7.9	7.6	7.8	6.9	6.8	7.1	6.6	5.8	5.9	4.9	5.3
Psychedelics [#]	4.7	5.6	6.9	10.9	7.7	8.5	7.9	6.7	--	4.7	3.6	3.7	3.1	3.1
LSD	--	--	--	--	--	--	--	--	--	4.4	3.1	3.1	2.5	2.4
Others	--	--	--	--	--	--	--	--	--	1.6	1.6	1.8	1.6	1.6
Heroin	1.1	1.5	1.7	2.0	1.7	1.6	1.3	1.4	1.3	1.1	1.4	1.4	1.4	1.4
Steroids	2.0	2.2	1.8	1.8	1.7	1.8	1.7	2.1	1.9	1.7	2.0	1.9	1.6	1.4
Injected Drugs	--	--	--	--	--	--	--	--	0.9	1.1	1.4	1.4	1.3	1.3
MDMA	--	--	--	--	--	--	--	--	--	3.9	3.1	3.0	2.5	2.7
Rohypnol	--	--	--	--	--	--	--	--	--	1.1	1.2	1.3	1.2	1.3
GHB	--	--	--	--	--	--	--	--	--	0.7	0.8	0.8	0.8	0.8

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 9th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*
Cigarettes	26.2	29.4	30.8	34.4	30.5	28.9	25.4	24.9	21.8	19.1	17.9	18.2	16.4	16.5	
Cigars	--	--	--	--	--	15.5	13.2	12.6	14.1	9.2	8.3	9.1	8.6	8.8	
Pipe	--	--	--	--	--	4.5	4.1	4.2	--	2.9	2.7	3.2	2.7	3.0	
Smokeless Tobacco	12.2	13.3	13.2	12.5	9.8	8.3	6.5	6.5	5.4	5.2	4.9	5.6	5.2	5.7	0.5
Alcohol	38.2	39.6	38.5	40.8	36.0	36.7	36.2	35.3	33.2	30.5	30.8	29.4	26.6	28.5	1.9
Marijuana	10.6	15.2	15.9	22.8	17.2	18.4	16.2	15.6	16.1	14.6	13.9	12.1	11.5	11.4	
Cocaine	0.1	1.6	1.7	2.4	2.2	2.4	1.4	1.6	1.6	1.6	1.8	2.0	1.7	1.7	
Crack	0.1	1.4	1.3	2.1	1.5	1.9	1.2	1.3	1.2	1.1	1.2	1.4	1.3	1.1	
Inhalants	5.2	7.3	6.9	7.2	6.1	5.8	4.4	4.2	3.7	3.0	3.0	3.9	3.6	3.6	
Amphetamines	6.4	7.8	7.2	8.9	6.8	5.9	4.8	5.1	5.2	4.3	3.6	3.8	3.0	3.0	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	-0.3
Ritalin	--	--	--	--	--	3.0	2.0	2.5	2.4	1.9	1.8	1.9	2.0	2.0	
Tranquilizers	5.2	7.1	6.2	7.7	7.9	7.5	7.2	7.4	7.3	6.7	5.9	6.3	5.1	5.7	
Narcotics	2.9	3.7	3.8	4.9	4.2	3.8	3.9	4.2	3.9	3.9	3.3	3.1	2.8	2.9	
Psychedelics [#]	2.7	3.7	4.9	6.5	4.6	5.0	4.3	3.6	--	2.4	1.8	1.9	1.7	1.6	
LSD	--	--	--	--	--	--	--	--	--	2.0	1.5	1.6	1.4	1.3	
Others	--	--	--	--	--	--	--	--	--	1.0	0.8	1.0	0.9	0.8	
Heroin	0.1	1.1	0.9	1.3	1.0	0.9	0.6	0.7	0.8	0.7	0.9	0.9	0.9	0.8	
Steroids	1.3	1.4	1.2	1.2	1.1	1.1	0.9	1.0	1.1	1.0	1.2	1.1	0.9	0.9	
Injected Drugs	--	--	--	--	--	--	--	--	--	0.6	0.6	0.8	0.8	0.8	
MDMA	--	--	--	--	--	--	--	--	--	--	1.9	1.5	1.6	1.3	1.5
Rohypnol	--	--	--	--	--	--	--	--	--	--	0.7	0.6	0.7	0.7	0.8
GHB	--	--	--	--	--	--	--	--	--	--	0.5	0.5	0.5	0.5	0.5

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 9th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*
CIGARETTES														
Daily Use	16.0	18.3	18.4	22.0	19.0	18.5	15.5	14.7	12.4	11.6	10.3	10.3	9.4	9.6
1/2+ Pack per Day	9.9	11.7	11.2	14.0	11.8	11.6	9.2	8.4	7.0	7.0	6.1	5.7	5.1	5.4
CIGARS														
--	--	--	--	--	--	3.3	2.8	2.8	4.3	2.1	1.7	2.2	1.8	1.9
SMOKELESS TOBACCO														
3.8	4.2	4.1	3.9	2.5	2.4	1.8	2.0	1.5	1.6	1.4	1.8	1.5	1.8	0.3
ALCOHOL														
Daily Use	2.8	3.2	3.1	3.6	3.1	3.6	3.6	3.4	3.0	3.1	2.8	2.9	2.4	2.7
Binge Drinking	23.3	23.6	23.3	25.5	23.1	25.7	24.3	24.3	19.7	19.8	17.9	17.3	14.8	15.6
MARIJUANA														
1.8	3.3	3.7	6.4	4.1	4.9	4.6	4.5	4.2	4.5	3.8	3.2	3.3	3.2	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 10th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*	National##	2005
Cigarettes	60.8	61.5	61.8	63.6	64.1	61.4	60.5	57.4	55.5	51.5	47.2	46.7	43.9	-0.9	38.9	38.9
Cigars	--	--	--	--	--	37.5	36.4	32.4	36.4	30.6	27.2	27.8	26.9	26.8	--	--
Pipe	--	--	--	--	--	12.8	12.1	10.6	--	9.0	8.3	7.7	7.8	8.6	0.8	--
Smokeless Tobacco	27.9	28.0	28.7	30.0	26.2	22.1	19.6	17.3	16.0	16.2	13.8	14.1	14.3	15.4	1.1	14.5
Alcohol	76.5	76.2	73.2	74.3	75.2	68.4	69.2	66.6	71.3	68.4	67.2	65.5	63.5	62.3	-1.2	63.2
Marijuana	25.6	30.5	34.2	40.2	40.1	38.2	37.4	34.1	37.8	36.9	34.8	33.5	31.6	30.1	-1.5	34.1
Cocaine	3.4	4.3	4.9	6.9	6.7	6.6	6.0	5.1	6.1	6.5	6.0	6.0	6.1	5.9	5.2	5.2
Crack	1.9	2.9	3.6	4.8	4.7	5.1	4.1	3.4	3.9	3.5	3.7	3.8	3.8	3.4	-0.4	2.5
Inhalants	15.0	16.1	17.5	17.8	17.5	15.9	12.8	11.4	11.9	10.7	10.9	11.3	11.2	11.1	13.1	13.1
Amphetamines	17.3	17.2	17.7	18.8	17.6	15.1	13.5	12.2	13.7	13.2	12.3	11.3	10.2	9.0	-1.2	11.1
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	4.5	3.5
Ritalin	--	--	--	--	--	6.8	6.6	6.8	7.5	7.5	6.5	6.1	6.5	6.5	6.5	4.1
Tranquilizers	13.0	14.6	13.5	16.3	16.3	16.1	15.5	14.9	16.4	17.4	16.5	16.0	15.0	14.3	-0.7	7.1
Narcotics	9.0	9.8	9.7	11.7	12.6	11.9	11.1	10.1	11.7	12.2	11.6	11.0	9.9	9.8	--	--
Psychedelics#	7.7	8.3	11.5	14.1	13.2	12.9	12.6	10.9	--	9.4	7.0	6.1	5.7	5.3	5.8	5.8
LSD	--	--	--	--	--	--	--	--	--	8.6	6.0	5.1	4.7	4.3	2.5	2.5
Others	--	--	--	--	--	--	--	--	--	3.2	3.1	2.9	2.9	2.7	5.2	5.2
Heroin	1.4	2.0	1.9	2.8	2.2	2.5	2.0	1.9	2.1	2.2	2.2	2.1	2.1	2.0	1.5	1.5
Steroids	2.4	2.8	2.2	2.9	2.4	3.1	2.3	2.7	2.5	3.0	2.8	2.7	2.2	2.2	2.0	2.0
Injected Drugs	--	--	--	--	--	--	--	--	1.6	1.8	1.9	1.9	2.1	2.0	--	--
MDMA	--	--	--	--	--	--	--	--	--	6.9	5.7	4.7	4.6	4.4	4.0	4.0
Rohypnol	--	--	--	--	--	--	--	--	--	1.8	1.9	1.8	1.8	1.9	1.0	1.0
GHB	--	--	--	--	--	--	--	--	--	1.3	1.1	1.2	1.2	1.1	--	--

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

Monitoring the Future Study, the University of Michigan, 2005



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 10th Grade Students: 1993-2006**

DRUG	National##						2005
	1993	1994	1995	1996	1997	1998	
Cigarettes	46.8	48.0	48.9	50.8	52.4	47.9	45.5
Cigars	--	--	--	--	--	32.1	30.4
Pipe	--	--	--	--	--	8.9	8.3
Smokeless Tobacco	21.6	21.8	22.2	22.9	20.3	16.4	14.6
Alcohol	68.0	67.7	65.3	65.9	67.3	61.8	63.0
Marijuana	22.2	27.6	30.8	36.1	35.1	33.7	32.8
Cocaine	2.6	3.5	4.0	5.7	5.5	5.4	5.0
Crack	1.5	2.3	3.0	4.0	3.8	4.0	3.2
Inhalants	10.3	11.3	12.4	11.3	11.7	10.7	8.1
Amphetamines	13.9	14.4	14.4	15.4	14.5	12.1	10.5
Methamphetamine	--	--	--	--	--	--	--
Ritalin	--	--	--	--	--	5.2	4.9
Tranquilizers	10.4	12.1	11.2	13.5	13.5	12.8	12.1
Narcotics	6.9	7.8	7.6	9.3	10.2	9.8	9.0
Psychedelics#	6.4	7.1	10.2	12.3	11.4	10.5	8.6
LSD	--	--	--	--	--	--	--
Others	--	--	--	--	--	--	--
Heroin	1.2	1.6	2.3	1.8	1.8	1.6	1.2
Steroids	1.8	2.1	1.7	2.2	1.7	2.2	1.6
Injected Drugs	--	--	--	--	--	--	--
MDMA	--	--	--	--	--	--	--
Rohypnol	--	--	--	--	--	--	--
GHB	--	--	--	--	--	--	--

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

Monitoring the Future Study, the University of Michigan, 2005



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 10th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*	National##	2005		
Cigarettes	30.4	33.0	34.4	36.7	37.2	33.9	31.5	28.9	27.0	23.8	22.2	22.1	20.8	20.6	14.9			
Cigars	--	--	--	--	--	19.0	16.5	14.6	17.8	11.9	11.5	12.4	11.8	11.6	--			
Pipe	--	--	--	--	--	4.8	4.3	3.6	--	3.2	3.3	2.9	3.2	3.4	--			
Smokeless Tobacco	13.2	13.6	13.8	14.6	12.3	9.5	8.2	7.4	6.9	7.1	6.0	6.4	7.0	7.8	0.8	5.6		
Alcohol	43.8	44.9	42.7	43.7	43.9	41.3	41.6	39.7	39.7	38.1	36.9	34.1	33.0	33.0	33.2	33.2		
Marijuana	14.3	18.9	21.5	24.9	23.7	21.8	21.1	19.0	21.4	19.2	18.2	17.2	16.0	14.6	-1.4	15.2		
Cocaine	1.4	1.6	2.2	2.6	2.6	2.9	2.3	1.7	2.4	2.4	2.3	2.4	2.2	2.1	1.5	1.5		
Crack	0.1	1.2	1.6	1.8	1.9	2.1	1.5	1.1	1.5	1.1	1.4	1.4	1.3	1.2	0.7	0.7		
Inhalants	5.0	5.6	5.8	5.1	5.3	4.9	3.4	2.9	3.3	2.9	2.9	3.3	3.0	3.0	2.2	2.2		
Amphetamines	7.5	8.3	7.9	7.8	7.8	6.5	5.6	5.4	6.5	5.9	5.2	4.8	4.0	3.7	3.7	3.7		
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	1.6	1.2	-0.4	1.1
Ritalin	--	--	--	--	--	2.5	2.1	2.4	2.6	2.2	2.1	2.0	2.1	2.2	--	--	--	--
Tranquilizers	5.7	6.9	6.2	7.0	7.4	7.5	7.1	7.0	8.0	7.6	7.1	7.0	6.4	5.9	-0.5	2.3	2.3	
Narcotics	3.4	4.0	4.1	4.4	5.0	5.2	4.4	4.4	5.5	4.6	4.2	4.3	3.6	3.8	--	--	--	--
Psychedelics#	3.7	4.1	7.0	6.8	6.5	6.0	5.5	4.2	--	2.5	2.1	2.0	1.8	1.9	1.5	1.5		
LSD	--	--	--	--	--	--	--	--	--	2.2	1.7	1.5	1.4	1.5	0.6	0.6	0.6	
Others	--	--	--	--	--	--	--	--	--	1.1	1.1	1.0	1.0	1.0	1.3	1.3	1.3	
Heroin	0.1	0.9	1.0	1.3	1.0	1.3	0.9	0.7	1.0	0.8	0.9	0.9	0.9	0.8	0.5	0.5	0.5	
Steroids	1.0	1.3	1.1	1.4	0.9	1.4	0.9	0.9	1.0	1.1	1.0	1.1	0.9	1.1	0.6	0.6	0.6	
Injected Drugs	--	--	--	--	--	--	--	--	0.8	0.7	0.7	0.8	0.9	0.9	--	--	--	
MDMA	--	--	--	--	--	--	--	--	--	2.4	1.8	1.5	1.5	1.5	1.0	1.0	1.0	
Rohypnol	--	--	--	--	--	--	--	--	--	0.6	0.7	0.7	0.7	0.8	0.2	0.2	0.2	
GHB	--	--	--	--	--	--	--	--	--	0.5	0.5	0.6	0.5	0.5	--	--	--	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

Monitoring the Future Study, the University of Michigan, 2005



**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 10th Grade Students: 1993-2006**

DRUG								National##								
	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*	2005
CIGARETTES																
Daily Use	19.0	21.9	22.4	24.4	23.9	21.5	20.6	17.9	16.5	14.6	13.7	13.3	12.5	12.2	7.5	
1/2+ Pack per Day	12.3	14.4	14.6	16.2	15.5	13.4	12.9	10.8	9.8	9.4	8.1	7.6	7.1	7.1	3.1	
CIGARS																
--	--	--	--	--	--	3.7	2.9	2.9	5.7	2.7	2.4	2.4	2.5	2.5	2.5	--
SMOKELESS TOBACCO																
4.7	5.2	5.0	5.6	4.1	3.3	2.5	2.3	2.2	2.7	2.7	2.2	2.4	2.5	3.2	0.7	1.9
ALCOHOL																
Daily Use	3.5	3.6	3.5	4.1	3.9	4.2	4.5	4.1	4.2	4.0	4.0	3.5	3.6	3.6	3.6	1.3
Binge Drinking	27.3	26.8	26.1	27.3	27.8	29.0	28.8	27.1	24.2	24.9	21.8	20.2	19.3	19.9	21.0	
MARIJUANA																
2.7	4.3	5.1	7.4	6.9	6.3	6.5	5.7	6.3	6.3	5.6	5.4	5.0	4.5	-0.5	3.1	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).

Monitoring the Future Study, the University of Michigan, 2005



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 11th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*
Cigarettes	64.7	64.5	66.5	67.7	66.9	66.9	65.3	62.3	58.7	54.8	51.2	50.1	46.9	46.8	
Cigars	--	--	--	--	--	43.9	42.1	39.3	40.7	34.2	31.5	31.8	30.9	30.7	
Pipe	--	--	--	--	--	14.5	13.1	12.4	--	9.4	8.8	9.3	8.1	9.2	1.1
Smokeless Tobacco	31.9	31.0	32.2	31.9	29.1	25.6	23.0	22.9	17.4	17.6	16.3	18.8	15.8	17.2	1.4
Alcohol	82.2	79.7	78.5	78.1	77.5	74.9	73.3	71.9	74.3	72.9	70.2	68.7	66.6	66.7	
Marijuana	31.1	33.2	36.6	44.1	42.3	43.7	42.1	39.8	41.0	41.1	38.4	35.9	34.3	34.4	
Cocaine	4.2	4.9	5.2	7.5	7.8	8.1	7.6	7.5	7.1	7.8	8.0	7.8	7.1	7.8	
Crack	2.0	2.7	3.2	4.1	4.5	5.3	4.2	3.8	3.8	4.0	3.8	4.6	3.7	3.8	
Inhalants	14.0	15.7	16.0	17.6	15.4	16.2	12.3	13.0	11.3	10.5	9.9	11.1	9.8	10.1	
Amphetamines	18.8	18.5	18.2	18.3	18.5	16.8	15.7	15.5	15.1	14.8	14.3	13.6	11.1	10.8	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	4.2
Ritalin	--	--	--	--	--	7.5	7.3	8.0	7.3	7.6	7.0	5.9	6.6	6.7	
Tranquilizers	12.9	13.5	13.9	16.6	16.3	17.5	16.0	16.7	17.0	18.6	16.7	15.9	14.6	14.5	
Narcotics	9.7	11.3	10.8	13.2	13.9	13.7	13.1	13.3	12.8	14.1	13.1	12.6	11.1	11.3	
Psychedelics [#]	10.5	10.1	12.7	16.8	14.9	16.4	15.6	14.0	--	11.4	9.0	7.5	6.3	6.3	
LSD	--	--	--	--	--	--	--	--	--	10.7	7.9	6.4	4.9	4.9	
Others	--	--	--	--	--	--	--	--	--	3.6	3.9	3.7	3.8	3.8	
Heroin	1.2	1.5	1.7	1.9	1.9	2.4	2.0	2.5	2.1	2.1	2.4	2.1	2.3		
Steroids	2.0	2.6	2.5	2.3	2.4	2.7	2.2	2.7	2.3	2.5	2.5	2.3	2.1	1.9	
Injected Drugs	--	--	--	--	--	--	--	--	1.5	1.5	1.8	1.9	2.1	2.1	
MDMA	--	--	--	--	--	--	--	--	--	8.1	6.5	5.7	4.9	5.4	
Rohypnol	--	--	--	--	--	--	--	--	--	1.5	1.5	1.9	1.8	2.0	
GHB	--	--	--	--	--	--	--	--	--	1.4	1.2	1.5	1.2	1.2	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 11th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*
Cigarettes	51.0	50.5	53.3	54.8	54.1	53.5	51.0	47.1	42.5	39.0	35.4	35.1	33.1	33.7
Cigars	--	--	--	--	--	37.2	35.1	30.9	31.7	24.5	22.9	23.6	22.9	23.1
Pipe	--	--	--	--	--	9.6	8.7	7.5	--	6.3	6.1	6.5	5.5	6.6
Smokeless Tobacco	23.5	23.1	23.9	22.9	21.1	18.2	15.9	16.3	12.4	12.5	11.9	13.1	11.8	13.1
Alcohol	73.6	70.2	70.9	70.0	70.0	68.8	67.2	65.6	65.8	63.5	59.8	58.6	56.0	57.0
Marijuana	26.3	28.7	32.8	38.6	36.3	37.5	35.2	32.9	34.1	33.7	29.9	27.5	25.7	26.5
Cocaine	2.9	3.6	4.3	5.8	6.3	6.7	5.9	5.5	5.4	5.7	5.8	5.7	4.8	5.3
Crack	1.6	1.9	2.5	3.2	3.6	4.2	2.9	2.3	2.6	2.9	2.5	3.0	2.3	2.4
Inhalants	8.6	10.1	9.9	10.1	9.4	9.6	6.9	7.1	6.0	5.4	5.1	6.0	5.0	5.2
Amphetamines	14.5	13.9	14.2	13.9	14.1	12.5	11.6	12.2	11.9	10.8	10.2	9.6	7.2	7.6
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	3.0
Ritalin	--	--	--	--	--	5.3	4.8	5.4	4.7	4.4	4.3	3.4	3.9	4.2
Tranquilizers	9.8	10.5	11.0	13.2	13.1	13.6	12.7	13.2	13.6	13.8	11.9	11.9	9.9	10.0
Narcotics	7.0	8.0	8.1	10.3	10.8	11.0	10.4	10.4	10.0	10.0	8.9	9.0	7.2	7.6
Psychedelics [#]	8.6	8.2	11.4	14.1	12.6	13.0	12.5	10.9	--	6.6	5.2	4.8	4.1	4.2
LSD	--	--	--	--	--	--	--	--	--	6.1	4.0	3.8	2.9	3.1
Others	--	--	--	--	--	--	--	--	--	2.4	2.7	2.4	2.5	2.6
Heroin	0.1	1.1	1.3	1.5	1.6	1.8	1.5	1.7	1.5	1.2	1.4	1.5	1.3	1.6
Steroids	1.4	1.8	1.6	1.7	1.5	1.7	1.5	1.8	1.6	1.6	1.7	1.6	1.5	1.3
Injected Drugs	--	--	--	--	--	--	--	--	1.1	1.0	1.2	1.1	1.3	1.3
MDMA	--	--	--	--	--	--	--	--	--	5.6	4.2	3.7	2.9	3.7
Rohypnol	--	--	--	--	--	--	--	--	--	0.9	1.0	1.2	1.0	1.2
GHB	--	--	--	--	--	--	--	--	--	0.9	0.8	1.0	0.7	0.8

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 11th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*
Cigarettes	33.7	34.4	39.3	40.1	39.2	39.8	37.4	34.1	30.2	27.1	25.2	25.1	23.5	23.8	
Cigars	--	--	--	--	--	21.0	18.6	16.8	19.8	13.4	13.1	13.9	13.0	12.8	
Pipe	--	--	--	--	--	4.5	4.3	3.8	--	3.2	3.2	3.0	3.5	0.5	
Smokeless Tobacco	14.4	14.6	15.4	14.2	12.0	8.9	9.7	7.0	7.5	7.2	8.1	7.3	8.0		
Alcohol	49.4	47.0	47.7	48.2	45.8	47.8	45.6	43.2	42.1	41.4	38.1	36.5	34.5	35.8	
Marijuana	16.7	19.6	21.8	25.7	22.3	24.6	22.0	20.5	21.7	21.2	18.4	17.2	15.3	15.7	
Cocaine	1.2	1.6	2.0	3.0	2.9	3.3	2.4	2.2	2.6	2.7	2.5	3.1	2.1	2.7	0.6
Crack	0.1	1.0	1.3	1.6	1.6	1.9	1.0	1.0	1.2	1.1	1.1	1.8	1.1	1.3	
Inhalants	3.8	4.2	4.3	4.3	3.6	3.9	2.7	2.8	2.3	2.1	2.2	2.7	2.1	2.3	
Amphetamines	7.2	7.0	7.0	6.4	7.0	6.0	5.8	6.6	6.4	5.6	5.3	5.4	3.5	3.9	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	1.5	1.5
Ritalin	--	--	--	--	--	2.4	1.9	2.2	2.0	2.0	1.8	1.6	1.8	2.0	
Tranquilizers	4.6	5.3	5.7	6.9	7.0	7.1	6.9	7.3	7.7	7.6	6.7	6.5	5.1	5.6	
Narcotics	3.4	3.5	3.8	4.8	4.7	5.2	4.7	5.3	5.0	4.8	4.4	4.7	3.6	4.2	
Psychedelics [#]	4.5	4.1	7.6	6.4	6.9	6.0	4.9	--	2.5	2.3	2.1	1.8	2.1		
LSD	--	--	--	--	--	--	--	--	--	2.1	1.7	1.7	1.3	1.6	
Others	--	--	--	--	--	--	--	--	--	1.1	1.3	1.1	1.1	1.3	
Heroin	0.1	0.6	0.8	0.8	1.0	0.7	0.9	0.7	0.6	0.8	1.1	0.8	1.0		
Steroids	0.1	1.1	1.0	0.8	0.8	1.0	0.6	0.9	0.7	0.8	1.0	0.9	0.8		
Injected Drugs	--	--	--	--	--	--	--	--	0.7	0.6	0.7	0.8	0.8	0.8	
MDMA	--	--	--	--	--	--	--	--	--	2.3	1.7	1.8	1.3	1.8	0.5
Rohypnol	--	--	--	--	--	--	--	--	--	0.6	0.6	0.7	0.5	0.7	
GHB	--	--	--	--	--	--	--	--	--	0.4	0.5	0.6	0.5	0.6	

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.



**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 11th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*
CIGARETTES														
Daily Use	22.0	22.8	26.0	27.3	26.6	27.0	25.2	23.3	20.0	17.9	16.0	16.4	14.8	14.8
1/2+ Pack per Day	15.0	15.3	17.2	18.9	18.1	17.3	16.2	15.0	12.8	11.9	9.9	10.3	8.8	8.5
CIGARS														
--	--	--	--	--	3.5	3.4	3.3	6.8	2.8	2.8	2.6	2.7	2.6	2.6
SMOKELESS TOBACCO														
5.8	6.4	6.3	6.2	4.7	4.1	3.5	3.9	2.9	3.3	2.7	3.2	2.9	3.4	0.5
ALCOHOL														
Daily Use	4.1	4.5	5.0	5.1	4.5	5.6	4.9	4.9	4.3	4.7	4.2	4.3	3.6	3.5
Binge Drinking	30.6	28.9	29.6	30.1	29.8	32.3	31.3	30.0	25.7	26.5	24.0	23.3	21.2	21.8
MARIJUANA														
3.9	5.1	6.1	8.6	7.1	7.6	7.5	6.9	7.0	7.4	6.2	6.2	5.3	4.9	-0.4

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 12th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*	National##	2005	
Cigarettes	66.4	65.6	67.9	66.4	68.6	67.2	68.1	65.8	62.9	60.5	55.9	53.9	51.9	50.9	50.0	50.0	
Cigars	--	--	--	--	--	50.1	48.9	45.3	46.7	41.2	39.4	37.3	38.9	37.7	--	--	
Pipe	--	--	--	--	--	15.9	15.6	14.0	--	10.9	9.7	10.0	10.5	10.7	--	--	
Smokeless Tobacco	34.0	33.9	35.8	31.3	31.4	29.4	26.9	26.3	22.9	21.2	18.7	19.1	18.8	20.4	1.6	17.5	
Alcohol	85.0	82.4	81.4	80.6	80.8	76.1	78.5	75.9	78.6	77.3	74.8	72.4	72.3	70.2	-2.1	75.1	
Marijuana	35.3	38.1	40.0	43.4	45.6	46.5	45.7	44.4	45.5	44.8	42.3	40.5	40.1	37.1	-3.0	44.8	
Cocaine	5.0	6.1	7.1	8.9	9.3	8.4	8.7	9.0	8.6	8.7	8.8	8.8	8.8	8.6	8.0	8.0	
Crack	2.0	2.8	4.3	4.1	5.6	5.6	4.5	4.2	4.1	4.1	4.0	4.4	3.9	4.0	3.5	3.5	
Inhalants	14.0	15.1	15.5	13.7	15.5	15.3	12.8	12.3	11.6	10.3	9.6	9.5	9.6	10.1	11.4		
Amphetamines	19.8	18.8	18.0	16.5	19.2	16.6	15.6	15.4	15.5	14.9	14.4	13.3	12.5	11.3	-1.2	13.1	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	5.5	5.0	4.5
Ritalin	--	--	--	--	--	5.9	7.4	7.4	7.8	7.2	6.9	6.3	7.1	7.0	--	--	--
Tranquilizers	12.6	13.2	14.2	16.2	16.4	15.8	15.7	16.5	17.3	16.6	15.6	15.5	14.5	-1.0	9.9		
Narcotics	11.3	11.5	11.4	11.7	14.5	14.7	13.6	13.8	14.7	14.5	13.4	12.6	12.5	12.0	--	--	
Psychedelics#	11.6	12.0	14.4	15.5	17.2	17.8	17.1	15.4	--	12.8	10.6	8.1	7.3	7.0	8.8		
LSD	--	--	--	--	--	--	--	--	--	11.9	9.3	6.6	5.5	5.3	3.5		
Others	--	--	--	--	--	--	--	--	--	4.9	4.6	4.3	4.4	4.6	8.1		
Heroin	1.3	1.8	2.0	2.1	2.3	2.7	2.1	2.3	2.3	1.9	2.1	2.3	2.0	2.3	1.5		
Steroids	2.4	2.8	2.6	2.1	2.5	2.6	2.5	2.5	2.3	2.4	2.3	2.3	2.1	2.3	2.6		
Injected Drugs	--	--	--	--	--	--	--	--	1.8	1.4	1.7	1.8	1.7	2.2	0.5	--	
MDMA	--	--	--	--	--	--	--	--	--	9.5	7.4	6.4	5.9	5.6	5.4		
Rohypnol	--	--	--	--	--	--	--	--	--	1.4	1.5	1.6	1.4	1.7	--		
GHB	--	--	--	--	--	--	--	--	--	1.6	1.4	1.2	1.1	1.3	--		

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

Monitoring the Future Study, the University of Michigan, 2005



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 12th Grade Students: 1993-2006**

DRUG	National##						2005
	1993	1994	1995	1996	1997	1998	
Cigarettes	52.2	51.8	54.5	52.7	55.7	53.4	54.6
Cigars	--	--	--	--	--	42.9	40.4
Pipe	--	--	--	--	--	11.1	10.2
Smokeless Tobacco	23.8	24.6	25.9	21.6	21.4	20.2	18.2
Alcohol	76.1	73.3	72.9	72.1	73.1	70.9	72.4
Marijuana	28.3	31.9	34.6	37.4	37.1	38.8	37.1
Cocaine	3.3	4.1	5.6	5.7	6.9	7.1	6.2
Crack	1.3	2.0	3.4	3.0	4.2	4.1	2.9
Inhalants	7.1	8.9	9.0	7.4	8.3	8.1	6.3
Amphetamines	13.8	13.5	13.1	12.1	13.6	11.9	10.8
Methamphetamine	--	--	--	--	--	--	--
Ritalin	--	--	--	--	--	4.0	4.3
Tranquilizers	9.3	9.7	9.5	11.1	12.3	12.8	11.9
Narcotics	7.3	7.8	8.1	8.5	10.7	11.2	9.9
Psychedelics#	9.1	9.3	12.2	12.5	13.6	13.6	12.6
LSD	--	--	--	--	--	--	--
Others	--	--	--	--	--	--	--
Heroin	0.1	1.2	1.5	1.8	1.7	1.9	1.4
Steroids	1.5	1.8	1.6	1.5	1.7	1.9	1.7
Injected Drugs	--	--	--	--	--	--	--
MDMA	--	--	--	--	--	--	--
Rohypnol	--	--	--	--	--	--	--
GHB	--	--	--	--	--	--	--

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

Monitoring the Future Study, the University of Michigan, 2005



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 12th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 Change*	National##	2005
Cigarettes	35.6	37.3	40.6	39.8	41.6	41.6	40.5	38.9	35.1	30.5	28.8	27.4	26.5	26.9	23.2	
Cigars	--	--	--	--	--	25.8	23.0	20.2	23.9	16.8	17.0	16.8	17.6	16.6	-1.0	--
Pipe	--	--	--	--	--	4.8	5.2	4.4	--	3.3	3.3	3.4	3.7	4.6	0.9	--
Smokeless Tobacco	15.4	15.5	16.5	14.4	13.5	12.1	11.2	11.3	9.4	8.3	7.9	8.7	8.6	9.8	1.2	7.6
Alcohol	53.2	53.7	51.2	52.5	50.1	52.0	51.7	50.1	49.9	48.1	46.1	42.2	41.8	42.2	47.0	
Marijuana	17.8	21.6	24.0	25.4	23.5	24.5	23.5	22.7	23.5	20.5	19.8	18.3	17.8	17.2	19.8	
Cocaine	1.5	2.0	2.5	3.0	3.3	3.5	2.4	2.7	2.8	2.9	2.5	2.9	2.9	3.1	2.3	
Crack	0.1	0.9	1.7	1.5	1.8	2.0	1.1	1.1	1.1	1.1	1.1	1.4	1.2	1.4	1.0	
Inhalants	2.9	3.8	3.8	3.1	3.3	3.2	2.6	2.4	2.0	1.9	1.7	1.8	1.9	2.2	2.0	
Amphetamines	6.5	7.2	6.5	6.1	6.7	5.8	5.1	5.9	5.7	5.3	4.9	4.5	3.8	4.1	3.9	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	1.7	1.5
Ritalin	--	--	--	--	--	1.7	1.5	1.9	2.0	1.4	1.5	1.4	1.7	1.8	--	--
Tranquilizers	4.5	4.6	4.9	6.0	6.2	6.6	5.9	6.5	6.7	6.0	5.9	5.4	5.2	5.3	2.9	
Narcotics	3.1	3.6	4.0	4.3	4.5	4.6	4.5	4.5	4.5	4.6	4.0	4.0	3.8	4.0	--	--
Psychedelics#	4.5	4.4	8.2	6.0	7.2	7.0	5.7	4.5	--	2.0	2.2	2.0	1.8	2.1	1.9	
LSD	--	--	--	--	--	--	--	--	--	1.5	1.5	1.4	1.2	1.3	0.7	
Others	--	--	--	--	--	--	--	--	--	1.1	1.3	1.2	1.5	0.3	1.6	
Heroin	0.1	0.7	0.9	1.1	1.0	1.1	0.7	0.9	0.8	0.6	0.6	0.8	0.7	1.0	0.3	0.5
Steroids	0.1	0.9	0.8	0.8	1.0	1.1	0.9	0.6	0.7	0.6	0.7	0.8	0.8	1.0	0.9	
Injected Drugs	--	--	--	--	--	--	--	--	0.6	0.6	0.6	0.7	0.6	0.9	0.3	--
MDMA	--	--	--	--	--	--	--	--	--	2.1	1.6	1.4	1.3	1.7	0.4	1.0
Rohypnol	--	--	--	--	--	--	--	--	--	0.5	0.4	0.5	0.5	0.7	--	--
GHB	--	--	--	--	--	--	--	--	--	0.4	0.4	0.4	0.4	0.5	--	--

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p<.05$).

Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

Monitoring the Future Study, the University of Michigan, 2005



**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 12th Grade Students: 1993-2006**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*	National##	2005
	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	Change*	National##	2005
CIGARETTES																	
Daily Use	23.2	25.1	27.6	27.0	28.6	28.5	27.6	26.5	23.3	19.9	18.7	18.0	16.4	16.9	13.6		
1/2+ Pack per Day	16.0	17.9	18.9	18.6	19.3	20.0	18.1	17.0	15.3	13.3	11.9	10.9	9.9	10.5	6.9		
CIGARS																	
--	--	--	--	--	--	3.8	3.7	3.4	7.5	2.8	2.9	3.0	3.1	3.1	--	--	
SMOKELESS TOBACCO																	
6.9	7.0	7.6	6.7	6.1	5.3	5.0	4.9	4.1	3.8	3.4	4.0	3.5	4.4	0.9	2.5	--	
ALCOHOL																	
Daily Use	6.0	6.9	6.4	7.5	5.9	8.1	7.1	6.8	6.2	6.5	6.0	5.6	5.4	5.6	3.1		
Binge Drinking	34.3	34.8	33.3	35.6	33.2	37.5	36.3	35.4	31.9	31.5	29.3	27.0	25.9	27.3	1.4	28.1	
MARIJUANA																	
4.2	6.0	7.4	7.6	8.1	8.5	8.2	7.5	8.2	7.3	7.4	6.7	6.3	6.1	5.0			

NOTES: -- Data not available

* Statistically significant changes between 2005 and 2006 prevalence rates ($p < .05$).

Monitoring the Future Study, the University of Michigan, 2005



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	0.7	0.4	0.5	0.5	0.4	0.4	0.3
	Never	87.4	80.0	69.3	63.3	56.6	52.8	48.7
	Once or Twice	8.5	11.8	15.4	14.9	16.7	16.7	17.8
	Occasionally	1.6	3.7	6.6	8.8	11.2	12.2	13.9
	Past Regularly	1.1	2.1	3.5	4.6	5.1	5.3	5.1
	Current Regularly	0.8	2.1	4.6	7.9	10.0	12.6	14.2
Cigars	No Answer	0.2	0.3	0.4	0.4	0.3	0.4	0.4
	Never	94.4	91.4	85.0	79.2	72.9	68.8	61.9
	1-5 times	4.2	6.1	9.8	11.9	15.1	16.7	19.1
	6-19 times	0.5	0.9	2.0	3.5	4.7	5.9	7.5
	20-40 times	0.2	0.5	0.9	1.8	2.8	3.3	4.7
	40+ times	0.5	0.8	1.9	3.2	4.1	4.9	6.4
Pipe	No Answer	0.6	0.8	1.2	1.4	1.5	2.0	2.0
	Never	97.7	96.2	93.9	91.7	89.9	88.8	87.3
	1-5 times	1.3	2.1	3.2	4.2	5.2	5.6	6.2
	6-19 times	0.2	0.4	0.8	1.1	1.5	1.4	1.9
	20-40 times	0.0	0.2	0.3	0.6	0.6	0.7	0.8
	40+ times	0.2	0.3	0.5	0.9	1.3	1.4	1.7
Smokeless Tobacco	No Answer	0.3	0.3	0.4	0.4	0.3	0.4	0.2
	Never	97.1	94.7	91.3	88.0	84.2	82.4	79.4
	1-5 times	2.0	3.6	5.4	6.4	8.0	8.9	9.8
	6-19 times	0.3	0.6	1.0	1.7	2.0	2.3	2.6
	20-40 times	0.1	0.3	0.6	1.0	1.3	1.3	1.8
	40+ times	0.3	0.5	1.3	2.5	4.2	4.8	6.2
Alcohol	No Answer	0.7	0.8	0.9	1.1	1.0	1.2	1.3
	Never	74.0	65.9	52.5	43.5	36.7	32.1	28.5
	1-5 times	20.6	23.2	27.9	26.6	25.8	24.0	20.8
	6-19 times	3.1	6.1	10.0	13.4	14.9	15.6	15.4
	20-40 times	0.8	2.0	4.2	7.0	9.1	11.0	11.8
	40+ times	0.8	2.1	4.4	8.5	12.4	16.1	22.3
Marijuana	No Answer	0.5	0.5	0.7	0.8	0.8	0.7	1.1
	Never	96.7	92.7	83.7	76.7	69.1	64.8	61.8
	1-5 times	1.7	3.9	7.6	9.2	11.5	12.5	12.9
	6-19 times	0.4	1.1	2.7	3.9	5.3	6.4	6.2
	20-40 times	0.3	0.6	1.8	2.7	3.5	4.1	4.5
	40+ times	0.5	1.2	3.5	6.7	9.7	11.5	13.4
Cocaine	No Answer	0.3	0.2	0.3	0.4	0.4	0.4	0.4
	Never	99.1	98.3	97.1	95.2	93.7	91.8	91.0
	1-5 times	0.3	0.9	1.5	2.6	3.5	4.5	4.4
	6-19 times	0.1	0.2	0.4	0.8	1.0	1.4	1.6
	20-40 times	0.0	0.1	0.2	0.5	0.6	0.9	1.1
	40+ times	0.2	0.3	0.3	0.6	0.8	1.1	1.3

General



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Crack	No Answer	0.4	0.3	0.4	0.4	0.5	0.4	0.5
	Never	98.9	98.1	97.3	96.6	96.1	95.9	95.5
	1-5 times	0.4	1.0	1.3	1.7	2.0	2.1	2.2
	6-19 times	0.1	0.2	0.4	0.6	0.5	0.6	0.8
	20-40 times	0.1	0.1	0.2	0.3	0.2	0.4	0.3
Inhalants	40+ times	0.2	0.3	0.3	0.4	0.6	0.6	0.7
	No Answer	0.9	0.5	0.5	0.4	0.4	0.4	0.5
	Never	92.6	90.5	89.1	88.3	88.5	89.5	89.5
	1-5 times	4.8	6.4	7.0	7.3	7.0	6.2	6.2
	6-19 times	0.9	1.3	1.9	2.3	2.1	2.1	2.1
Amphetamines	20-40 times	0.3	0.5	0.6	0.8	0.9	0.8	0.8
	40+ times	0.5	0.8	0.9	1.0	1.0	1.0	1.0
	No Answer	1.6	0.8	0.5	0.5	0.6	0.5	0.6
	Never	97.9	97.8	96.3	92.8	90.4	88.7	88.1
	1-5 times	0.3	0.8	1.9	3.5	4.2	4.8	4.8
Methamphetamines	6-19 times	0.1	0.3	0.7	1.4	2.4	2.7	2.8
	20-40 times	0.0	0.1	0.3	0.8	1.1	1.5	1.7
	40+ times	0.2	0.2	0.3	1.1	1.3	1.8	2.1
	No Answer	0.9	0.5	0.4	0.4	0.5	0.5	0.6
	Never	98.6	98.3	97.6	96.6	96.0	95.2	94.4
Ritalin	1-5 times	0.2	0.6	1.2	1.7	1.9	2.0	2.3
	6-19 times	0.1	0.2	0.4	0.5	0.5	0.8	1.0
	20-40 times	0.0	0.1	0.2	0.3	0.4	0.6	0.6
	40+ times	0.1	0.2	0.2	0.4	0.7	0.8	1.1
	No Answer	1.5	0.7	0.6	0.6	0.8	0.7	0.7
Tranquilizers	Never	97.6	97.5	96.2	94.0	92.8	92.7	92.3
	1-5 times	0.5	1.1	1.9	3.2	3.6	3.7	3.7
	6-19 times	0.1	0.4	0.6	1.1	1.4	1.4	1.4
	20-40 times	0.1	0.1	0.3	0.5	0.7	0.6	0.9
	40+ times	0.2	0.3	0.4	0.7	0.8	1.0	1.0
Narcotics	No Answer	0.9	0.5	0.6	0.4	0.6	0.5	0.7
	Never	95.3	93.5	90.5	87.1	85.2	85.0	84.8
	1-5 times	3.0	4.3	5.8	6.9	7.8	7.3	7.0
	6-19 times	0.5	0.9	1.7	2.9	3.2	3.7	3.5
	20-40 times	0.1	0.4	0.7	1.2	1.5	1.6	1.8
General	40+ times	0.2	0.5	0.7	1.4	1.8	1.9	2.2
	No Answer	1.3	0.6	0.5	0.5	0.7	0.5	0.6
	Never	97.9	97.4	95.4	92.1	89.5	88.2	87.5
	1-5 times	0.4	1.1	2.4	3.6	4.8	5.0	5.1
	6-19 times	0.1	0.4	0.8	1.7	2.3	2.9	2.8
Narcotics	20-40 times	0.0	0.1	0.4	1.0	1.3	1.5	1.9
	40+ times	0.1	0.3	0.5	1.1	1.5	1.9	2.3

General



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Psychedelics (All)	No Answer	1.2	0.5	0.3	0.2	0.4	0.3	0.4
	Never	98.2	98.0	96.9	95.6	94.3	93.4	92.6
	1-5 times	0.3	0.9	1.7	2.5	3.2	3.7	3.9
	6-19 times	0.1	0.2	0.4	0.7	0.9	1.2	1.5
	20-40 times	0.0	0.1	0.3	0.4	0.5	0.7	0.7
	40+ times	0.2	0.2	0.4	0.7	0.7	0.8	1.0
LSD Only	No Answer	1.4	0.6	0.4	0.4	0.6	0.4	0.5
	Never	98.1	98.3	97.3	96.2	95.1	94.7	94.2
	1-5 times	0.2	0.7	1.4	2.0	2.7	2.8	3.2
	6-19 times	0.1	0.2	0.4	0.5	0.7	0.8	0.9
	20-40 times	0.0	0.1	0.2	0.3	0.3	0.6	0.5
	40+ times	0.2	0.2	0.3	0.5	0.6	0.6	0.8
Other Psychedelics	No Answer	1.9	0.8	0.6	0.5	0.6	0.5	0.6
	Never	97.7	98.3	98.1	97.3	96.7	95.7	94.8
	1-5 times	0.2	0.4	0.8	1.2	1.5	2.2	2.3
	6-19 times	0.1	0.2	0.2	0.4	0.5	0.8	1.2
	20-40 times	0.0	0.1	0.1	0.2	0.3	0.3	0.5
	40+ times	0.1	0.2	0.2	0.4	0.4	0.4	0.6
Heroin	No Answer	1.1	0.4	0.4	0.4	0.5	0.4	0.5
	Never	98.4	98.5	98.1	97.7	97.5	97.2	97.1
	1-5 times	0.2	0.6	0.8	0.9	1.0	1.0	0.9
	6-19 times	0.1	0.2	0.2	0.3	0.3	0.4	0.4
	20-40 times	0.0	0.1	0.2	0.3	0.2	0.3	0.4
	40+ times	0.2	0.2	0.2	0.4	0.5	0.6	0.7
Steroids	No Answer	0.7	0.6	0.5	0.4	0.5	0.5	0.7
	Never	98.5	98.3	98.0	97.7	97.3	97.5	97.0
	1-5 times	0.4	0.6	0.8	0.9	0.9	0.8	1.0
	6-19 times	0.1	0.1	0.3	0.3	0.4	0.3	0.3
	20-40 times	0.1	0.1	0.1	0.2	0.2	0.2	0.3
	40+ times	0.2	0.3	0.3	0.5	0.7	0.6	0.8
Injected Drugs	No Answer	0.9	0.6	0.6	0.6	0.6	0.6	0.7
	Never	98.2	98.1	97.8	97.7	97.4	97.3	97.1
	1-5 times	0.6	0.7	0.9	0.8	0.9	0.9	0.8
	6-19 times	0.1	0.3	0.3	0.3	0.3	0.3	0.4
	20-40 times	0.0	0.1	0.2	0.2	0.2	0.2	0.2
	40+ times	0.2	0.2	0.4	0.5	0.6	0.7	0.8
MDMA	No Answer	1.8	0.7	0.5	0.5	0.6	0.5	0.5
	Never	97.8	98.0	97.2	95.8	95.0	94.1	93.9
	1-5 times	0.2	0.7	1.5	2.4	2.8	3.4	3.3
	6-19 times	0.1	0.2	0.4	0.6	0.7	0.9	0.9
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.5	0.5
	40+ times	0.1	0.2	0.3	0.5	0.5	0.6	0.8

General

**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Rohypnol	No Answer	2.0	0.9	0.6	0.5	0.6	0.4	0.6
	Never	97.7	98.3	98.2	97.7	97.5	97.6	97.7
	1-5 times	0.1	0.4	0.6	1.1	1.0	1.1	0.7
	6-19 times	0.1	0.1	0.2	0.3	0.3	0.4	0.3
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.2	0.2
	40+ times	0.1	0.2	0.2	0.2	0.3	0.3	0.4
GHB	No Answer	2.0	1.0	0.6	0.5	0.6	0.5	0.6
	Never	97.7	98.5	98.6	98.5	98.2	98.3	98.1
	1-5 times	0.1	0.2	0.4	0.5	0.6	0.6	0.5
	6-19 times	0.0	0.1	0.1	0.2	0.2	0.2	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.2
	40+ times	0.1	0.2	0.2	0.2	0.3	0.3	0.4

General



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	0.9	0.5	0.6	0.6	0.4	0.5	0.3
	Never	86.1	79.5	69.4	63.2	55.6	52.1	45.3
	Once or Twice	9.3	12.2	15.6	15.3	17.1	16.7	17.7
	Occasionally	1.6	3.2	6.1	8.2	11.6	12.7	15.0
	Past Regularly	1.2	2.4	3.9	5.0	5.1	5.0	5.8
	Current Regularly	0.9	2.2	4.4	7.7	10.2	13.1	15.9
Cigars	No Answer	0.3	0.3	0.4	0.5	0.4	0.5	0.4
	Never	92.9	89.5	81.2	72.8	63.4	56.7	48.1
	1-5 times	5.5	7.3	11.9	14.3	17.9	19.9	20.9
	6-19 times	0.5	1.2	2.7	5.2	7.3	9.3	11.7
	20-40 times	0.2	0.7	1.3	2.7	4.7	5.5	8.2
	40+ times	0.6	1.0	2.5	4.4	6.4	8.1	10.6
Pipe	No Answer	0.6	0.9	1.2	1.5	1.5	1.7	1.7
	Never	97.2	95.4	92.5	88.9	85.5	84.2	81.2
	1-5 times	1.6	2.7	4.0	5.8	7.6	8.2	9.5
	6-19 times	0.3	0.5	1.0	1.5	2.3	2.4	3.2
	20-40 times	0.1	0.2	0.5	0.9	1.0	1.3	1.4
	40+ times	0.3	0.4	0.8	1.5	2.1	2.4	3.0
Smokeless Tobacco	No Answer	0.3	0.3	0.5	0.5	0.5	0.5	0.2
	Never	95.7	92.2	87.0	81.3	74.5	70.9	65.6
	1-5 times	3.0	5.2	7.6	8.9	11.3	12.6	14.4
	6-19 times	0.5	0.9	1.6	2.8	3.5	4.1	4.7
	20-40 times	0.2	0.5	1.0	1.9	2.4	2.4	3.3
	40+ times	0.4	0.9	2.2	4.6	7.9	9.4	11.8
Alcohol	No Answer	0.7	0.8	1.0	1.0	1.1	0.9	1.2
	Never	71.8	65.2	53.9	44.6	38.0	33.3	27.9
	1-5 times	21.9	23.4	26.3	24.3	22.5	20.7	17.7
	6-19 times	3.7	6.3	9.7	12.6	13.7	14.1	13.4
	20-40 times	0.9	2.0	4.2	7.2	9.3	10.6	10.6
	40+ times	0.9	2.3	4.9	10.4	15.5	20.4	29.2
Marijuana	No Answer	0.6	0.6	0.7	0.9	0.8	0.8	1.0
	Never	95.9	91.5	82.7	75.0	67.2	62.7	57.3
	1-5 times	2.0	4.5	7.6	8.7	11.0	12.1	13.1
	6-19 times	0.5	1.2	2.9	4.0	5.2	6.0	6.3
	20-40 times	0.3	0.7	1.8	2.9	3.8	3.7	4.9
	40+ times	0.7	1.4	4.2	8.5	12.0	14.7	17.3
Cocaine	No Answer	0.3	0.3	0.4	0.5	0.5	0.5	0.4
	Never	99.0	98.1	97.0	94.7	92.5	90.3	88.5
	1-5 times	0.3	0.9	1.4	2.6	4.1	5.1	5.6
	6-19 times	0.1	0.2	0.4	0.9	1.1	1.6	2.1
	20-40 times	0.1	0.1	0.2	0.5	0.7	0.9	1.5
	40+ times	0.2	0.4	0.5	0.8	1.2	1.6	1.9

Male



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Crack	No Answer	0.4	0.4	0.4	0.6	0.6	0.4	0.6
	Never	98.8	98.0	97.4	96.2	95.3	94.8	94.0
	1-5 times	0.4	0.9	1.2	1.7	2.2	2.2	2.8
	6-19 times	0.1	0.2	0.4	0.6	0.8	0.9	1.1
	20-40 times	0.1	0.1	0.2	0.3	0.3	0.6	0.4
Inhalants	40+ times	0.2	0.4	0.4	0.6	0.9	1.0	1.0
	No Answer	0.9	0.6	0.5	0.5	0.5	0.5	0.5
	Never	91.9	90.6	89.6	88.7	88.2	89.0	86.6
	1-5 times	5.1	6.3	6.4	6.8	6.8	6.1	7.7
	6-19 times	1.1	1.2	1.8	2.1	2.1	2.2	2.8
Amphetamines	20-40 times	0.3	0.5	0.6	0.8	1.0	0.9	1.0
	40+ times	0.7	0.8	1.1	1.1	1.4	1.3	1.5
	No Answer	1.8	1.0	0.6	0.5	0.7	0.5	0.5
	Never	97.6	97.7	96.6	93.0	90.6	88.9	86.9
	1-5 times	0.3	0.6	1.5	3.2	3.6	4.7	4.9
Methamphetamines	6-19 times	0.1	0.2	0.7	1.4	2.4	2.2	3.3
	20-40 times	0.0	0.2	0.2	0.8	1.1	1.5	1.6
	40+ times	0.2	0.3	0.4	1.1	1.6	2.2	2.7
	No Answer	1.0	0.6	0.5	0.6	0.6	0.6	0.7
	Never	98.4	98.3	97.6	96.3	95.5	94.6	93.2
Ritalin	1-5 times	0.2	0.6	1.0	1.8	1.9	2.0	2.6
	6-19 times	0.1	0.2	0.4	0.5	0.6	0.9	1.3
	20-40 times	0.0	0.1	0.2	0.3	0.5	0.8	0.7
	40+ times	0.1	0.2	0.3	0.6	0.9	1.1	1.5
	No Answer	1.6	0.9	0.7	0.9	0.8	0.8	0.8
Tranquilizers	Never	97.5	97.4	96.4	93.8	92.2	91.9	90.7
	1-5 times	0.5	0.9	1.5	2.9	3.5	3.4	4.4
	6-19 times	0.1	0.3	0.6	1.1	1.7	1.7	1.7
	20-40 times	0.1	0.1	0.3	0.5	0.9	0.7	1.0
	40+ times	0.2	0.3	0.6	0.9	0.9	1.5	1.4
Narcotics	No Answer	1.0	0.6	0.6	0.5	0.7	0.6	0.8
	Never	95.3	94.5	92.5	89.2	86.6	86.0	83.4
	1-5 times	2.8	3.3	4.3	5.3	6.3	5.9	7.0
	6-19 times	0.5	0.7	1.4	2.5	3.1	3.7	4.0
	20-40 times	0.1	0.3	0.5	0.9	1.3	1.6	2.3
	40+ times	0.3	0.5	0.7	1.5	2.0	2.1	2.5
	No Answer	1.3	0.7	0.6	0.6	0.6	0.6	0.7
	Never	97.9	97.4	95.4	91.9	89.0	86.7	84.3
	1-5 times	0.4	1.0	2.1	3.4	4.5	5.1	5.5
	6-19 times	0.1	0.4	0.8	1.8	2.5	3.3	3.7
	20-40 times	0.1	0.1	0.4	0.9	1.4	1.8	2.6
	40+ times	0.2	0.3	0.6	1.3	1.9	2.5	3.3

Male



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Psychedelics (All)	No Answer	1.3	0.6	0.3	0.3	0.5	0.3	0.4
	Never	98.0	97.9	96.5	94.5	92.4	91.0	89.6
	1-5 times	0.4	0.9	1.8	3.1	4.1	4.6	5.1
	6-19 times	0.1	0.3	0.5	0.8	1.3	1.8	2.2
	20-40 times	0.1	0.1	0.3	0.5	0.6	1.0	1.1
	40+ times	0.2	0.3	0.6	0.9	1.2	1.3	1.6
LSD Only	No Answer	1.6	0.7	0.5	0.5	0.7	0.5	0.7
	Never	97.9	98.1	96.8	95.1	93.4	92.9	91.8
	1-5 times	0.2	0.8	1.6	2.6	3.5	3.6	4.3
	6-19 times	0.1	0.2	0.4	0.7	1.0	1.3	1.3
	20-40 times	0.0	0.1	0.2	0.4	0.4	0.8	0.7
	40+ times	0.2	0.3	0.5	0.7	1.0	1.0	1.2
Other Psychedelics	No Answer	2.1	1.0	0.7	0.6	0.8	0.6	0.8
	Never	97.5	98.2	98.0	96.7	95.7	94.1	92.5
	1-5 times	0.2	0.3	0.7	1.5	1.9	2.9	3.2
	6-19 times	0.1	0.1	0.2	0.4	0.7	1.2	1.7
	20-40 times	0.0	0.1	0.1	0.2	0.4	0.5	0.8
	40+ times	0.1	0.2	0.3	0.5	0.5	0.8	1.0
Heroin	No Answer	1.1	0.5	0.4	0.5	0.7	0.5	0.7
	Never	98.3	98.3	97.9	97.2	96.9	96.2	96.0
	1-5 times	0.3	0.6	0.8	1.0	1.0	1.3	1.3
	6-19 times	0.1	0.2	0.3	0.5	0.4	0.6	0.5
	20-40 times	0.0	0.1	0.2	0.3	0.3	0.4	0.5
	40+ times	0.2	0.3	0.3	0.5	0.7	0.9	1.0
Steroids	No Answer	0.8	0.7	0.6	0.5	0.7	0.7	0.9
	Never	98.2	97.7	97.4	96.9	96.0	96.0	95.3
	1-5 times	0.6	0.8	1.0	1.1	1.3	1.3	1.6
	6-19 times	0.1	0.2	0.4	0.6	0.6	0.6	0.5
	20-40 times	0.1	0.2	0.2	0.3	0.3	0.3	0.5
	40+ times	0.2	0.5	0.5	0.7	1.1	1.1	1.2
Injected Drugs	No Answer	1.1	0.8	0.7	0.7	0.9	0.9	0.8
	Never	98.0	97.9	97.7	97.2	96.7	96.3	96.1
	1-5 times	0.5	0.7	0.8	0.8	1.0	1.0	1.1
	6-19 times	0.1	0.2	0.3	0.3	0.5	0.4	0.5
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.4	0.2
	40+ times	0.2	0.3	0.4	0.6	0.8	1.0	1.2
MDMA	No Answer	1.9	0.9	0.6	0.7	0.8	0.6	0.7
	Never	97.7	97.8	97.2	95.3	94.2	93.1	92.1
	1-5 times	0.2	0.7	1.3	2.5	2.9	3.6	4.1
	6-19 times	0.1	0.1	0.4	0.7	0.7	1.1	1.3
	20-40 times	0.0	0.1	0.2	0.3	0.5	0.7	0.6
	40+ times	0.1	0.3	0.4	0.6	0.8	0.9	1.2

Male



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Rohypnol	No Answer	2.2	1.0	0.6	0.7	0.7	0.6	0.8
	Never	97.5	98.2	98.1	97.3	97.0	97.0	96.8
	1-5 times	0.1	0.3	0.6	1.2	1.2	1.2	0.9
	6-19 times	0.1	0.1	0.3	0.3	0.4	0.5	0.4
	20-40 times	0.0	0.1	0.2	0.2	0.2	0.2	0.3
	40+ times	0.1	0.2	0.2	0.4	0.5	0.4	0.7
GHB	No Answer	2.2	1.2	0.7	0.7	0.8	0.6	0.9
	Never	97.6	98.1	98.4	98.1	97.6	97.5	97.2
	1-5 times	0.1	0.2	0.4	0.6	0.7	0.9	0.8
	6-19 times	0.0	0.1	0.2	0.2	0.3	0.4	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.2	0.2
	40+ times	0.1	0.2	0.3	0.3	0.5	0.5	0.6

Male



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	0.4	0.3	0.4	0.4	0.3	0.2	0.3
	Never	88.7	80.4	69.3	63.3	57.6	53.5	52.3
	Once or Twice	7.7	11.5	15.3	14.6	16.3	16.6	17.9
	Occasionally	1.6	4.1	7.1	9.4	10.9	11.8	12.7
	Past Regularly	0.9	1.8	3.1	4.2	5.2	5.6	4.3
	Current Regularly	0.7	1.9	4.8	8.1	9.8	12.3	12.5
Cigars	No Answer	0.2	0.2	0.3	0.3	0.3	0.3	0.3
	Never	96.0	93.3	88.9	85.1	81.6	79.8	75.6
	1-5 times	2.9	4.8	7.7	9.7	12.5	13.8	17.2
	6-19 times	0.4	0.7	1.3	1.9	2.4	2.8	3.4
	20-40 times	0.1	0.4	0.6	1.0	1.1	1.3	1.3
	40+ times	0.3	0.7	1.2	2.0	2.1	2.0	2.2
Pipe	No Answer	0.5	0.8	1.2	1.4	1.4	2.3	2.2
	Never	98.1	97.2	95.4	94.4	94.0	93.1	93.5
	1-5 times	1.0	1.5	2.5	2.8	3.1	3.3	3.0
	6-19 times	0.1	0.3	0.6	0.7	0.7	0.6	0.6
	20-40 times	0.0	0.1	0.2	0.4	0.3	0.2	0.3
	40+ times	0.1	0.1	0.3	0.3	0.5	0.5	0.4
Smokeless Tobacco	No Answer	0.2	0.2	0.2	0.2	0.2	0.2	0.1
	Never	98.4	97.3	95.7	94.3	93.1	92.9	93.1
	1-5 times	1.0	1.9	3.1	4.1	5.0	5.5	5.3
	6-19 times	0.1	0.3	0.4	0.6	0.6	0.6	0.6
	20-40 times	0.0	0.1	0.1	0.3	0.3	0.3	0.3
	40+ times	0.2	0.2	0.4	0.5	0.7	0.5	0.6
Alcohol	No Answer	0.7	0.8	0.8	1.1	1.0	1.5	1.3
	Never	76.3	66.6	51.2	42.4	35.5	31.0	29.1
	1-5 times	19.2	23.0	29.6	28.8	28.9	27.0	24.0
	6-19 times	2.5	5.8	10.3	14.3	16.0	16.9	17.3
	20-40 times	0.7	2.0	4.3	6.7	9.0	11.4	12.9
	40+ times	0.6	1.8	3.8	6.7	9.7	12.1	15.5
Marijuana	No Answer	0.3	0.3	0.7	0.7	0.7	0.7	1.1
	Never	97.4	94.0	84.8	78.3	71.0	66.8	66.5
	1-5 times	1.4	3.3	7.5	9.7	11.9	12.9	12.8
	6-19 times	0.4	0.9	2.5	3.8	5.5	6.7	6.0
	20-40 times	0.2	0.5	1.7	2.5	3.3	4.4	4.2
	40+ times	0.3	1.0	2.9	5.1	7.5	8.5	9.5
Cocaine	No Answer	0.2	0.1	0.2	0.3	0.3	0.3	0.3
	Never	99.3	98.5	97.3	95.6	94.8	93.3	93.8
	1-5 times	0.2	0.9	1.6	2.5	3.0	3.9	3.2
	6-19 times	0.1	0.2	0.4	0.7	0.9	1.1	1.1
	20-40 times	0.0	0.1	0.3	0.5	0.5	0.8	0.8
	40+ times	0.1	0.2	0.1	0.4	0.5	0.6	0.8

Female



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Crack	No Answer	0.3	0.2	0.4	0.3	0.4	0.3	0.3
	Never	98.9	98.3	97.4	96.9	96.9	96.9	97.1
	1-5 times	0.5	1.0	1.4	1.7	1.8	1.9	1.6
	6-19 times	0.1	0.2	0.4	0.5	0.3	0.4	0.4
	20-40 times	0.0	0.1	0.2	0.3	0.2	0.2	0.2
Inhalants	40+ times	0.1	0.2	0.2	0.2	0.4	0.3	0.4
	No Answer	0.8	0.4	0.5	0.3	0.3	0.3	0.4
	Never	93.3	90.5	88.7	87.9	88.9	90.0	92.4
	1-5 times	4.5	6.5	7.5	7.8	7.2	6.2	4.8
	6-19 times	0.7	1.5	2.0	2.4	2.1	1.9	1.4
Amphetamines	20-40 times	0.3	0.5	0.7	0.8	0.8	0.7	0.6
	40+ times	0.4	0.7	0.7	0.9	0.7	0.8	0.5
	No Answer	1.3	0.5	0.4	0.4	0.5	0.4	0.5
	Never	98.2	98.0	96.1	92.6	90.2	88.7	89.5
	1-5 times	0.2	0.9	2.2	3.7	4.8	4.9	4.7
Methamphetamines	6-19 times	0.1	0.3	0.7	1.3	2.3	3.1	2.2
	20-40 times	0.0	0.1	0.4	0.9	1.2	1.6	1.7
	40+ times	0.1	0.2	0.2	1.0	1.0	1.4	1.4
	No Answer	0.8	0.3	0.4	0.3	0.3	0.5	0.5
	Never	98.8	98.5	97.7	97.0	96.5	95.9	95.9
Ritalin	1-5 times	0.2	0.7	1.3	1.7	1.8	2.0	1.9
	6-19 times	0.1	0.2	0.3	0.4	0.5	0.6	0.6
	20-40 times	0.0	0.1	0.2	0.3	0.4	0.4	0.5
	40+ times	0.1	0.2	0.2	0.3	0.5	0.5	0.6
	No Answer	1.3	0.5	0.6	0.4	0.7	0.4	0.5
Tranquilizers	Never	97.8	97.5	96.1	94.1	93.4	93.5	94.2
	1-5 times	0.6	1.2	2.2	3.4	3.7	3.9	3.0
	6-19 times	0.1	0.3	0.6	1.1	1.1	1.2	1.1
	20-40 times	0.1	0.1	0.3	0.5	0.5	0.5	0.7
	40+ times	0.1	0.3	0.3	0.5	0.6	0.5	0.6
Narcotics	No Answer	0.8	0.3	0.5	0.4	0.4	0.4	0.5
	Never	95.2	92.5	88.6	85.2	83.9	84.1	86.4
	1-5 times	3.2	5.3	7.3	8.4	9.1	8.4	6.9
	6-19 times	0.5	1.1	2.1	3.2	3.3	3.8	3.0
	20-40 times	0.1	0.4	0.9	1.5	1.6	1.6	1.4
	40+ times	0.2	0.4	0.6	1.4	1.6	1.6	1.9
	No Answer	1.3	0.4	0.3	0.4	0.6	0.4	0.4
	Never	97.9	97.4	95.4	92.4	90.0	89.7	90.8
	1-5 times	0.4	1.2	2.6	3.7	5.0	4.8	4.6
	6-19 times	0.1	0.5	0.9	1.6	2.1	2.5	1.8
	20-40 times	0.0	0.1	0.4	1.0	1.2	1.2	1.2
	40+ times	0.1	0.2	0.4	0.9	1.0	1.4	1.2

Female



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Psychedelics (All)	No Answer	1.1	0.4	0.2	0.2	0.3	0.2	0.2
	Never	98.4	98.2	97.4	96.6	96.0	95.7	95.7
	1-5 times	0.2	0.9	1.6	1.9	2.5	2.8	2.6
	6-19 times	0.1	0.2	0.4	0.6	0.5	0.6	0.8
	20-40 times	0.0	0.1	0.3	0.3	0.3	0.4	0.2
	40+ times	0.2	0.2	0.1	0.4	0.3	0.3	0.4
LSD Only	No Answer	1.3	0.5	0.3	0.3	0.4	0.3	0.3
	Never	98.4	98.5	97.9	97.2	96.7	96.5	96.7
	1-5 times	0.2	0.6	1.2	1.5	2.0	2.2	2.0
	6-19 times	0.1	0.1	0.3	0.4	0.4	0.4	0.5
	20-40 times	0.0	0.0	0.2	0.2	0.3	0.4	0.2
	40+ times	0.1	0.2	0.1	0.3	0.2	0.2	0.4
Other Psychedelics	No Answer	1.6	0.6	0.4	0.4	0.5	0.4	0.4
	Never	98.0	98.5	98.3	98.0	97.7	97.3	97.3
	1-5 times	0.2	0.5	0.8	1.0	1.2	1.6	1.4
	6-19 times	0.1	0.2	0.2	0.3	0.3	0.4	0.6
	20-40 times	0.0	0.1	0.2	0.2	0.2	0.1	0.2
	40+ times	0.1	0.1	0.1	0.2	0.2	0.1	0.2
Heroin	No Answer	1.0	0.3	0.4	0.2	0.4	0.3	0.3
	Never	98.6	98.7	98.4	98.3	98.1	98.2	98.5
	1-5 times	0.2	0.5	0.8	0.9	0.9	0.7	0.5
	6-19 times	0.1	0.2	0.1	0.2	0.2	0.3	0.2
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.2	0.2
	40+ times	0.1	0.1	0.1	0.2	0.2	0.2	0.3
Steroids	No Answer	0.7	0.4	0.3	0.3	0.4	0.4	0.4
	Never	98.9	98.9	98.7	98.6	98.6	98.9	98.9
	1-5 times	0.3	0.5	0.6	0.6	0.6	0.4	0.3
	6-19 times	0.1	0.1	0.2	0.1	0.2	0.0	0.1
	20-40 times	0.0	0.1	0.1	0.2	0.1	0.0	0.1
	40+ times	0.1	0.1	0.2	0.2	0.2	0.2	0.2
Injected Drugs	No Answer	0.7	0.4	0.4	0.4	0.4	0.4	0.4
	Never	98.5	98.3	98.0	98.1	98.1	98.3	98.4
	1-5 times	0.6	0.8	0.9	0.8	0.8	0.7	0.5
	6-19 times	0.1	0.3	0.2	0.3	0.3	0.2	0.1
	20-40 times	0.0	0.1	0.2	0.1	0.1	0.1	0.2
	40+ times	0.1	0.2	0.2	0.4	0.3	0.4	0.4
MDMA	No Answer	1.7	0.5	0.3	0.3	0.4	0.3	0.2
	Never	98.0	98.3	97.3	96.3	95.7	95.0	95.9
	1-5 times	0.1	0.7	1.6	2.3	2.7	3.2	2.6
	6-19 times	0.1	0.2	0.4	0.5	0.7	0.8	0.6
	20-40 times	0.0	0.1	0.2	0.2	0.2	0.3	0.3
	40+ times	0.1	0.1	0.2	0.4	0.2	0.4	0.3

Female



**Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Rohypnol	No Answer	1.7	0.8	0.5	0.4	0.5	0.2	0.4
	Never	97.9	98.5	98.4	98.1	98.1	98.2	98.7
	1-5 times	0.1	0.5	0.7	1.0	0.9	1.1	0.5
	6-19 times	0.1	0.1	0.2	0.2	0.2	0.2	0.1
	20-40 times	0.0	0.1	0.1	0.2	0.1	0.2	0.2
	40+ times	0.1	0.1	0.1	0.1	0.2	0.2	0.1
GHB	No Answer	1.8	0.8	0.5	0.3	0.5	0.4	0.3
	Never	97.9	98.8	98.9	99.0	98.8	99.0	99.2
	1-5 times	0.1	0.2	0.3	0.4	0.5	0.3	0.2
	6-19 times	0.1	0.0	0.1	0.1	0.1	0.1	0.1
	20-40 times	0.0	0.0	0.0	0.0	0.1	0.1	0.1
	40+ times	0.1	0.1	0.1	0.2	0.1	0.1	0.2

Female



Finding sustainable resources is easy

Whether you need program planning,
evaluation, training, or grants consulting,
our staff can connect you to the resources
you need to sustain drug prevention in your community.

as calling 800-346-3077 or
visit www.drugs.indiana.edu.



**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	1.3	1.0	1.1	1.0	1.1	1.0	1.1
	None	92.6	87.3	79.7	74.0	68.9	65.3	62.0
	Few times	4.0	7.2	11.3	12.9	15.4	16.5	18.1
	1-5 cigarettes/day	1.3	2.8	4.2	5.9	6.7	7.7	7.6
	1/2 pack/day	0.4	0.8	1.7	3.2	3.9	4.8	5.3
	1 pack/day	0.2	0.5	1.1	1.9	2.7	3.2	4.4
	1 & 1/2 pack/day	0.0	0.2	0.3	0.5	0.6	0.7	0.7
	2+ pack/day	0.3	0.3	0.5	0.6	0.7	0.7	0.8
Cigars	No Answer	2.2	2.5	2.7	3.3	3.3	3.7	3.7
	Never	94.4	91.7	87.0	81.3	76.6	73.2	67.2
	1-5 times	2.3	3.6	6.0	8.6	11.2	12.8	15.9
	6-19 times	0.5	0.9	1.8	2.8	3.7	4.5	5.8
	20-40 times	0.2	0.5	0.9	1.4	2.1	2.1	3.1
	40+ times	0.3	0.7	1.6	2.5	3.2	3.6	4.2
Pipe	No Answer	2.4	3.0	3.3	4.2	4.2	4.8	5.0
	Never	96.5	94.9	93.3	90.7	89.6	88.6	87.2
	1-5 times	0.7	1.3	2.0	2.7	3.5	3.7	4.2
	6-19 times	0.2	0.4	0.6	1.0	1.1	1.2	1.5
	20-40 times	0.1	0.2	0.4	0.5	0.7	0.6	0.8
	40+ times	0.1	0.2	0.5	0.9	1.0	1.1	1.4
Smokeless Tobacco	No Answer	2.1	2.4	2.6	3.3	3.2	3.5	3.6
	Never	95.9	93.7	90.9	87.2	84.6	83.3	81.0
	1-5 times	1.2	2.4	3.7	4.8	5.4	6.0	6.7
	6-19 times	0.3	0.7	1.1	1.4	1.8	1.9	2.1
	20-40 times	0.1	0.3	0.5	1.0	1.2	1.1	1.3
	40+ times	0.3	0.5	1.2	2.3	3.8	4.2	5.3
Alcohol	No Answer	2.5	2.8	3.0	3.6	3.5	3.9	3.9
	Never	80.1	71.6	58.7	49.1	42.8	39.1	34.6
	1-5 times	14.0	17.6	24.0	25.4	25.7	25.6	23.8
	6-19 times	2.2	5.0	8.3	11.7	13.4	14.2	14.8
	20-40 times	0.6	1.7	3.2	5.1	7.0	8.4	9.9
	40+ times	0.5	1.2	2.7	5.2	7.6	8.9	12.9
Marijuana	No Answer	2.3	2.8	3.0	3.7	3.6	4.0	4.2
	Never	95.4	91.6	84.3	77.9	72.5	69.5	68.3
	1-5 times	1.3	3.1	6.1	7.8	9.4	10.7	10.9
	6-19 times	0.4	1.0	2.5	3.5	4.7	5.1	4.7
	20-40 times	0.3	0.6	1.6	2.5	3.1	3.2	3.1
	40+ times	0.3	0.9	2.6	4.7	6.6	7.5	8.9

General



**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	2.4	2.7	3.0	3.7	3.7	4.2	4.3
	Never	97.2	96.2	95.1	93.0	91.9	90.5	89.6
	1-5 times	0.2	0.7	1.1	1.9	2.6	3.1	3.5
	6-19 times	0.1	0.2	0.4	0.7	0.9	1.1	1.1
	20-40 times	0.0	0.2	0.2	0.3	0.4	0.6	0.7
	40+ times	0.1	0.1	0.2	0.5	0.5	0.6	0.8
Crack	No Answer	2.5	2.7	3.0	3.7	3.8	4.3	4.3
	Never	97.0	96.2	95.3	94.1	93.9	93.3	93.2
	1-5 times	0.3	0.6	1.0	1.2	1.2	1.3	1.3
	6-19 times	0.1	0.3	0.3	0.4	0.5	0.6	0.5
	20-40 times	0.0	0.1	0.2	0.2	0.2	0.3	0.3
	40+ times	0.1	0.1	0.2	0.4	0.3	0.3	0.4
Inhalants	No Answer	2.9	2.9	3.0	3.7	3.8	4.3	4.3
	Never	92.6	90.9	89.7	88.8	89.6	90.6	91.0
	1-5 times	3.2	4.1	4.7	4.9	4.3	3.3	2.9
	6-19 times	0.9	1.2	1.6	1.6	1.2	1.1	1.0
	20-40 times	0.2	0.5	0.5	0.5	0.6	0.4	0.4
	40+ times	0.3	0.5	0.5	0.6	0.5	0.4	0.4
Amphetamines	No Answer	3.4	3.1	3.1	3.7	3.9	4.2	4.3
	Never	96.2	95.8	94.4	91.2	89.2	88.1	88.2
	1-5 times	0.2	0.6	1.5	2.6	3.4	3.8	3.6
	6-19 times	0.1	0.2	0.6	1.4	1.9	2.1	1.7
	20-40 times	0.0	0.2	0.2	0.5	0.9	1.0	1.2
	40+ times	0.1	0.1	0.2	0.6	0.6	0.8	1.0
Methamphetamines	No Answer	3.1	3.0	3.0	3.8	3.9	4.3	4.4
	Never	96.6	96.1	95.6	94.1	93.7	93.1	92.5
	1-5 times	0.2	0.5	0.9	1.2	1.2	1.3	1.5
	6-19 times	0.1	0.2	0.3	0.4	0.5	0.7	0.6
	20-40 times	0.0	0.1	0.1	0.2	0.3	0.4	0.5
	40+ times	0.0	0.1	0.1	0.3	0.3	0.4	0.5
Ritalin	No Answer	3.5	3.2	3.1	3.8	4.0	4.4	4.5
	Never	95.9	95.5	94.5	92.4	91.6	91.5	91.6
	1-5 times	0.3	0.7	1.5	2.1	2.6	2.3	2.1
	6-19 times	0.1	0.3	0.4	0.9	1.0	1.2	1.0
	20-40 times	0.0	0.1	0.2	0.4	0.5	0.3	0.3
	40+ times	0.1	0.2	0.2	0.4	0.4	0.4	0.4
Tranquilizers	No Answer	3.0	3.0	3.1	3.8	3.8	4.4	4.4
	Never	94.3	92.4	90.1	86.6	85.4	85.6	85.8
	1-5 times	2.0	2.9	4.2	5.5	5.9	5.3	5.1
	6-19 times	0.4	1.1	1.6	2.4	2.7	2.8	2.6
	20-40 times	0.1	0.3	0.6	0.9	1.2	1.0	1.1
	40+ times	0.1	0.3	0.5	0.8	0.9	0.9	0.9

General



**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	3.5	3.0	3.1	3.8	3.9	4.4	4.4
	Never	96.0	95.5	93.9	90.9	89.0	88.0	88.0
	1-5 times	0.3	0.8	1.7	2.8	3.7	3.8	3.6
	6-19 times	0.1	0.3	0.7	1.3	1.7	2.0	1.9
	20-40 times	0.0	0.2	0.3	0.7	0.8	0.9	1.1
	40+ times	0.1	0.2	0.2	0.6	0.8	1.0	1.0
Psychedelics (All)	No Answer	3.3	2.9	2.8	3.5	3.6	4.1	4.2
	Never	96.4	96.0	95.1	93.4	92.7	91.7	91.3
	1-5 times	0.2	0.6	1.3	1.8	2.3	2.5	2.8
	6-19 times	0.1	0.2	0.3	0.6	0.7	0.9	0.9
	20-40 times	0.0	0.1	0.2	0.3	0.4	0.4	0.4
	40+ times	0.1	0.2	0.2	0.4	0.4	0.4	0.4
LSD Only	No Answer	3.5	3.2	3.1	3.9	3.9	4.3	4.4
	Never	96.2	96.0	95.2	93.7	93.2	92.7	92.5
	1-5 times	0.2	0.5	1.0	1.5	1.8	1.7	2.0
	6-19 times	0.0	0.1	0.3	0.5	0.5	0.7	0.6
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.3	0.2
	40+ times	0.1	0.1	0.2	0.3	0.3	0.3	0.3
Other Psychedelics	No Answer	3.9	3.4	3.2	3.8	4.0	4.3	4.5
	Never	95.9	96.1	95.7	94.6	94.1	93.1	92.3
	1-5 times	0.1	0.3	0.7	0.9	1.1	1.6	2.0
	6-19 times	0.0	0.1	0.2	0.3	0.5	0.6	0.7
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.2	0.3
	40+ times	0.0	0.1	0.1	0.2	0.2	0.2	0.3
Heroin	No Answer	3.2	2.9	3.1	3.7	3.9	4.4	4.5
	Never	96.5	96.3	95.8	94.8	94.8	94.1	94.0
	1-5 times	0.1	0.5	0.6	0.7	0.7	0.8	0.7
	6-19 times	0.1	0.2	0.2	0.3	0.3	0.4	0.3
	20-40 times	0.1	0.1	0.1	0.2	0.2	0.2	0.2
	40+ times	0.0	0.1	0.1	0.2	0.2	0.3	0.2
Steroids	No Answer	2.9	3.0	3.1	3.9	4.0	4.4	4.5
	Never	96.6	96.1	95.8	94.7	94.5	94.3	94.1
	1-5 times	0.3	0.4	0.6	0.7	0.7	0.5	0.6
	6-19 times	0.1	0.2	0.2	0.2	0.4	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.2	0.2
	40+ times	0.1	0.2	0.1	0.3	0.3	0.3	0.3
Injected Drugs	No Answer	3.2	3.3	3.4	4.2	4.1	4.5	4.6
	Never	96.2	95.8	95.4	94.5	94.5	94.2	94.1
	1-5 times	0.3	0.5	0.7	0.7	0.6	0.5	0.4
	6-19 times	0.1	0.2	0.2	0.2	0.3	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.2	0.2
	40+ times	0.1	0.2	0.2	0.3	0.3	0.3	0.3

General



**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	3.9	3.3	3.2	3.9	3.9	4.3	4.5
	Never	95.9	95.8	95.1	93.5	93.1	92.0	92.0
	1-5 times	0.1	0.5	1.1	1.6	1.9	2.3	2.3
	6-19 times	0.1	0.2	0.3	0.5	0.6	0.6	0.5
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.3	0.3
	40+ times	0.0	0.1	0.1	0.2	0.3	0.3	0.3
Rohypnol	No Answer	4.0	3.4	3.3	3.9	3.9	4.4	4.5
	Never	95.8	96.1	95.9	94.8	94.7	94.4	94.5
	1-5 times	0.1	0.3	0.5	0.7	0.7	0.6	0.5
	6-19 times	0.0	0.1	0.2	0.3	0.3	0.3	0.2
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.2	0.1
	40+ times	0.0	0.1	0.1	0.2	0.2	0.1	0.2
GHB	No Answer	4.1	3.5	3.3	3.9	3.9	4.4	4.5
	Never	95.7	96.1	96.2	95.3	95.2	94.8	94.7
	1-5 times	0.1	0.2	0.3	0.3	0.4	0.4	0.3
	6-19 times	0.1	0.1	0.1	0.2	0.2	0.2	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.1	0.1	0.1	0.1	0.1	0.2

General



**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	1.4	1.1	1.3	1.1	1.2	1.3	1.2
	None	92.1	87.6	80.7	74.6	67.9	64.0	58.0
	Few times	4.1	6.7	10.1	12.0	15.6	16.6	19.5
	1-5 cigarettes/day	1.4	2.7	3.9	5.5	6.3	7.1	7.8
	1/2 pack/day	0.5	0.8	1.7	3.1	3.9	5.0	5.6
	1 pack/day	0.2	0.5	1.2	2.2	3.2	4.0	5.7
	1 & 1/2 pack/day	0.1	0.2	0.3	0.6	0.8	0.9	1.0
Cigars	2+ pack/day	0.3	0.4	0.7	0.8	1.0	1.1	1.1
	No Answer	3.0	3.3	3.3	4.4	4.2	5.0	4.7
	Never	92.7	89.8	84.3	75.3	68.2	62.6	54.0
	1-5 times	3.0	4.4	7.0	10.8	13.9	15.9	19.8
	6-19 times	0.6	1.1	2.2	4.3	5.8	7.3	9.5
	20-40 times	0.3	0.7	1.2	1.9	3.3	3.6	5.3
	40+ times	0.5	0.8	2.0	3.3	4.6	5.6	6.7
Pipe	No Answer	3.3	3.7	3.9	5.2	5.1	6.2	6.0
	Never	95.4	93.7	91.7	87.3	85.4	83.6	81.5
	1-5 times	0.8	1.7	2.4	3.8	5.0	5.4	6.4
	6-19 times	0.2	0.5	0.7	1.6	1.8	1.9	2.4
	20-40 times	0.1	0.2	0.5	0.7	1.0	1.1	1.3
	40+ times	0.2	0.2	0.7	1.4	1.6	1.9	2.3
	No Answer	3.0	3.2	3.3	4.3	4.1	4.9	4.6
Smokeless Tobacco	Never	94.2	91.1	87.2	80.6	75.8	72.6	69.0
	1-5 times	1.7	3.5	5.0	6.9	7.9	8.9	10.2
	6-19 times	0.5	1.0	1.7	2.3	3.1	3.3	3.9
	20-40 times	0.2	0.5	0.8	1.8	2.1	2.2	2.4
	40+ times	0.4	0.7	2.0	4.2	7.1	8.2	9.9
	No Answer	3.3	3.6	3.7	4.5	4.3	5.1	4.9
	Never	78.3	71.5	60.9	50.6	44.3	39.6	33.4
Alcohol	1-5 times	14.6	17.0	21.5	22.2	21.6	21.2	19.3
	6-19 times	2.5	4.9	7.7	11.3	12.5	13.5	13.5
	20-40 times	0.7	1.6	3.2	5.0	7.4	8.8	10.8
	40+ times	0.6	1.4	3.0	6.3	9.9	11.7	18.1
	No Answer	3.2	3.7	3.8	4.8	4.6	5.4	5.2
	Never	94.0	90.1	83.3	76.1	70.6	67.4	64.1
	1-5 times	1.4	3.4	6.0	7.1	8.5	9.6	10.6
Marijuana	6-19 times	0.6	1.2	2.4	3.4	4.7	4.7	5.1
	20-40 times	0.4	0.6	1.6	2.8	3.4	3.2	3.4
	40+ times	0.4	1.1	2.9	5.9	8.3	9.8	11.7
	No Answer	3.2	3.7	3.8	4.8	4.6	5.4	5.2
	Never	94.0	90.1	83.3	76.1	70.6	67.4	64.1

Male



**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	3.3	3.5	3.9	4.9	4.9	5.8	5.6
	Never	96.2	95.3	94.4	91.6	90.3	87.8	86.5
	1-5 times	0.2	0.6	1.0	1.9	2.7	3.5	4.5
	6-19 times	0.1	0.3	0.3	0.7	1.0	1.3	1.5
	20-40 times	0.1	0.2	0.2	0.4	0.5	0.8	0.9
	40+ times	0.1	0.1	0.2	0.6	0.6	0.7	1.0
Crack	No Answer	3.3	3.5	3.8	4.9	5.0	5.9	5.6
	Never	96.1	95.4	94.6	92.9	92.3	90.9	91.1
	1-5 times	0.3	0.6	0.8	1.0	1.3	1.4	1.6
	6-19 times	0.1	0.3	0.4	0.4	0.7	0.8	0.6
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.4	0.5
	40+ times	0.1	0.2	0.2	0.5	0.4	0.5	0.6
Inhalants	No Answer	3.7	3.6	3.8	4.9	5.0	6.0	5.6
	Never	91.5	90.8	90.0	88.8	88.8	88.8	88.3
	1-5 times	3.3	3.7	3.9	3.7	3.7	3.1	3.5
	6-19 times	0.9	0.9	1.3	1.4	1.1	1.2	1.4
	20-40 times	0.1	0.5	0.4	0.5	0.7	0.4	0.6
	40+ times	0.4	0.5	0.5	0.7	0.6	0.6	0.6
Amphetamines	No Answer	4.4	3.9	3.9	5.0	5.1	6.0	5.6
	Never	95.1	95.1	94.0	90.4	88.7	87.0	86.3
	1-5 times	0.2	0.5	1.1	2.2	2.7	3.1	3.8
	6-19 times	0.1	0.2	0.5	1.2	1.8	2.1	1.9
	20-40 times	0.0	0.1	0.2	0.5	0.9	0.9	1.2
	40+ times	0.1	0.2	0.3	0.7	0.8	1.0	1.2
Methamphetamines	No Answer	4.0	3.8	3.8	5.0	5.1	6.0	5.6
	Never	95.6	95.4	95.0	92.7	92.3	90.9	90.5
	1-5 times	0.2	0.4	0.7	1.2	1.2	1.4	1.8
	6-19 times	0.1	0.1	0.3	0.4	0.6	0.8	0.7
	20-40 times	0.0	0.1	0.1	0.3	0.3	0.4	0.7
	40+ times	0.1	0.2	0.2	0.4	0.4	0.5	0.6
Ritalin	No Answer	4.5	4.1	3.9	5.1	5.2	6.2	5.9
	Never	94.9	94.8	94.1	91.5	90.2	89.5	89.5
	1-5 times	0.3	0.6	1.2	1.8	2.5	2.0	2.5
	6-19 times	0.1	0.2	0.4	0.9	1.1	1.4	1.1
	20-40 times	0.1	0.1	0.2	0.3	0.5	0.4	0.4
	40+ times	0.1	0.2	0.2	0.4	0.5	0.6	0.5
Tranquilizers	No Answer	4.1	3.9	3.9	5.0	5.0	6.1	5.8
	Never	93.5	92.8	91.3	87.7	86.3	85.3	84.1
	1-5 times	1.7	2.1	2.9	3.9	4.4	4.3	5.1
	6-19 times	0.4	0.7	1.1	1.9	2.3	2.5	3.0
	20-40 times	0.1	0.2	0.4	0.7	1.0	1.0	1.1
	40+ times	0.2	0.3	0.4	0.8	0.9	0.8	1.0

Male



**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	4.5	3.9	3.9	5.0	5.2	6.1	5.7
	Never	95.0	94.9	93.5	89.8	87.9	85.5	84.5
	1-5 times	0.3	0.6	1.4	2.7	3.3	4.0	4.4
	6-19 times	0.1	0.3	0.7	1.3	1.9	2.2	2.5
	20-40 times	0.0	0.1	0.3	0.6	0.9	1.0	1.5
	40+ times	0.1	0.2	0.3	0.7	0.9	1.2	1.4
Psychedelics (All)	No Answer	4.1	3.7	3.6	4.7	4.7	5.8	5.5
	Never	95.4	95.2	94.2	91.4	90.4	88.6	88.1
	1-5 times	0.3	0.6	1.2	2.3	2.8	3.2	3.9
	6-19 times	0.1	0.2	0.4	0.7	0.9	1.3	1.3
	20-40 times	0.1	0.1	0.2	0.4	0.5	0.6	0.6
	40+ times	0.1	0.2	0.3	0.5	0.6	0.6	0.6
LSD Only	No Answer	4.5	4.1	4.0	5.2	5.2	6.0	5.8
	Never	95.1	95.0	94.1	91.7	91.0	89.7	89.8
	1-5 times	0.2	0.5	1.1	1.9	2.2	2.2	2.8
	6-19 times	0.1	0.1	0.3	0.5	0.7	1.0	0.9
	20-40 times	0.0	0.1	0.2	0.3	0.4	0.5	0.3
	40+ times	0.1	0.2	0.3	0.4	0.5	0.5	0.4
Other Psychedelics	No Answer	4.8	4.2	4.1	5.1	5.1	6.0	5.9
	Never	95.0	95.2	94.9	92.9	92.3	90.3	89.5
	1-5 times	0.1	0.3	0.6	1.1	1.3	2.2	2.8
	6-19 times	0.1	0.1	0.2	0.4	0.6	0.8	0.9
	20-40 times	0.0	0.1	0.1	0.2	0.3	0.4	0.4
	40+ times	0.1	0.1	0.2	0.3	0.3	0.3	0.5
Heroin	No Answer	4.0	3.8	4.0	5.0	5.1	6.3	5.8
	Never	95.6	95.4	94.9	93.3	93.3	91.6	92.0
	1-5 times	0.1	0.5	0.6	0.7	0.6	1.0	1.0
	6-19 times	0.1	0.1	0.2	0.4	0.4	0.4	0.5
	20-40 times	0.1	0.1	0.1	0.3	0.2	0.3	0.3
	40+ times	0.1	0.1	0.2	0.3	0.3	0.4	0.4
Steroids	No Answer	3.8	3.9	3.9	5.2	5.3	6.2	5.9
	Never	95.5	94.9	94.7	92.9	92.4	91.7	91.8
	1-5 times	0.4	0.5	0.7	0.9	0.8	0.8	1.0
	6-19 times	0.1	0.2	0.3	0.4	0.6	0.5	0.5
	20-40 times	0.0	0.2	0.1	0.3	0.4	0.3	0.4
	40+ times	0.2	0.3	0.2	0.4	0.5	0.5	0.5
Injected Drugs	No Answer	4.3	4.2	4.3	5.7	5.5	6.3	6.0
	Never	95.1	94.9	94.6	92.9	92.8	92.0	92.3
	1-5 times	0.3	0.5	0.6	0.7	0.7	0.5	0.6
	6-19 times	0.1	0.2	0.2	0.2	0.4	0.4	0.5
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.4	0.2
	40+ times	0.1	0.2	0.2	0.4	0.4	0.4	0.5

Male

**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	4.8	4.2	4.0	5.2	5.2	6.2	5.9
	Never	94.9	94.9	94.4	91.8	91.6	89.6	89.7
	1-5 times	0.1	0.6	1.0	1.7	1.9	2.5	2.8
	6-19 times	0.1	0.1	0.3	0.6	0.7	0.7	0.7
	20-40 times	0.0	0.1	0.2	0.3	0.3	0.5	0.5
	40+ times	0.1	0.1	0.2	0.3	0.4	0.4	0.5
Rohypnol	No Answer	5.0	4.3	4.2	5.2	5.1	6.2	5.9
	Never	94.8	95.1	95.0	93.4	93.2	92.3	92.6
	1-5 times	0.1	0.3	0.4	0.7	0.7	0.7	0.6
	6-19 times	0.1	0.1	0.2	0.3	0.5	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.3	0.2	0.3	0.2
	40+ times	0.0	0.1	0.1	0.2	0.3	0.2	0.4
GHB	No Answer	5.0	4.4	4.2	5.3	5.1	6.1	5.9
	Never	94.8	95.1	95.3	93.7	93.7	92.7	93.0
	1-5 times	0.0	0.2	0.3	0.4	0.5	0.5	0.5
	6-19 times	0.1	0.1	0.1	0.2	0.3	0.2	0.2
	20-40 times	0.0	0.0	0.1	0.2	0.2	0.2	0.2
	40+ times	0.0	0.1	0.1	0.2	0.2	0.2	0.2

Male



**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	1.1	0.9	0.9	0.9	0.9	0.7	1.0
	None	93.1	87.0	78.8	73.5	69.8	66.4	66.0
	Few times	4.0	7.7	12.5	13.6	15.2	16.5	16.7
	1-5 cigarettes/day	1.2	2.9	4.5	6.2	7.1	8.2	7.4
	1/2 pack/day	0.2	0.7	1.8	3.2	3.9	4.7	5.0
	1 pack/day	0.1	0.4	1.0	1.7	2.3	2.6	3.0
	1 & 1/2 pack/day	0.0	0.1	0.3	0.5	0.4	0.6	0.3
Cigars	2+ pack/day	0.2	0.2	0.3	0.4	0.4	0.3	0.4
	No Answer	1.4	1.8	2.0	2.3	2.3	2.3	2.4
	Never	96.2	93.7	89.8	87.0	84.2	82.9	80.4
	1-5 times	1.6	2.8	5.0	6.4	8.9	10.1	12.1
	6-19 times	0.4	0.8	1.5	1.5	1.7	2.1	2.3
	20-40 times	0.2	0.3	0.5	0.9	1.0	0.7	1.0
	40+ times	0.2	0.6	1.2	1.7	1.9	1.8	1.8
Pipe	No Answer	1.6	2.2	2.5	3.1	3.2	3.4	3.6
	Never	97.6	96.2	94.9	94.0	93.4	93.2	93.1
	1-5 times	0.6	1.0	1.7	1.7	2.1	2.2	2.0
	6-19 times	0.1	0.2	0.4	0.5	0.5	0.6	0.6
	20-40 times	0.0	0.1	0.3	0.2	0.3	0.2	0.2
	40+ times	0.1	0.2	0.2	0.5	0.4	0.3	0.4
	No Answer	1.3	1.6	1.9	2.2	2.3	2.3	2.4
Smokeless Tobacco	Never	97.6	96.4	94.7	93.4	92.7	93.0	93.1
	1-5 times	0.7	1.3	2.3	2.9	3.2	3.3	3.2
	6-19 times	0.1	0.4	0.5	0.6	0.6	0.6	0.4
	20-40 times	0.1	0.1	0.2	0.3	0.4	0.2	0.2
	40+ times	0.1	0.2	0.3	0.6	0.8	0.6	0.7
	No Answer	1.8	2.1	2.2	2.6	2.7	2.7	2.7
	Never	82.0	71.8	56.6	47.7	41.6	38.7	35.9
Alcohol	1-5 times	13.3	18.2	26.6	28.4	29.4	29.5	28.2
	6-19 times	2.0	5.1	9.0	12.1	14.2	14.9	16.2
	20-40 times	0.5	1.7	3.3	5.1	6.6	8.0	9.1
	40+ times	0.4	1.1	2.4	4.1	5.6	6.3	7.9
	No Answer	1.5	1.9	2.1	2.6	2.7	2.7	2.9
	Never	96.8	93.3	85.3	79.7	74.4	71.4	72.8
	1-5 times	1.1	2.8	6.3	8.5	10.3	11.8	11.2
Marijuana	6-19 times	0.2	0.9	2.6	3.5	4.7	5.5	4.3
	20-40 times	0.2	0.5	1.5	2.2	2.8	3.2	2.8
	40+ times	0.1	0.7	2.2	3.5	5.1	5.4	6.1
	No Answer	1.5	1.9	2.1	2.6	2.7	2.7	2.9
	Never	96.8	93.3	85.3	79.7	74.4	71.4	72.8

Female



**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	1.4	1.9	2.1	2.6	2.6	2.6	2.8
	Never	98.3	97.0	96.0	94.4	93.5	93.0	93.0
	1-5 times	0.1	0.7	1.3	1.9	2.5	2.7	2.5
	6-19 times	0.1	0.2	0.4	0.6	0.7	0.9	0.7
	20-40 times	0.0	0.1	0.2	0.2	0.4	0.4	0.5
	40+ times	0.0	0.1	0.1	0.3	0.3	0.4	0.5
Crack	No Answer	1.6	1.8	2.1	2.5	2.7	2.7	2.7
	Never	98.0	97.0	96.1	95.4	95.4	95.6	95.6
	1-5 times	0.3	0.7	1.1	1.3	1.2	1.2	1.0
	6-19 times	0.1	0.2	0.4	0.4	0.3	0.3	0.3
	20-40 times	0.0	0.1	0.2	0.2	0.2	0.1	0.1
	40+ times	0.0	0.1	0.1	0.2	0.2	0.2	0.2
Inhalants	No Answer	2.2	2.1	2.1	2.5	2.7	2.7	2.7
	Never	93.6	91.0	89.6	88.8	90.5	92.3	94.0
	1-5 times	3.0	4.4	5.4	5.9	4.7	3.5	2.3
	6-19 times	0.8	1.6	1.9	1.8	1.2	1.0	0.5
	20-40 times	0.2	0.5	0.6	0.5	0.5	0.4	0.2
	40+ times	0.2	0.4	0.4	0.4	0.3	0.2	0.2
Amphetamines	No Answer	2.4	2.2	2.2	2.5	2.8	2.6	2.8
	Never	97.2	96.5	94.8	92.0	89.8	89.3	90.4
	1-5 times	0.2	0.8	1.8	3.0	4.0	4.3	3.4
	6-19 times	0.1	0.3	0.7	1.5	2.0	2.1	1.6
	20-40 times	0.0	0.2	0.3	0.5	0.9	1.0	1.1
	40+ times	0.1	0.1	0.1	0.5	0.5	0.7	0.8
Methamphetamines	No Answer	2.1	2.2	2.1	2.6	2.7	2.7	2.9
	Never	97.6	96.9	96.2	95.4	95.1	95.1	94.9
	1-5 times	0.2	0.6	1.1	1.2	1.2	1.1	1.1
	6-19 times	0.0	0.2	0.3	0.4	0.5	0.5	0.5
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.3	0.3
	40+ times	0.0	0.1	0.1	0.2	0.3	0.3	0.3
Ritalin	No Answer	2.6	2.3	2.2	2.6	2.8	2.7	2.8
	Never	97.0	96.3	95.1	93.3	92.9	93.3	94.1
	1-5 times	0.3	0.8	1.8	2.5	2.8	2.5	1.8
	6-19 times	0.1	0.3	0.5	1.0	0.9	1.0	0.8
	20-40 times	0.1	0.2	0.4	0.4	0.4	0.3	0.3
	40+ times	0.0	0.2	0.2	0.3	0.3	0.3	0.3
Tranquilizers	No Answer	2.0	2.1	2.2	2.6	2.7	2.7	2.8
	Never	95.0	92.1	88.9	85.6	84.7	86.0	87.8
	1-5 times	2.3	3.7	5.5	7.0	7.2	6.2	5.1
	6-19 times	0.5	1.4	2.1	3.0	3.0	3.1	2.3
	20-40 times	0.1	0.4	0.8	1.1	1.5	1.0	1.1
	40+ times	0.1	0.3	0.6	0.8	0.8	0.9	0.8

Female



**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	2.6	2.2	2.3	2.5	2.7	2.8	2.8
	Never	97.0	96.2	94.4	92.1	90.2	90.3	91.9
	1-5 times	0.3	1.0	2.1	2.9	4.1	3.5	2.7
	6-19 times	0.1	0.3	0.8	1.3	1.6	1.9	1.3
	20-40 times	0.0	0.2	0.3	0.7	0.8	0.7	0.7
	40+ times	0.0	0.1	0.2	0.5	0.7	0.8	0.6
Psychedelics (All)	No Answer	2.4	2.1	2.0	2.3	2.5	2.5	2.6
	Never	97.3	96.9	96.0	95.4	94.8	94.7	94.8
	1-5 times	0.2	0.7	1.3	1.4	1.8	1.8	1.7
	6-19 times	0.0	0.2	0.3	0.5	0.5	0.5	0.5
	20-40 times	0.0	0.1	0.3	0.2	0.2	0.3	0.2
	40+ times	0.1	0.1	0.1	0.2	0.2	0.2	0.2
LSD Only	No Answer	2.6	2.2	2.2	2.6	2.8	2.6	2.8
	Never	97.2	97.0	96.4	95.7	95.2	95.5	95.5
	1-5 times	0.1	0.5	0.9	1.0	1.3	1.2	1.2
	6-19 times	0.0	0.1	0.3	0.4	0.3	0.3	0.3
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.2	0.1
	40+ times	0.1	0.1	0.1	0.2	0.2	0.2	0.1
Other Psychedelics	No Answer	2.9	2.4	2.3	2.6	2.8	2.7	2.8
	Never	96.9	97.0	96.6	96.2	95.8	95.6	95.5
	1-5 times	0.1	0.4	0.7	0.7	0.9	1.1	1.1
	6-19 times	0.0	0.1	0.2	0.3	0.3	0.4	0.4
	20-40 times	0.0	0.1	0.2	0.1	0.1	0.1	0.1
	40+ times	0.0	0.1	0.0	0.1	0.1	0.1	0.1
Heroin	No Answer	2.4	2.0	2.2	2.5	2.7	2.6	2.8
	Never	97.4	97.2	96.8	96.4	96.2	96.4	96.4
	1-5 times	0.1	0.5	0.5	0.6	0.7	0.5	0.4
	6-19 times	0.0	0.2	0.3	0.3	0.2	0.3	0.2
	20-40 times	0.0	0.1	0.1	0.2	0.1	0.1	0.1
	40+ times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
Steroids	No Answer	2.0	2.1	2.3	2.6	2.7	2.8	2.7
	Never	97.8	97.4	97.0	96.5	96.4	96.8	96.7
	1-5 times	0.2	0.3	0.5	0.5	0.5	0.2	0.2
	6-19 times	0.0	0.1	0.2	0.1	0.2	0.1	0.1
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.1	0.2	0.1	0.1	0.1
Injected Drugs	No Answer	2.1	2.3	2.4	2.7	2.8	2.8	2.9
	Never	97.4	96.7	96.3	96.2	96.1	96.3	96.4
	1-5 times	0.3	0.5	0.7	0.6	0.6	0.5	0.2
	6-19 times	0.1	0.3	0.3	0.2	0.3	0.1	0.1
	20-40 times	0.1	0.1	0.1	0.1	0.1	0.0	0.2
	40+ times	0.1	0.1	0.1	0.2	0.2	0.2	0.1

Female

**Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	2.9	2.3	2.2	2.6	2.6	2.7	2.8
	Never	96.9	96.8	95.9	95.1	94.6	94.2	94.6
	1-5 times	0.1	0.5	1.2	1.5	2.0	2.2	1.8
	6-19 times	0.0	0.2	0.4	0.5	0.5	0.5	0.3
	20-40 times	0.0	0.1	0.2	0.2	0.2	0.2	0.2
	40+ times	0.0	0.1	0.1	0.2	0.2	0.3	0.1
Rohypnol	No Answer	3.0	2.4	2.2	2.7	2.7	2.7	2.8
	Never	96.8	97.1	96.8	96.2	96.2	96.3	96.7
	1-5 times	0.1	0.3	0.6	0.7	0.7	0.6	0.3
	6-19 times	0.0	0.1	0.2	0.2	0.2	0.2	0.1
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.1	0.0	0.1	0.1	0.1	0.0
GHB	No Answer	3.2	2.6	2.3	2.6	2.7	2.8	2.9
	Never	96.7	97.1	97.2	96.9	96.8	96.8	96.8
	1-5 times	0.1	0.2	0.3	0.3	0.3	0.2	0.2
	6-19 times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.0	0.0
	40+ times	0.0	0.0	0.0	0.1	0.0	0.1	0.0

Female



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	1.1	0.9	1.0	0.9	0.9	1.0	1.0
	None	95.6	92.4	87.4	82.6	78.5	75.2	72.2
	Few times	1.9	3.6	5.8	7.0	8.4	9.0	9.9
	1-5 cigarettes/day	0.6	1.7	2.9	4.1	5.1	6.4	6.4
	1/2 pack/day	0.2	0.5	1.3	2.5	3.1	4.0	4.8
	1 pack/day	0.2	0.4	1.0	1.8	2.6	3.1	4.2
	1 & 1/2 pack/day	0.1	0.1	0.2	0.5	0.6	0.7	0.7
	2+ pack/day	0.2	0.3	0.5	0.7	0.7	0.7	0.8
Cigars	No Answer	3.1	3.4	3.5	4.3	4.3	4.7	4.6
	Never	95.1	93.3	90.5	86.8	84.1	82.5	78.8
	1-5 times	1.3	2.1	3.7	5.5	7.1	8.3	10.7
	6-19 times	0.3	0.5	0.9	1.3	2.0	2.0	2.7
	20-40 times	0.1	0.3	0.5	0.7	0.8	0.8	1.0
	40+ times	0.1	0.3	0.8	1.3	1.7	1.7	2.1
Pipe	No Answer	3.2	3.6	3.9	4.7	4.9	5.2	5.2
	Never	96.2	95.1	94.1	92.4	91.6	91.3	90.2
	1-5 times	0.4	0.8	1.4	1.9	2.1	2.1	2.7
	6-19 times	0.1	0.3	0.3	0.6	0.6	0.5	0.9
	20-40 times	0.1	0.1	0.1	0.2	0.3	0.3	0.4
	40+ times	0.1	0.1	0.2	0.3	0.5	0.5	0.6
Smokeless Tobacco	No Answer	3.0	3.3	3.5	4.2	4.3	4.7	4.6
	Never	95.9	94.4	92.6	90.1	87.9	87.3	85.6
	1-5 times	0.8	1.5	2.2	2.9	3.4	3.3	3.8
	6-19 times	0.1	0.4	0.6	1.0	1.3	1.2	1.6
	20-40 times	0.1	0.2	0.4	0.7	1.0	1.1	1.3
	40+ times	0.1	0.2	0.6	1.2	2.2	2.4	3.1
Alcohol	No Answer	3.2	3.6	3.6	4.2	4.4	4.5	4.3
	Never	89.2	83.5	75.5	67.3	62.6	59.7	53.5
	1-5 times	6.4	10.1	16.0	20.7	22.8	24.7	26.2
	6-19 times	0.8	1.9	3.3	5.1	6.7	7.6	10.4
	20-40 times	0.2	0.5	0.9	1.6	2.1	2.1	3.2
	40+ times	0.2	0.3	0.7	1.1	1.4	1.4	2.3
Marijuana	No Answer	3.1	3.7	3.8	4.6	4.5	4.8	4.7
	Never	95.5	92.7	88.1	84.0	80.9	79.5	78.1
	1-5 times	0.8	2.2	4.4	5.7	7.0	7.5	7.4
	6-19 times	0.2	0.7	1.8	2.5	3.2	3.2	3.7
	20-40 times	0.2	0.4	0.9	1.4	2.0	2.1	2.3
	40+ times	0.1	0.3	1.1	1.7	2.5	2.8	3.8

General



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	3.2	3.5	3.8	4.5	4.8	5.0	5.2
	Never	96.5	95.7	95.0	93.8	93.1	92.3	91.8
	1-5 times	0.2	0.5	0.7	1.0	1.4	1.8	1.9
	6-19 times	0.0	0.1	0.3	0.4	0.5	0.5	0.7
	20-40 times	0.0	0.1	0.1	0.2	0.1	0.3	0.3
	40+ times	0.0	0.0	0.1	0.1	0.1	0.2	0.2
Crack	No Answer	3.3	3.7	3.9	4.6	4.8	5.0	5.1
	Never	96.4	95.5	95.1	94.3	94.0	93.7	93.5
	1-5 times	0.2	0.5	0.6	0.7	0.7	0.8	0.8
	6-19 times	0.0	0.2	0.2	0.3	0.3	0.3	0.4
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.1	0.0	0.1	0.1	0.1	0.1
Inhalants	No Answer	3.7	3.9	3.9	4.6	4.8	5.2	5.2
	Never	93.4	92.5	92.0	91.8	92.2	92.5	92.6
	1-5 times	2.2	2.5	3.0	2.8	2.2	1.6	1.6
	6-19 times	0.4	0.7	0.7	0.6	0.5	0.5	0.4
	20-40 times	0.1	0.2	0.2	0.2	0.2	0.1	0.1
	40+ times	0.1	0.2	0.2	0.1	0.1	0.1	0.1
Amphetamines	No Answer	4.4	4.1	4.0	4.7	4.9	5.1	5.2
	Never	95.4	95.1	94.6	92.3	91.4	91.1	90.7
	1-5 times	0.2	0.5	1.0	2.1	2.5	2.4	2.6
	6-19 times	0.0	0.2	0.3	0.6	0.7	0.9	1.0
	20-40 times	0.0	0.1	0.1	0.2	0.3	0.3	0.4
	40+ times	0.1	0.0	0.0	0.1	0.1	0.2	0.2
Methamphetamines	No Answer	4.0	3.8	4.0	4.7	4.8	5.1	5.2
	Never	95.8	95.6	95.1	94.1	94.0	93.4	93.3
	1-5 times	0.1	0.3	0.6	0.8	0.7	0.9	0.9
	6-19 times	0.0	0.2	0.2	0.3	0.3	0.4	0.4
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.0	0.1	0.0	0.1	0.1
Ritalin	No Answer	4.4	4.0	4.0	4.8	4.8	5.1	5.3
	Never	95.3	95.1	94.7	93.2	93.0	92.9	92.9
	1-5 times	0.2	0.5	0.9	1.4	1.4	1.3	1.2
	6-19 times	0.0	0.2	0.3	0.4	0.5	0.5	0.4
	20-40 times	0.0	0.1	0.0	0.1	0.1	0.2	0.1
	40+ times	0.0	0.0	0.0	0.1	0.1	0.1	0.1
Tranquilizers	No Answer	3.9	3.9	4.0	4.7	4.8	5.1	5.3
	Never	94.7	93.3	91.9	89.7	89.3	89.3	89.3
	1-5 times	1.2	2.1	3.0	4.0	3.9	3.9	3.7
	6-19 times	0.2	0.5	0.8	1.1	1.3	1.1	1.1
	20-40 times	0.0	0.1	0.2	0.4	0.4	0.4	0.4
	40+ times	0.0	0.1	0.1	0.2	0.3	0.2	0.2

General



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	4.4	4.0	3.9	4.8	4.8	5.1	5.2
	Never	95.3	95.2	94.2	92.3	91.4	90.7	90.8
	1-5 times	0.2	0.5	1.2	1.9	2.6	2.5	2.5
	6-19 times	0.1	0.2	0.4	0.6	0.8	1.1	0.9
	20-40 times	0.0	0.1	0.1	0.2	0.3	0.4	0.4
	40+ times	0.0	0.0	0.1	0.2	0.1	0.2	0.3
Psychedelics (All)	No Answer	4.1	3.8	3.7	4.4	4.7	4.9	5.0
	Never	95.7	95.5	95.1	93.9	93.5	92.9	92.9
	1-5 times	0.2	0.4	0.7	1.1	1.3	1.4	1.4
	6-19 times	0.0	0.2	0.3	0.3	0.3	0.4	0.4
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.1	0.2
	40+ times	0.0	0.1	0.1	0.1	0.1	0.2	0.1
LSD Only	No Answer	4.4	4.0	4.0	4.7	4.9	5.2	5.3
	Never	95.4	95.4	95.1	94.0	93.5	93.2	93.4
	1-5 times	0.1	0.3	0.6	0.9	1.0	1.0	0.9
	6-19 times	0.0	0.1	0.2	0.2	0.2	0.4	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.1	0.1
	40+ times	0.0	0.1	0.1	0.0	0.1	0.1	0.1
Other Psychedelics	No Answer	4.6	4.3	4.2	4.7	5.0	5.2	5.2
	Never	95.2	95.3	95.2	94.5	94.0	93.5	93.3
	1-5 times	0.1	0.2	0.4	0.5	0.7	0.9	1.1
	6-19 times	0.0	0.1	0.2	0.2	0.2	0.3	0.3
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.0	0.1	0.1	0.1	0.1
Heroin	No Answer	4.0	3.9	4.0	4.8	4.9	5.2	5.2
	Never	95.8	95.6	95.3	94.4	94.2	93.8	93.8
	1-5 times	0.2	0.3	0.4	0.4	0.5	0.6	0.5
	6-19 times	0.0	0.1	0.2	0.2	0.2	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.0	0.0	0.1	0.1	0.1
Steroids	No Answer	3.9	3.9	4.2	4.8	5.0	5.3	5.3
	Never	95.9	95.5	95.1	94.3	93.9	93.8	93.7
	1-5 times	0.2	0.3	0.4	0.5	0.5	0.4	0.4
	6-19 times	0.0	0.1	0.1	0.2	0.2	0.2	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.1	0.1
	40+ times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
Injected Drugs	No Answer	4.1	4.2	4.4	5.1	5.2	5.7	5.6
	Never	95.5	95.2	94.7	94.1	93.9	93.5	93.4
	1-5 times	0.2	0.4	0.5	0.4	0.5	0.5	0.5
	6-19 times	0.1	0.2	0.2	0.2	0.2	0.2	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.1	0.1	0.1	0.1	0.1	0.1

General



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	4.7	4.2	4.1	4.8	5.0	5.2	5.2
	Never	95.1	95.2	94.9	93.8	93.5	93.0	93.2
	1-5 times	0.1	0.4	0.7	1.1	1.1	1.4	1.1
	6-19 times	0.0	0.1	0.2	0.2	0.2	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.2	0.2
	40+ times	0.0	0.1	0.0	0.0	0.0	0.0	0.1
Rohypnol	No Answer	4.8	4.3	4.2	4.7	5.0	5.2	5.3
	Never	95.1	95.4	95.3	94.5	94.2	94.1	94.0
	1-5 times	0.1	0.2	0.4	0.5	0.5	0.4	0.3
	6-19 times	0.0	0.1	0.1	0.2	0.2	0.1	0.2
	20-40 times	0.0	0.0	0.0	0.0	0.1	0.1	0.1
	40+ times	0.0	0.0	0.0	0.0	0.1	0.1	0.0
GHB	No Answer	5.0	4.3	4.2	4.8	4.9	5.3	5.3
	Never	95.0	95.4	95.5	94.7	94.6	94.1	94.2
	1-5 times	0.0	0.2	0.2	0.3	0.3	0.4	0.2
	6-19 times	0.0	0.1	0.1	0.1	0.1	0.1	0.2
	20-40 times	0.0	0.1	0.0	0.1	0.1	0.0	0.1
	40+ times	0.0	0.0	0.0	0.0	0.0	0.1	0.1

General



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	1.2	1.0	1.1	1.0	1.1	1.2	1.2
	None	95.4	92.4	87.9	82.6	77.2	73.7	68.2
	Few times	1.8	3.4	5.3	6.6	8.7	9.6	11.1
	1-5 cigarettes/day	0.6	1.6	2.7	3.8	5.0	5.7	6.8
	1/2 pack/day	0.3	0.6	1.1	2.6	3.1	4.2	5.1
	1 pack/day	0.3	0.4	1.0	1.9	3.0	3.7	5.6
	1 & 1/2 pack/day	0.1	0.2	0.2	0.5	0.8	0.9	0.9
Cigars	2+ pack/day	0.3	0.4	0.6	0.9	1.1	1.0	1.1
	No Answer	3.9	4.4	4.4	5.6	5.5	6.4	5.8
	Never	93.8	91.7	88.7	82.6	77.8	74.4	68.7
	1-5 times	1.7	2.6	4.2	7.3	10.0	12.1	16.1
	6-19 times	0.3	0.7	1.2	2.0	3.2	3.2	4.6
	20-40 times	0.1	0.3	0.6	1.0	1.1	1.3	1.6
	40+ times	0.1	0.4	1.0	1.6	2.3	2.6	3.2
Pipe	No Answer	4.0	4.6	4.7	6.1	6.3	7.0	6.5
	Never	95.1	93.7	92.8	89.8	88.4	87.6	85.9
	1-5 times	0.5	1.1	1.6	2.6	3.2	3.1	4.3
	6-19 times	0.1	0.3	0.4	0.8	0.9	0.8	1.6
	20-40 times	0.1	0.1	0.2	0.3	0.5	0.5	0.6
	40+ times	0.1	0.2	0.3	0.5	0.8	0.9	1.0
	No Answer	3.9	4.3	4.4	5.5	5.5	6.5	5.9
Smokeless Tobacco	Never	94.6	92.5	90.0	85.0	80.9	79.2	76.5
	1-5 times	1.0	2.1	3.1	4.5	5.5	5.3	6.4
	6-19 times	0.2	0.6	0.9	1.7	2.3	2.3	2.9
	20-40 times	0.2	0.3	0.6	1.2	1.7	2.0	2.5
	40+ times	0.1	0.3	1.1	2.1	4.1	4.8	5.8
	No Answer	4.0	4.5	4.5	5.6	5.6	6.0	5.3
	Never	87.9	83.3	76.3	67.1	60.8	57.2	49.0
Alcohol	1-5 times	6.9	9.4	14.3	18.4	20.9	22.9	24.7
	6-19 times	0.9	1.9	3.2	5.7	7.9	8.8	12.8
	20-40 times	0.2	0.5	0.9	1.7	2.6	2.9	4.5
	40+ times	0.2	0.4	0.9	1.5	2.1	2.2	3.7
	No Answer	4.1	4.8	4.6	6.1	5.8	6.4	5.9
	Never	94.2	91.3	87.1	81.9	78.3	76.2	73.6
	1-5 times	1.0	2.3	4.2	5.2	6.5	7.0	7.6
Marijuana	6-19 times	0.3	0.8	1.8	2.8	3.7	3.5	4.6
	20-40 times	0.2	0.3	1.0	1.7	2.4	2.6	2.9
	40+ times	0.2	0.4	1.3	2.4	3.4	4.3	5.3

Male



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	4.2	4.5	4.8	6.0	6.3	6.8	6.4
	Never	95.4	94.6	94.2	92.1	91.1	89.9	89.3
	1-5 times	0.2	0.5	0.6	1.1	1.7	2.1	2.6
	6-19 times	0.1	0.1	0.2	0.4	0.6	0.6	1.0
	20-40 times	0.1	0.1	0.1	0.2	0.2	0.3	0.4
	40+ times	0.0	0.0	0.1	0.2	0.2	0.3	0.3
Crack	No Answer	4.2	4.9	4.9	6.2	6.3	6.8	6.4
	Never	95.5	94.3	94.2	92.6	92.1	91.4	91.8
	1-5 times	0.2	0.5	0.5	0.7	0.9	0.9	0.9
	6-19 times	0.0	0.2	0.2	0.3	0.4	0.5	0.6
	20-40 times	0.1	0.1	0.1	0.1	0.1	0.1	0.2
	40+ times	0.0	0.1	0.1	0.1	0.1	0.2	0.1
Inhalants	No Answer	4.7	4.9	4.9	6.1	6.3	7.1	6.6
	Never	92.4	92.0	91.6	91.0	90.8	90.4	90.7
	1-5 times	2.2	2.2	2.5	2.2	2.0	1.6	1.9
	6-19 times	0.5	0.5	0.6	0.5	0.5	0.6	0.6
	20-40 times	0.1	0.2	0.2	0.2	0.2	0.2	0.2
	40+ times	0.1	0.2	0.2	0.1	0.2	0.1	0.1
Amphetamines	No Answer	5.4	5.1	5.0	6.2	6.4	6.9	6.5
	Never	94.3	94.1	93.9	91.2	90.2	89.4	89.0
	1-5 times	0.1	0.4	0.7	1.6	2.2	2.1	2.8
	6-19 times	0.0	0.2	0.2	0.6	0.8	0.9	1.2
	20-40 times	0.1	0.0	0.1	0.3	0.3	0.3	0.3
	40+ times	0.1	0.0	0.0	0.1	0.2	0.2	0.2
Methamphetamines	No Answer	5.0	4.8	5.1	6.2	6.2	7.0	6.5
	Never	94.7	94.6	94.1	92.5	92.4	91.2	91.5
	1-5 times	0.1	0.3	0.5	0.8	0.8	1.0	1.1
	6-19 times	0.1	0.2	0.2	0.3	0.4	0.5	0.6
	20-40 times	0.0	0.0	0.1	0.1	0.2	0.2	0.2
	40+ times	0.0	0.0	0.0	0.1	0.0	0.2	0.1
Ritalin	No Answer	5.5	5.1	5.0	6.3	6.2	7.0	6.6
	Never	94.3	94.3	94.0	91.9	91.4	90.6	91.2
	1-5 times	0.1	0.3	0.7	1.2	1.6	1.4	1.3
	6-19 times	0.0	0.3	0.2	0.5	0.5	0.6	0.7
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.2	0.1
	40+ times	0.0	0.0	0.0	0.1	0.2	0.1	0.1
Tranquilizers	No Answer	4.9	5.0	5.0	6.2	6.3	7.0	6.7
	Never	94.0	93.0	92.4	89.6	88.8	88.0	87.5
	1-5 times	0.9	1.5	1.8	2.7	3.2	3.4	3.9
	6-19 times	0.2	0.3	0.5	0.9	1.0	1.0	1.3
	20-40 times	0.0	0.1	0.2	0.4	0.4	0.4	0.4
	40+ times	0.1	0.1	0.1	0.2	0.3	0.1	0.2

Male



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	5.3	5.0	4.9	6.4	6.3	7.0	6.6
	Never	94.4	94.2	93.6	90.7	89.7	88.5	88.0
	1-5 times	0.2	0.4	1.0	1.9	2.7	2.5	3.3
	6-19 times	0.1	0.2	0.3	0.6	0.8	1.2	1.2
	20-40 times	0.1	0.0	0.1	0.2	0.3	0.5	0.5
	40+ times	0.0	0.1	0.1	0.2	0.2	0.3	0.4
Psychedelics (All)	No Answer	5.0	4.8	4.6	5.9	6.1	6.7	6.3
	Never	94.7	94.6	94.2	92.0	91.4	90.1	90.6
	1-5 times	0.2	0.3	0.7	1.4	1.7	2.0	2.0
	6-19 times	0.0	0.2	0.2	0.4	0.4	0.7	0.7
	20-40 times	0.0	0.1	0.2	0.3	0.2	0.3	0.2
	40+ times	0.0	0.1	0.1	0.1	0.2	0.2	0.2
LSD Only	No Answer	5.4	5.0	5.0	6.3	6.5	7.1	6.7
	Never	94.4	94.4	94.0	92.1	91.4	90.4	91.3
	1-5 times	0.2	0.3	0.6	1.1	1.4	1.5	1.2
	6-19 times	0.0	0.1	0.2	0.3	0.3	0.6	0.5
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.1	0.2
	40+ times	0.0	0.1	0.1	0.0	0.2	0.2	0.1
Other Psychedelics	No Answer	5.6	5.3	5.2	6.1	6.4	7.1	6.6
	Never	94.2	94.3	94.2	92.7	92.2	90.9	91.2
	1-5 times	0.1	0.2	0.3	0.7	1.0	1.4	1.5
	6-19 times	0.0	0.1	0.1	0.3	0.3	0.4	0.5
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.0	0.1	0.1	0.1	0.1
Heroin	No Answer	5.0	4.9	5.0	6.3	6.4	7.1	6.7
	Never	94.8	94.6	94.3	92.6	92.4	91.5	91.9
	1-5 times	0.1	0.3	0.5	0.6	0.7	0.8	0.7
	6-19 times	0.0	0.2	0.1	0.3	0.2	0.4	0.5
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.2
	40+ times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
Steroids	No Answer	5.0	5.0	5.2	6.4	6.5	7.2	6.7
	Never	94.6	94.2	93.8	92.3	91.8	91.3	91.7
	1-5 times	0.3	0.4	0.5	0.7	0.8	0.6	0.7
	6-19 times	0.0	0.2	0.2	0.3	0.4	0.4	0.5
	20-40 times	0.0	0.1	0.1	0.1	0.3	0.2	0.2
	40+ times	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Injected Drugs	No Answer	5.3	5.3	5.5	6.7	6.9	7.7	7.3
	Never	94.4	94.1	93.7	92.3	91.9	91.1	91.4
	1-5 times	0.2	0.3	0.4	0.5	0.6	0.7	0.7
	6-19 times	0.1	0.2	0.1	0.3	0.3	0.3	0.4
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.1	0.1	0.1	0.2	0.1	0.2

Male

**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	5.7	5.2	5.1	6.3	6.5	7.1	6.5
	Never	94.1	94.1	94.0	92.1	91.8	90.4	91.3
	1-5 times	0.1	0.4	0.6	1.2	1.2	1.7	1.3
	6-19 times	0.0	0.1	0.2	0.3	0.3	0.4	0.5
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.2	0.2
	40+ times	0.0	0.0	0.0	0.0	0.0	0.0	0.1
Rohypnol	No Answer	5.7	5.4	5.3	6.2	6.5	7.1	6.9
	Never	94.1	94.2	94.2	92.9	92.5	91.9	92.1
	1-5 times	0.1	0.2	0.3	0.5	0.7	0.5	0.5
	6-19 times	0.0	0.1	0.1	0.2	0.2	0.3	0.3
	20-40 times		0.1	0.0	0.1	0.1	0.1	0.2
	40+ times		0.0	0.0	0.0	0.1	0.1	0.0
GHB	No Answer	6.0	5.4	5.2	6.3	6.4	7.2	6.8
	Never	93.9	94.2	94.4	93.1	92.9	91.9	92.5
	1-5 times	0.0	0.2	0.3	0.3	0.5	0.5	0.3
	6-19 times	0.0	0.1	0.1	0.2	0.2	0.2	0.3
	20-40 times	0.0	0.1	0.0	0.1	0.1		0.1
	40+ times		0.0	0.0	0.0	0.0	0.1	0.1

Male



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	1.0	0.7	0.7	0.7	0.8	0.7	0.8
	None	95.8	92.5	87.0	82.6	79.7	76.7	76.1
	Few times	2.0	3.9	6.3	7.3	8.1	8.4	8.8
	1-5 cigarettes/day	0.6	1.8	3.1	4.4	5.2	7.0	6.0
	1/2 pack/day	0.1	0.4	1.4	2.4	3.2	3.8	4.6
	1 pack/day	0.2	0.3	0.9	1.7	2.3	2.5	2.8
	1 & 1/2 pack/day	0.1	0.1	0.2	0.4	0.4	0.6	0.5
	2+ pack/day	0.2	0.3	0.3	0.4	0.4	0.3	0.4
Cigars	No Answer	2.2	2.4	2.6	3.0	3.2	3.0	3.2
	Never	96.5	95.1	92.5	90.9	89.9	90.0	89.0
	1-5 times	0.9	1.6	3.3	3.9	4.5	4.8	5.5
	6-19 times	0.2	0.4	0.6	0.8	0.9	0.9	0.8
	20-40 times	0.1	0.2	0.4	0.4	0.5	0.4	0.4
	40+ times	0.1	0.3	0.6	1.0	1.0	0.9	1.0
Pipe	No Answer	2.2	2.6	3.0	3.3	3.6	3.6	3.7
	Never	97.3	96.6	95.4	94.9	94.7	94.6	94.6
	1-5 times	0.4	0.5	1.1	1.2	1.1	1.3	1.1
	6-19 times	0.1	0.2	0.3	0.3	0.3	0.3	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.1	0.2	0.2	0.1	0.2
Smokeless Tobacco	No Answer	2.1	2.3	2.6	3.0	3.2	3.1	3.1
	Never	97.2	96.4	95.3	94.9	94.3	94.7	94.9
	1-5 times	0.5	0.8	1.4	1.5	1.5	1.4	1.3
	6-19 times	0.1	0.3	0.3	0.3	0.3	0.3	0.2
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.2	0.1
	40+ times	0.0	0.1	0.1	0.2	0.4	0.3	0.4
Alcohol	No Answer	2.4	2.7	2.6	2.9	3.3	3.1	3.1
	Never	90.7	83.8	74.7	67.5	64.2	62.1	58.2
	1-5 times	5.9	10.8	17.8	22.8	24.5	26.3	27.7
	6-19 times	0.8	1.9	3.4	4.6	5.6	6.6	8.2
	20-40 times	0.1	0.6	0.9	1.5	1.6	1.3	2.0
	40+ times	0.1	0.2	0.6	0.6	0.8	0.7	0.9
Marijuana	No Answer	2.2	2.6	2.8	3.1	3.3	3.3	3.3
	Never	96.9	94.2	89.1	86.1	83.3	82.5	82.8
	1-5 times	0.7	2.1	4.5	6.2	7.4	8.0	7.2
	6-19 times	0.2	0.5	1.8	2.2	2.7	3.0	2.8
	20-40 times	0.1	0.4	0.8	1.2	1.7	1.7	1.8
	40+ times	0.0	0.3	0.8	1.2	1.7	1.4	2.2

Female



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	2.2	2.5	2.8	3.1	3.3	3.3	3.6
	Never	97.6	96.8	96.0	95.4	95.0	94.5	94.5
	1-5 times	0.1	0.5	0.8	0.9	1.2	1.4	1.2
	6-19 times	0.0	0.1	0.3	0.3	0.3	0.4	0.3
	20-40 times		0.1	0.1	0.2	0.1	0.2	0.2
	40+ times	0.0	0.0	0.0	0.1	0.1	0.0	0.1
Crack	No Answer	2.4	2.6	2.8	3.1	3.4	3.3	3.5
	Never	97.4	96.6	96.1	95.9	95.8	95.9	95.6
	1-5 times	0.2	0.5	0.8	0.6	0.5	0.5	0.6
	6-19 times	0.0	0.2	0.2	0.2	0.3	0.2	0.1
	20-40 times		0.1	0.1	0.1	0.1	0.0	0.0
	40+ times	0.0	0.0	0.0	0.1	0.1	0.0	0.1
Inhalants	No Answer	2.8	2.8	2.9	3.2	3.4	3.4	3.5
	Never	94.4	93.0	92.4	92.6	93.5	94.5	94.9
	1-5 times	2.3	2.9	3.5	3.3	2.3	1.5	1.2
	6-19 times	0.4	0.9	0.8	0.7	0.5	0.4	0.2
	20-40 times	0.1	0.3	0.2	0.2	0.1	0.1	0.0
	40+ times	0.1	0.1	0.2	0.1	0.1	0.0	0.0
Amphetamines	No Answer	3.2	2.9	2.8	3.3	3.4	3.3	3.5
	Never	96.5	96.2	95.4	93.3	92.6	92.6	92.8
	1-5 times	0.2	0.6	1.4	2.5	2.8	2.7	2.4
	6-19 times	0.0	0.2	0.4	0.6	0.7	0.8	0.7
	20-40 times		0.1	0.1	0.2	0.3	0.4	0.4
	40+ times	0.0	0.0	0.0	0.1	0.1	0.2	0.2
Methamphetamines	No Answer	3.0	2.8	2.9	3.2	3.4	3.4	3.5
	Never	96.8	96.7	96.1	95.6	95.7	95.4	95.5
	1-5 times	0.1	0.4	0.7	0.8	0.6	0.8	0.8
	6-19 times	0.0	0.1	0.2	0.3	0.2	0.3	0.1
	20-40 times		0.0	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.0	0.0	0.0	0.1	0.1
Ritalin	No Answer	3.3	3.0	3.0	3.3	3.4	3.4	3.6
	Never	96.4	96.1	95.4	94.5	94.6	95.0	95.0
	1-5 times	0.2	0.6	1.2	1.6	1.3	1.1	1.1
	6-19 times	0.0	0.2	0.3	0.4	0.5	0.3	0.2
	20-40 times	0.0	0.1	0.0	0.1	0.1	0.1	0.1
	40+ times	0.0	0.1	0.1	0.1	0.1	0.0	0.0
Tranquilizers	No Answer	2.8	2.7	2.9	3.2	3.3	3.4	3.6
	Never	95.4	93.6	91.4	89.7	89.8	90.5	91.5
	1-5 times	1.5	2.7	4.1	5.1	4.7	4.3	3.4
	6-19 times	0.2	0.7	1.1	1.4	1.6	1.2	0.9
	20-40 times	0.0	0.2	0.3	0.4	0.3	0.4	0.4
	40+ times	0.0	0.1	0.2	0.2	0.3	0.2	0.1

Female



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	3.4	2.9	2.9	3.2	3.4	3.4	3.5
	Never	96.2	96.2	95.0	93.8	93.1	92.7	94.0
	1-5 times	0.3	0.6	1.4	1.9	2.4	2.5	1.7
	6-19 times	0.1	0.2	0.5	0.6	0.8	0.9	0.5
	20-40 times	0.1	0.1	0.2	0.3	0.3	0.3	0.3
	40+ times	0.0	0.0	0.1	0.2	0.1	0.1	0.1
Psychedelics (All)	No Answer	3.1	2.8	2.7	3.0	3.3	3.2	3.4
	Never	96.7	96.5	96.1	95.8	95.4	95.6	95.6
	1-5 times	0.1	0.4	0.8	0.8	0.9	0.8	0.8
	6-19 times	0.0	0.1	0.3	0.2	0.2	0.2	0.1
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.0	0.1
	40+ times	0.0	0.1	0.1	0.1	0.0	0.1	0.1
LSD Only	No Answer	3.4	3.0	3.0	3.2	3.5	3.4	3.5
	Never	96.5	96.5	96.2	95.9	95.6	95.8	95.8
	1-5 times	0.1	0.3	0.6	0.6	0.6	0.6	0.5
	6-19 times		0.1	0.2	0.2	0.2	0.1	0.1
	20-40 times	0.0	0.1	0.0	0.1	0.1	0.0	0.0
	40+ times	0.0	0.0	0.1	0.0		0.1	0.1
Other Psychedelics	No Answer	3.6	3.2	3.1	3.3	3.5	3.4	3.5
	Never	96.3	96.4	96.3	96.1	95.8	95.9	95.8
	1-5 times	0.1	0.3	0.4	0.4	0.5	0.5	0.6
	6-19 times	0.0	0.1	0.2	0.1	0.1	0.1	0.0
	20-40 times		0.0	0.1		0.1	0.0	0.1
	40+ times	0.0	0.0	0.0	0.1	0.0	0.0	0.0
Heroin	No Answer	3.1	2.8	3.0	3.3	3.5	3.5	3.5
	Never	96.8	96.7	96.4	96.1	96.0	96.0	96.0
	1-5 times	0.1	0.3	0.4	0.3	0.3	0.3	0.3
	6-19 times	0.0	0.1	0.2	0.2	0.1	0.2	0.1
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.0	0.1
	40+ times	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Steroids	No Answer	2.7	2.8	3.0	3.2	3.5	3.6	3.6
	Never	97.1	96.8	96.5	96.3	96.0	96.2	96.1
	1-5 times	0.1	0.2	0.3	0.2	0.3	0.1	0.2
	6-19 times	0.0	0.1	0.1	0.1	0.1	0.0	0.1
	20-40 times		0.0	0.0	0.1	0.1	0.1	0.0
	40+ times	0.0	0.0	0.0	0.1	0.0	0.0	0.1
Injected Drugs	No Answer	3.0	3.0	3.3	3.5	3.6	3.7	3.7
	Never	96.7	96.3	95.8	95.9	95.7	95.8	95.8
	1-5 times	0.2	0.4	0.5	0.4	0.4	0.3	0.2
	6-19 times	0.1	0.2	0.3	0.2	0.1	0.1	0.1
	20-40 times	0.0	0.1	0.0	0.0	0.1	0.0	0.1
	40+ times	0.0		0.1	0.0	0.1	0.0	0.1


Female



**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	3.7	3.0	3.0	3.3	3.5	3.4	3.5
	Never	96.2	96.4	95.9	95.4	95.2	95.4	95.4
	1-5 times	0.1	0.4	0.8	1.0	1.0	1.0	0.9
	6-19 times		0.1	0.2	0.1	0.2	0.1	0.1
	20-40 times		0.1	0.1	0.1	0.0	0.1	0.0
	40+ times	0.0	0.0	0.0	0.0	0.0	0.0	0.1
Rohypnol	No Answer	3.7	3.2	3.0	3.2	3.6	3.4	3.5
	Never	96.2	96.5	96.4	96.1	95.8	96.1	96.3
	1-5 times	0.1	0.2	0.5	0.5	0.4	0.3	0.2
	6-19 times	0.0	0.0	0.1	0.1	0.1	0.0	0.1
	20-40 times		0.0	0.1	0.0	0.0	0.1	0.0
	40+ times		0.0	0.0	0.0	0.0	0.0	0.0
GHB	No Answer	3.9	3.2	3.1	3.3	3.5	3.5	3.5
	Never	96.0	96.6	96.6	96.4	96.2	96.2	96.3
	1-5 times	0.1	0.1	0.2	0.2	0.2	0.2	0.1
	6-19 times	0.0	0.0	0.1	0.1	0.1	0.0	0.1
	20-40 times		0.0	0.0	0.0	0.0	0.0	0.0
	40+ times		0.0	0.0	0.0	0.0	0.0	0.0

Female



New books, videos and curricula on

- Addiction counseling
- Prevention planning
- Screening and brief intervention
- Medically assisted treatment protocols
- Gambling prevention

Visit our library at: <http://www.drugs.indiana.edu>



**Binge Drinking in the Past Two Weeks
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Binge drinking	No Answer	25.4	22.8	21.4	17.8	15.6	14.0	12.9
	None	70.4	70.2	67.6	66.5	64.5	64.2	59.8
	Once	2.3	3.4	5.5	6.9	8.2	9.1	10.1
	Twice	0.9	1.7	2.4	3.8	5.0	5.2	6.5
	3-5 times	0.6	1.1	1.7	2.8	3.9	4.4	6.7
	6-9 times	0.1	0.4	0.6	0.8	1.3	1.4	1.9
	10+ times	0.2	0.5	0.8	1.3	1.5	1.7	2.1

General

**Binge Drinking in the Past Two Weeks
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Binge drinking	No Answer	26.7	25.4	23.8	20.0	18.0	16.3	14.8
	None	68.8	67.7	65.7	64.1	60.5	58.6	53.0
	Once	2.5	3.3	5.1	6.4	7.8	9.3	10.0
	Twice	0.9	1.7	2.2	3.6	5.4	6.0	7.8
	3-5 times	0.7	1.0	1.5	3.0	4.5	5.3	8.5
	6-9 times	0.1	0.3	0.7	0.8	1.7	1.8	2.7
	10+ times	0.3	0.5	1.1	2.0	2.1	2.7	3.3

Male

**Binge Drinking in the Past Two Weeks
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Binge drinking	No Answer	24.1	20.1	18.9	15.6	13.3	11.8	10.7
	None	72.0	72.8	69.6	69.0	68.2	69.4	66.8
	Once	2.1	3.4	5.8	7.3	8.4	8.8	10.3
	Twice	1.0	1.8	2.7	4.0	4.7	4.5	5.3
	3-5 times	0.5	1.1	1.8	2.6	3.5	3.6	5.0
	6-9 times	0.2	0.4	0.6	0.9	1.0	1.0	1.0
	10+ times	0.2	0.4	0.6	0.6	0.8	0.8	0.9

Female



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	2.4	2.3	2.3	2.3	2.1	2.1	2.0
	Never	85.7	78.1	67.6	61.5	55.0	51.2	47.2
	7 years or less	2.4	2.3	2.8	2.5	2.5	1.9	1.7
	8-9 years	2.2	2.8	3.5	3.5	3.1	2.7	2.4
	10-11 years	4.2	5.5	6.0	5.1	4.5	4.1	3.4
	12-13 years	2.9	8.2	13.0	12.4	11.1	9.3	8.3
	14-15 years	0.0	0.7	4.7	12.0	16.6	14.8	12.9
	16-17 years	0.0	0.0	0.1	0.6	5.1	13.6	17.6
	Over 18 years	0.1	0.1	0.1	0.1	0.1	0.4	4.4
Cigars	No Answer	1.6	1.8	2.1	2.0	2.0	2.2	2.3
	Never	93.3	90.2	83.6	78.0	71.8	67.8	61.0
	7 years	1.2	1.2	1.5	1.3	1.4	1.2	1.1
	8-9 years	1.0	1.1	1.4	1.6	1.3	1.0	1.0
	10-11 years	1.7	2.1	2.7	2.4	2.3	1.7	1.5
	12-13 years	1.1	3.3	6.0	6.3	5.5	4.3	3.7
	14-15 years	0.0	0.4	2.5	7.9	11.6	10.4	8.7
	16-17 years	0.0	0.0	0.0	0.4	4.0	11.0	16.1
	Over 18 years	0.0	0.0	0.1	0.0	0.1	0.3	4.5
Pipe	No Answer	1.9	2.3	2.9	3.4	3.5	3.8	4.2
	Never	96.6	95.1	92.5	90.1	88.1	86.9	85.0
	7 years	0.4	0.4	0.5	0.5	0.7	0.5	0.6
	8-9 years	0.3	0.3	0.4	0.5	0.5	0.4	0.4
	10-11 years	0.5	0.7	0.9	0.9	0.8	0.7	0.5
	12-13 years	0.3	1.1	1.7	2.1	2.0	1.7	1.4
	14-15 years	0.0	0.1	0.9	2.3	3.2	2.9	2.4
	16-17 years	0.0		0.0	0.1	1.1	3.0	4.2
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	1.3
Smokeless Tobacco	No Answer	1.5	1.7	2.0	2.0	2.2	2.2	2.3
	Never	95.8	93.3	89.6	86.2	82.2	80.4	77.1
	7 years	0.6	0.8	1.0	1.0	1.2	1.1	1.1
	8-9 years	0.5	0.6	0.9	0.9	0.9	0.7	0.8
	10-11 years	0.9	1.1	1.5	1.4	1.4	1.2	1.2
	12-13 years	0.6	2.3	3.3	3.4	3.1	2.4	2.8
	14-15 years	0.0	0.2	1.7	4.7	6.8	5.5	4.8
	16-17 years	0.0		0.0	0.3	2.2	6.2	8.1
	Over 18 years	0.1	0.0	0.1	0.0	0.1	0.2	1.9

General



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Alcohol	No Answer	2.7	2.8	3.4	2.8	3.0	2.9	3.0
	Never	73.8	65.8	52.3	43.4	36.5	32.2	28.6
	7 years	5.3	4.6	4.5	3.8	3.7	3.1	3.1
	8-9 years	4.1	3.8	3.7	3.0	2.3	1.9	1.5
	10-11 years	8.2	7.7	7.5	6.0	4.9	3.7	3.0
	12-13 years	5.8	14.3	18.4	16.9	14.3	11.2	9.2
	14-15 years	0.0	0.9	10.2	23.1	27.0	24.9	21.4
	16-17 years	0.0	0.0	0.0	1.0	8.2	19.8	25.6
	Over 18 years	0.1	0.1	0.1	0.1	0.1	0.5	4.7
Marijuana	No Answer	1.7	1.9	2.4	2.5	2.7	2.7	3.0
	Never	95.5	91.4	82.4	75.6	68.1	63.8	60.9
	7 years	0.4	0.6	1.0	1.0	1.0	1.0	0.9
	8-9 years	0.4	0.5	0.8	0.9	0.8	0.7	0.6
	10-11 years	0.9	1.6	2.2	2.2	2.1	1.9	1.2
	12-13 years	1.0	3.7	6.9	7.5	7.4	6.1	5.4
	14-15 years	0.0	0.3	4.1	9.9	13.9	12.9	11.8
	16-17 years				0.5	3.8	10.7	13.9
	Over 18 years	0.0	0.0	0.1	0.0	0.1	0.2	2.2
Cocaine	No Answer	1.7	1.8	2.2	2.3	2.6	2.7	3.0
	Never	97.8	96.8	95.5	93.4	91.8	89.9	88.8
	7 years	0.1	0.2	0.2	0.2	0.3	0.3	0.3
	8-9 years	0.1	0.1	0.1	0.2	0.2	0.1	0.1
	10-11 years	0.1	0.3	0.3	0.3	0.2	0.3	0.2
	12-13 years	0.1	0.8	1.0	1.1	0.8	0.7	0.6
	14-15 years	0.0	0.1	0.8	2.3	2.7	2.2	1.4
	16-17 years	0.0	0.0	0.0	0.2	1.5	3.8	4.3
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.1	1.3
Crack	No Answer	1.8	1.9	2.2	2.4	2.6	2.8	3.0
	Never	97.6	96.6	95.8	94.8	94.2	93.7	93.2
	7 years	0.2	0.2	0.2	0.2	0.3	0.3	0.3
	8-9 years	0.1	0.1	0.2	0.2	0.1	0.2	0.1
	10-11 years	0.2	0.3	0.3	0.2	0.2	0.2	0.2
	12-13 years	0.2	0.8	0.8	0.8	0.5	0.4	0.3
	14-15 years	0.0	0.1	0.5	1.3	1.3	1.0	0.6
	16-17 years	0.0	0.0	0.0	0.1	0.7	1.4	1.8
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	0.5

General



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Inhalants	No Answer	2.4	2.1	2.3	2.4	2.7	2.8	3.0
	Never	91.7	89.8	88.2	87.1	87.1	87.9	87.9
	7 years	1.5	1.3	1.0	0.8	0.7	0.5	0.6
	8-9 years	1.0	1.1	0.9	0.8	0.6	0.5	0.3
	10-11 years	1.9	2.3	2.0	1.6	1.2	1.0	0.8
	12-13 years	1.3	3.2	4.1	3.7	2.9	2.2	1.6
	14-15 years	0.0	0.2	1.5	3.5	3.8	3.0	2.6
	16-17 years	0.0	0.0	0.0	0.1	1.0	2.0	2.6
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	0.5
Amphetamines	No Answer	2.8	2.2	2.3	2.5	2.7	2.7	3.1
	Never	96.7	96.4	94.7	91.3	88.7	86.9	86.4
	7 years	0.1	0.2	0.2	0.2	0.4	0.3	0.4
	8-9 years	0.1	0.1	0.1	0.2	0.2	0.1	0.1
	10-11 years	0.1	0.3	0.4	0.4	0.4	0.4	0.2
	12-13 years	0.1	0.8	1.4	2.2	1.9	1.6	1.1
	14-15 years	0.0	0.1	0.8	2.9	4.2	4.2	3.2
	16-17 years	0.0	0.0	0.0	0.2	1.5	3.6	4.8
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.1	0.7
Methamphetamines	No Answer	2.3	2.0	2.2	2.5	2.7	2.8	3.1
	Never	97.3	96.9	96.0	94.7	93.8	93.2	92.4
	7 years	0.1	0.1	0.2	0.2	0.2	0.2	0.2
	8-9 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.1	0.2	0.2	0.3	0.2	0.2	0.1
	12-13 years	0.1	0.6	0.7	0.7	0.5	0.6	0.4
	14-15 years	0.0	0.1	0.6	1.4	1.7	1.4	1.0
	16-17 years	0.0	0.0		0.1	0.6	1.6	2.1
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.0	0.6
Ritalin	No Answer	2.9	2.2	2.4	2.5	2.8	2.9	3.1
	Never	96.4	96.3	94.7	92.4	91.0	90.8	90.4
	7 years	0.2	0.2	0.2	0.3	0.3	0.3	0.3
	8-9 years	0.1	0.2	0.2	0.3	0.2	0.2	0.1
	10-11 years	0.2	0.3	0.5	0.5	0.4	0.4	0.3
	12-13 years	0.2	0.8	1.2	1.7	1.2	1.0	0.8
	14-15 years	0.0	0.0	0.7	2.1	2.9	2.5	2.0
	16-17 years	0.0	0.0	0.0	0.1	1.0	1.9	2.7
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	0.4

General



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Tranquilizers	No Answer	2.3	2.1	2.4	2.5	2.8	2.9	3.1
	Never	94.2	92.5	89.5	86.1	83.8	83.7	83.5
	7 years	0.6	0.6	0.5	0.5	0.6	0.4	0.4
	8-9 years	0.5	0.5	0.4	0.4	0.3	0.1	0.1
	10-11 years	1.2	1.1	1.1	1.0	0.7	0.6	0.4
	12-13 years	1.1	3.1	3.6	3.6	2.7	1.9	1.5
	14-15 years	0.0	0.2	2.3	5.6	6.5	5.6	3.8
	16-17 years	0.0		0.0	0.3	2.5	4.7	6.2
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	1.0
Narcotics	No Answer	2.8	2.2	2.3	2.6	2.8	3.0	3.3
	Never	96.6	96.0	94.1	90.7	88.1	86.7	86.0
	7 years	0.2	0.2	0.2	0.2	0.4	0.3	0.4
	8-9 years	0.1	0.1	0.2	0.2	0.1	0.1	0.1
	10-11 years	0.2	0.3	0.4	0.6	0.4	0.4	0.2
	12-13 years	0.2	0.9	1.6	2.0	1.7	1.3	0.8
	14-15 years	0.0	0.1	1.2	3.4	4.6	4.2	3.1
	16-17 years	0.0		0.0	0.2	1.8	4.0	5.3
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	0.8
LSD Only	No Answer	2.9	2.2	2.2	2.5	2.8	2.9	3.1
	Never	96.7	96.7	95.7	94.3	93.2	92.6	92.0
	7 years	0.1	0.2	0.2	0.2	0.3	0.2	0.3
	8-9 years	0.0	0.1	0.1	0.2	0.1	0.1	0.1
	10-11 years	0.1	0.2	0.2	0.2	0.2	0.3	0.1
	12-13 years	0.1	0.5	0.8	0.8	0.7	0.6	0.4
	14-15 years	0.0	0.1	0.7	1.7	1.8	1.5	1.0
	16-17 years	0.0		0.0	0.1	1.0	1.8	2.3
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.1	0.6
Other Psychedelics	No Answer	3.2	2.4	2.3	2.6	2.8	2.9	3.1
	Never	96.5	96.9	96.5	95.3	94.6	93.5	92.6
	7 years	0.1	0.1	0.1	0.2	0.2	0.2	0.2
	8-9 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.1	0.1	0.2	0.2	0.2	0.2	0.1
	12-13 years	0.1	0.4	0.5	0.5	0.4	0.4	0.2
	14-15 years	0.0	0.0	0.4	0.9	1.2	1.2	0.9
	16-17 years	0.0			0.1	0.6	1.3	2.1
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.5

General



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Heroin	No Answer	2.5	2.0	2.3	2.5	2.8	2.9	3.1
	Never	97.1	97.0	96.4	95.7	95.3	94.9	94.7
	7 years	0.1	0.2	0.2	0.1	0.2	0.2	0.3
	8-9 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.1	0.2	0.1	0.2	0.1	0.2	0.1
	12-13 years	0.1	0.5	0.5	0.4	0.3	0.3	0.2
	14-15 years	0.0	0.1	0.4	0.8	0.7	0.5	0.4
	16-17 years	0.0	0.0	0.0	0.1	0.4	0.8	0.7
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.1	0.3
Steroids	No Answer	2.2	2.2	2.3	2.5	2.8	2.8	3.2
	Never	97.1	96.8	96.3	95.7	95.1	95.4	94.7
	7 years	0.2	0.2	0.2	0.2	0.3	0.2	0.3
	8-9 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.2	0.2	0.2	0.2	0.2	0.2	0.1
	12-13 years	0.2	0.5	0.4	0.5	0.3	0.2	0.3
	14-15 years	0.0	0.1	0.4	0.7	0.7	0.4	0.3
	16-17 years	0.0		0.0	0.1	0.6	0.7	0.8
	Over 18 years	0.0	0.0	0.1	0.0	0.0	0.0	0.3
Injected Drugs	No Answer	2.5	2.4	2.5	3.0	3.0	3.1	3.3
	Never	96.7	96.4	96.0	95.4	95.1	95.0	94.7
	7 years	0.3	0.2	0.2	0.2	0.2	0.2	0.3
	8-9 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.2	0.2	0.2	0.2	0.1	0.2	0.1
	12-13 years	0.1	0.6	0.5	0.4	0.3	0.3	0.2
	14-15 years	0.0	0.1	0.4	0.7	0.6	0.5	0.4
	16-17 years	0.0	0.0	0.0	0.1	0.4	0.7	0.5
	Over 18 years	0.0	0.0	0.1	0.1	0.0	0.0	0.3
MDMA	No Answer	3.2	2.3	2.3	2.5	2.8	3.0	3.2
	Never	96.5	96.6	95.6	94.0	92.9	91.9	91.8
	7 years	0.1	0.2	0.2	0.2	0.2	0.2	0.3
	8-9 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.1	0.2	0.2	0.3	0.2	0.2	0.1
	12-13 years	0.1	0.7	0.8	0.9	0.7	0.6	0.3
	14-15 years	0.0	0.0	0.7	1.9	1.9	1.5	1.0
	16-17 years	0.0	0.0	0.0	0.2	1.2	2.6	2.4
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	0.9

General



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Rohypnol	No Answer	3.3	2.4	2.3	2.5	2.7	2.9	3.1
	Never	96.4	96.9	96.6	95.7	95.4	95.2	95.3
	7 years	0.1	0.1	0.1	0.1	0.2	0.2	0.2
	8-9 years	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.1	0.1	0.1	0.2	0.1	0.2	0.1
	12-13 years	0.1	0.4	0.4	0.5	0.3	0.3	0.3
	14-15 years	0.0	0.0	0.3	0.6	0.7	0.5	0.2
	16-17 years	0.0		0.0	0.1	0.4	0.5	0.5
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.1	0.2
GHB	No Answer	3.3	2.5	2.4	2.5	2.8	2.8	3.2
	Never	96.4	97.0	97.0	96.5	96.0	95.9	95.6
	7 years	0.1	0.1	0.1	0.1	0.2	0.2	0.2
	8-9 years	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	12-13 years	0.0	0.2	0.2	0.2	0.2	0.2	0.2
	14-15 years	0.0	0.0	0.2	0.3	0.4	0.3	0.2
	16-17 years	0.0		0.0	0.0	0.2	0.3	0.3
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.0	0.2



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	2.7	2.4	2.4	2.5	2.3	2.2	2.3
	Never	84.3	77.6	67.8	61.2	53.9	50.6	43.4
	7 years or less	2.9	2.9	3.2	3.1	3.0	2.3	2.2
	8-9 years	2.5	3.1	3.8	3.9	3.6	2.9	2.8
	10-11 years	4.5	5.6	6.2	5.4	5.0	4.3	3.9
	12-13 years	3.0	7.6	11.9	11.8	10.9	9.2	8.8
	14-15 years	0.0	0.7	4.5	11.2	15.3	14.0	13.2
	16-17 years	0.0	0.0	0.1	0.7	5.8	14.0	18.5
	Over 18 years	0.1	0.1	0.1	0.1	0.1	0.6	4.9
Cigars	No Answer	2.0	2.4	2.5	2.4	2.6	2.7	2.8
	Never	91.4	87.8	79.6	71.3	62.3	55.7	47.1
	7 years	1.5	1.7	1.9	1.8	1.9	1.7	1.5
	8-9 years	1.4	1.3	1.9	2.3	1.9	1.6	1.4
	10-11 years	2.2	2.5	3.4	3.2	3.4	2.5	2.4
	12-13 years	1.4	3.9	7.5	8.2	7.3	6.3	5.4
	14-15 years		0.4	3.2	10.1	14.9	14.5	12.4
	16-17 years	0.0		0.0	0.6	5.5	14.4	21.1
	Over 18 years	0.0	0.1	0.1	0.0	0.2	0.5	6.0
Pipe	No Answer	2.3	2.9	3.4	3.9	4.0	4.5	4.6
	Never	95.7	93.6	90.7	86.7	83.2	80.8	78.1
	7 years	0.5	0.6	0.7	0.7	0.9	0.7	0.8
	8-9 years	0.4	0.4	0.6	0.8	0.9	0.8	0.7
	10-11 years	0.7	0.9	1.3	1.4	1.2	1.2	0.8
	12-13 years	0.4	1.5	2.3	2.9	3.2	2.6	2.3
	14-15 years	0.0	0.1	1.0	3.3	4.8	4.6	3.9
	16-17 years	0.0			0.2	1.7	4.5	6.6
	Over 18 years	0.0	0.0	0.1	0.0	0.1	0.2	2.1
Smokeless Tobacco	No Answer	2.0	2.3	2.4	2.6	2.8	2.9	2.8
	Never	94.2	90.2	85.1	79.3	72.1	68.6	63.1
	7 years	0.8	1.1	1.5	1.6	1.9	1.9	1.7
	8-9 years	0.8	0.9	1.2	1.4	1.5	1.2	1.4
	10-11 years	1.3	1.6	2.1	2.2	2.2	2.1	2.0
	12-13 years	0.9	3.5	4.9	5.1	5.1	4.0	4.8
	14-15 years	0.0	0.3	2.7	7.3	10.7	8.8	8.1
	16-17 years	0.0		0.0	0.5	3.6	10.1	13.1
	Over 18 years	0.1	0.0	0.1	0.0	0.1	0.4	2.8

Male



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Alcohol	No Answer	3.0	3.4	3.9	3.1	3.6	3.4	3.4
	Never	71.7	65.1	53.6	44.7	37.9	33.6	27.8
	7 years	5.6	5.1	5.0	4.5	4.0	3.6	3.6
	8-9 years	4.6	4.1	3.8	3.3	2.5	2.3	1.9
	10-11 years	8.9	8.0	7.5	6.7	5.6	4.0	3.5
	12-13 years	6.1	13.1	16.3	15.6	13.7	11.2	9.5
	14-15 years	0.0	1.1	9.7	20.8	24.1	22.3	20.5
	16-17 years		0.0	0.1	1.2	8.4	19.0	24.8
	Over 18 years	0.1	0.1	0.1	0.1	0.2	0.6	4.9
	No Answer	2.3	2.5	2.9	2.9	3.4	3.3	3.5
Marijuana	Never	94.4	89.7	81.2	73.7	65.8	61.7	56.3
	7 years	0.5	0.7	1.3	1.3	1.3	1.3	1.0
	8-9 years	0.5	0.6	0.9	1.2	1.1	1.0	0.8
	10-11 years	1.0	1.8	2.6	2.7	2.7	2.4	1.7
	12-13 years	1.3	4.1	7.0	7.9	7.9	7.1	6.4
	14-15 years	0.0	0.5	4.1	9.6	13.6	12.7	12.9
	16-17 years				0.6	4.0	10.3	14.5
	Over 18 years	0.1	0.1	0.1	0.1	0.1	0.3	2.8
	No Answer	2.2	2.5	2.7	2.9	3.4	3.5	3.8
	Never	97.3	96.0	95.0	92.5	90.0	87.4	85.6
Cocaine	7 years	0.1	0.3	0.3	0.3	0.5	0.5	0.5
	8-9 years	0.1	0.1	0.1	0.2	0.2	0.1	0.2
	10-11 years	0.1	0.3	0.3	0.3	0.3	0.5	0.4
	12-13 years	0.1	0.7	0.9	1.2	0.9	0.8	0.8
	14-15 years	0.0	0.1	0.7	2.4	2.7	2.3	1.6
	16-17 years		0.0	0.0	0.2	1.9	4.7	5.3
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	1.9
	No Answer	2.3	2.6	2.7	3.0	3.4	3.6	3.9
	Never	97.0	95.8	95.4	94.0	92.8	91.9	91.0
	7 years	0.2	0.3	0.3	0.4	0.4	0.3	0.4
Crack	8-9 years	0.1	0.2	0.2	0.2	0.2	0.2	0.2
	10-11 years	0.2	0.3	0.2	0.3	0.2	0.3	0.2
	12-13 years	0.1	0.7	0.7	0.7	0.5	0.5	0.5
	14-15 years	0.0	0.1	0.5	1.4	1.4	1.2	0.9
	16-17 years	0.0	0.0	0.0	0.1	0.9	1.8	2.3
	Over 18 years	0.0	0.0	0.1	0.0	0.1	0.1	0.6

Male



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Inhalants	No Answer	2.9	2.7	2.8	2.9	3.5	3.8	3.8
	Never	90.7	89.5	88.4	87.4	86.3	86.8	84.8
	7 years	1.7	1.5	1.2	1.0	0.9	0.7	0.8
	8-9 years	1.1	1.1	1.0	1.0	0.7	0.7	0.5
	10-11 years	2.2	2.1	1.8	1.6	1.3	1.1	1.0
	12-13 years	1.4	2.8	3.2	3.0	2.8	2.0	2.0
	14-15 years		0.2	1.5	2.9	3.3	2.8	3.0
	16-17 years		0.0	0.0	0.2	1.1	2.1	3.3
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	0.7
Amphetamines	No Answer	3.4	2.9	2.7	3.0	3.5	3.6	4.0
	Never	96.1	95.7	94.7	91.1	88.4	86.5	84.5
	7 years	0.1	0.2	0.2	0.3	0.5	0.4	0.4
	8-9 years	0.1	0.1	0.2	0.4	0.3	0.2	0.2
	10-11 years	0.1	0.3	0.3	0.5	0.5	0.5	0.3
	12-13 years	0.1	0.6	1.2	1.8	1.8	1.4	1.2
	14-15 years	0.0	0.1	0.7	2.7	3.6	3.8	3.0
	16-17 years	0.0	0.0		0.2	1.4	3.6	5.5
	Over 18 years	0.0	0.0	0.1	0.0	0.1	0.1	1.0
Methamphetamines	No Answer	2.8	2.6	2.7	3.0	3.5	3.6	3.9
	Never	96.8	96.3	95.6	94.0	92.8	91.7	90.7
	7 years	0.1	0.2	0.2	0.2	0.4	0.3	0.3
	8-9 years	0.1	0.1	0.1	0.2	0.1	0.2	0.2
	10-11 years	0.1	0.2	0.2	0.4	0.2	0.3	0.1
	12-13 years	0.1	0.5	0.6	0.6	0.5	0.6	0.6
	14-15 years	0.0	0.1	0.5	1.4	1.7	1.5	1.1
	16-17 years		0.0		0.2	0.7	1.7	2.4
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.0	0.7
Ritalin	No Answer	3.5	2.8	2.9	3.0	3.6	3.8	3.9
	Never	95.7	95.7	94.5	92.0	89.8	89.4	88.2
	7 years	0.2	0.3	0.3	0.5	0.5	0.4	0.3
	8-9 years	0.1	0.2	0.2	0.3	0.3	0.3	0.2
	10-11 years	0.2	0.2	0.5	0.6	0.5	0.6	0.4
	12-13 years	0.2	0.7	0.9	1.4	1.3	1.0	1.0
	14-15 years	0.0	0.1	0.6	2.0	2.9	2.4	2.2
	16-17 years	0.0	0.0	0.0	0.1	1.1	2.1	3.1
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	0.5

Male



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Tranquilizers	No Answer	2.9	2.7	2.8	3.0	3.6	3.7	3.9
	Never	93.8	93.1	91.2	87.9	84.9	84.1	81.9
	7 years	0.6	0.6	0.5	0.5	0.7	0.4	0.5
	8-9 years	0.5	0.5	0.4	0.5	0.4	0.2	0.2
	10-11 years	1.3	0.7	0.9	1.0	0.7	0.7	0.4
	12-13 years	0.9	2.2	2.6	2.8	2.2	1.5	1.4
	14-15 years	0.0	0.2	1.7	4.1	5.2	4.9	3.7
	16-17 years			0.0	0.3	2.2	4.3	6.7
	Over 18 years	0.0	0.0	0.1	0.0	0.1	0.1	1.3
Narcotics	No Answer	3.2	2.9	2.7	3.2	3.6	3.8	4.2
	Never	96.1	95.4	93.9	90.0	87.1	84.8	82.5
	7 years	0.2	0.3	0.3	0.3	0.5	0.3	0.5
	8-9 years	0.1	0.1	0.2	0.2	0.2	0.2	0.2
	10-11 years	0.2	0.3	0.4	0.6	0.6	0.6	0.3
	12-13 years	0.2	0.8	1.4	1.9	1.7	1.5	1.0
	14-15 years	0.0	0.1	1.1	3.3	4.2	4.3	3.7
	16-17 years	0.0		0.0	0.3	2.1	4.3	6.5
	Over 18 years	0.0	0.0	0.1	0.0	0.1	0.1	1.3
LSD Only	No Answer	3.5	2.9	2.7	3.1	3.6	3.8	3.9
	Never	96.0	95.9	94.9	92.9	90.8	90.0	89.2
	7 years	0.2	0.2	0.3	0.2	0.5	0.3	0.5
	8-9 years	0.1	0.1	0.2	0.3	0.2	0.1	0.1
	10-11 years	0.1	0.2	0.3	0.3	0.3	0.5	0.2
	12-13 years	0.1	0.6	0.8	1.0	0.8	0.7	0.7
	14-15 years	0.0	0.1	0.8	2.1	2.4	1.8	1.2
	16-17 years				0.2	1.3	2.6	3.4
	Over 18 years	0.1	0.0	0.0	0.0	0.1	0.1	0.8
Other Psychedelics	No Answer	3.7	3.1	2.7	3.2	3.6	3.9	4.1
	Never	96.0	96.2	96.0	94.3	92.8	91.1	89.7
	7 years	0.1	0.1	0.2	0.2	0.3	0.3	0.3
	8-9 years	0.0	0.1	0.1	0.1	0.1	0.2	0.2
	10-11 years	0.1	0.1	0.1	0.3	0.2	0.2	0.2
	12-13 years	0.1	0.3	0.5	0.6	0.4	0.6	0.3
	14-15 years	0.0	0.1	0.3	1.2	1.5	1.7	1.3
	16-17 years				0.1	0.8	1.9	3.2
	Over 18 years	0.0	0.0	0.1	0.0	0.1	0.1	0.7

Male



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Heroin	No Answer	3.0	2.7	2.7	3.1	3.7	3.8	4.1
	Never	96.6	96.2	95.8	94.8	94.0	93.1	92.9
	7 years	0.1	0.2	0.2	0.2	0.3	0.3	0.3
	8-9 years	0.1	0.1	0.1	0.2	0.2	0.2	0.1
	10-11 years	0.1	0.1	0.2	0.3	0.2	0.3	0.1
	12-13 years	0.1	0.5	0.6	0.5	0.3	0.4	0.4
	14-15 years	0.0	0.1	0.4	0.9	0.7	0.7	0.6
	16-17 years	0.0		0.0	0.0	0.6	1.1	1.0
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	0.4
Steroids	No Answer	2.8	2.9	2.9	3.3	3.7	3.8	4.2
	Never	96.2	95.6	95.3	94.2	93.1	93.1	92.5
	7 years	0.2	0.2	0.3	0.3	0.4	0.3	0.4
	8-9 years	0.1	0.1	0.2	0.2	0.2	0.2	0.1
	10-11 years	0.3	0.3	0.2	0.2	0.2	0.3	0.2
	12-13 years	0.3	0.7	0.5	0.7	0.4	0.3	0.4
	14-15 years	0.0	0.1	0.5	0.9	0.9	0.7	0.5
	16-17 years	0.0			0.2	1.0	1.3	1.3
	Over 18 years	0.1	0.0	0.1	0.0	0.1	0.1	0.5
Injected Drugs	No Answer	3.3	3.1	3.1	3.8	4.0	3.9	4.3
	Never	95.9	95.7	95.5	94.3	93.6	93.3	93.0
	7 years	0.4	0.3	0.3	0.2	0.4	0.3	0.5
	8-9 years	0.1	0.1	0.1	0.1	0.1	0.2	0.1
	10-11 years	0.1	0.2	0.2	0.2	0.2	0.2	0.1
	12-13 years	0.1	0.6	0.5	0.4	0.3	0.5	0.3
	14-15 years	0.0	0.1	0.3	0.8	0.7	0.7	0.6
	16-17 years	0.0	0.0	0.0	0.1	0.5	0.9	0.7
	Over 18 years	0.0	0.0	0.1	0.1	0.1	0.1	0.4
MDMA	No Answer	3.7	3.0	2.8	3.1	3.8	3.9	4.2
	Never	95.9	95.7	95.1	93.1	91.5	90.0	89.5
	7 years	0.0	0.3	0.2	0.3	0.3	0.3	0.3
	8-9 years	0.1	0.1	0.2	0.2	0.1	0.1	0.2
	10-11 years	0.1	0.2	0.2	0.3	0.2	0.3	0.1
	12-13 years	0.1	0.6	0.8	0.9	0.6	0.6	0.4
	14-15 years	0.0	0.0	0.6	2.0	2.0	1.7	1.0
	16-17 years	0.0		0.0	0.2	1.4	2.9	3.1
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.0	1.2

Male



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Rohypnol	No Answer	3.8	3.1	2.9	3.3	3.5	4.0	3.9
	Never	95.8	96.1	96.1	94.8	94.1	93.7	93.8
	7 years	0.1	0.1	0.2	0.2	0.3	0.2	0.3
	8-9 years	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.1	0.1	0.1	0.3	0.2	0.4	0.1
	12-13 years	0.1	0.4	0.4	0.5	0.4	0.3	0.4
	14-15 years	0.0	0.1	0.2	0.7	0.8	0.5	0.4
	16-17 years	0.0			0.1	0.5	0.6	0.7
	Over 18 years	0.0	0.0	0.0	0.0	0.1	0.1	0.2
GHB	No Answer	3.9	3.2	2.8	3.2	3.7	3.6	4.0
	Never	95.9	96.2	96.4	95.6	94.6	94.4	94.2
	7 years	0.1	0.1	0.2	0.2	0.3	0.3	0.3
	8-9 years	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.0	0.1	0.1	0.1	0.2	0.2	0.1
	12-13 years	0.0	0.3	0.2	0.3	0.2	0.2	0.3
	14-15 years	0.0	0.0	0.2	0.4	0.6	0.5	0.4
	16-17 years				0.1	0.4	0.5	0.5
	Over 18 years	0.0	0.0	0.1	0.0	0.1	0.1	0.3

Male



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	2.2	2.1	2.2	2.0	1.9	2.0	1.7
	Never	87.1	78.8	67.6	61.8	56.0	51.9	51.1
	7 years or less	2.0	1.8	2.3	2.0	2.0	1.5	1.2
	8-9 years	2.0	2.6	3.1	3.1	2.7	2.5	2.0
	10-11 years	3.9	5.3	5.8	4.9	4.0	3.8	2.8
	12-13 years	2.8	8.8	14.0	12.9	11.2	9.3	7.7
	14-15 years	0.1	0.6	5.0	12.7	17.7	15.5	12.6
	16-17 years	0.0		0.0	0.5	4.3	13.3	16.8
	Over 18 years	0.1	0.0	0.1	0.0	0.0	0.3	4.0
Cigars	No Answer	1.2	1.1	1.7	1.6	1.5	1.7	1.7
	Never	95.1	92.7	87.7	84.3	80.5	78.8	74.9
	7 years	0.9	0.7	1.1	0.8	0.8	0.7	0.7
	8-9 years	0.7	0.9	1.0	0.9	0.8	0.4	0.7
	10-11 years	1.2	1.6	2.0	1.7	1.4	1.0	0.6
	12-13 years	0.8	2.7	4.5	4.6	3.8	2.5	2.0
	14-15 years	0.0	0.3	1.9	5.8	8.6	6.9	5.1
	16-17 years	0.0	0.0	0.0	0.3	2.6	8.1	11.2
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.2	3.1
Pipe	No Answer	1.4	1.6	2.4	2.9	3.1	3.1	3.5
	Never	97.5	96.7	94.4	93.4	92.6	92.6	92.1
	7 years	0.3	0.2	0.3	0.2	0.4	0.3	0.4
	8-9 years	0.2	0.2	0.3	0.3	0.2	0.1	0.2
	10-11 years	0.3	0.5	0.6	0.4	0.4	0.3	0.2
	12-13 years	0.2	0.7	1.2	1.3	1.0	0.7	0.5
	14-15 years	0.0	0.0	0.7	1.4	1.8	1.4	1.0
	16-17 years			0.0	0.1	0.6	1.6	1.9
	Over 18 years	0.0		0.0	0.0	0.0		0.4
Smokeless Tobacco	No Answer	1.1	1.0	1.5	1.5	1.6	1.6	1.6
	Never	97.5	96.4	94.2	92.7	91.4	91.1	91.0
	7 years	0.4	0.4	0.4	0.4	0.5	0.4	0.4
	8-9 years	0.2	0.3	0.5	0.4	0.4	0.2	0.2
	10-11 years	0.5	0.6	0.8	0.7	0.6	0.5	0.3
	12-13 years	0.3	1.1	1.8	1.8	1.3	0.9	0.7
	14-15 years	0.0	0.1	0.7	2.3	3.3	2.6	1.5
	16-17 years	0.0			0.0	0.9	2.7	3.2
	Over 18 years	0.0	0.0	0.0	0.1	0.0	0.0	0.9

Female



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Alcohol	No Answer	2.4	2.1	2.9	2.6	2.5	2.4	2.5
	Never	75.9	66.6	51.1	42.1	35.3	30.9	29.3
	7 years	5.0	4.0	4.0	3.0	3.4	2.6	2.6
	8-9 years	3.6	3.6	3.5	2.7	2.0	1.5	1.1
	10-11 years	7.5	7.3	7.4	5.4	4.3	3.5	2.5
	12-13 years	5.5	15.6	20.4	18.1	14.9	11.2	8.9
	14-15 years	0.0	0.8	10.7	25.3	29.6	27.1	22.2
	16-17 years	0.0		0.0	0.8	7.9	20.5	26.4
	Over 18 years	0.0	0.1	0.1	0.0	0.1	0.4	4.4
	No Answer	1.2	1.2	1.9	2.1	2.1	2.1	2.4
Marijuana	Never	96.7	93.2	83.8	77.4	70.4	65.8	65.7
	7 years	0.3	0.3	0.7	0.6	0.7	0.7	0.7
	8-9 years	0.3	0.4	0.6	0.6	0.6	0.4	0.3
	10-11 years	0.7	1.3	1.9	1.6	1.4	1.4	0.8
	12-13 years	0.8	3.3	6.9	7.0	6.9	5.4	4.3
	14-15 years	0.0	0.2	4.1	10.2	14.2	13.0	10.7
	16-17 years				0.4	3.7	11.1	13.4
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.2	1.7
	No Answer	1.2	1.1	1.6	1.7	1.8	1.8	2.0
	Never	98.4	97.6	96.1	94.3	93.4	92.1	92.2
Cocaine	7 years	0.1	0.1	0.2	0.1	0.2	0.1	0.1
	8-9 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	10-11 years	0.1	0.3	0.3	0.2	0.1	0.1	0.1
	12-13 years	0.1	0.8	1.0	1.1	0.7	0.6	0.4
	14-15 years		0.0	0.8	2.2	2.6	2.1	1.1
	16-17 years	0.0			0.1	1.1	3.0	3.4
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.7
	No Answer	1.2	1.1	1.7	1.8	1.9	2.0	2.0
	Never	98.2	97.3	96.2	95.6	95.5	95.5	95.7
	7 years	0.2	0.2	0.1	0.1	0.2	0.2	0.1
Crack	8-9 years	0.1	0.0	0.1	0.1	0.1	0.1	0.0
	10-11 years	0.1	0.4	0.4	0.2	0.1	0.1	0.1
	12-13 years	0.2	0.8	0.9	0.8	0.5	0.4	0.1
	14-15 years		0.1	0.6	1.2	1.1	0.7	0.4
	16-17 years				0.1	0.6	1.1	1.3
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.3

Female



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Inhalants	No Answer	1.9	1.4	1.7	1.8	2.0	1.9	2.0
	Never	92.8	90.2	88.1	86.8	87.9	89.0	91.3
	7 years	1.4	1.1	0.8	0.6	0.5	0.4	0.3
	8-9 years	0.9	1.1	0.8	0.5	0.5	0.3	0.1
	10-11 years	1.7	2.5	2.1	1.6	1.0	0.9	0.6
	12-13 years	1.2	3.6	4.9	4.3	3.0	2.4	1.2
	14-15 years	0.0	0.1	1.5	4.2	4.2	3.1	2.2
	16-17 years			0.0	0.1	0.9	1.9	2.0
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.3
Amphetamines	No Answer	2.2	1.4	1.8	2.0	1.9	1.9	2.2
	Never	97.3	97.0	94.8	91.4	89.1	87.3	88.4
	7 years	0.1	0.1	0.2	0.2	0.2	0.2	0.3
	8-9 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	10-11 years	0.1	0.3	0.5	0.4	0.3	0.3	0.2
	12-13 years	0.1	0.9	1.6	2.6	2.0	1.8	1.1
	14-15 years			0.0	0.9	3.1	4.8	4.5
	16-17 years			0.0	0.2	1.5	3.7	4.2
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.1	0.4
Methamphetamines	No Answer	1.8	1.3	1.7	1.9	2.0	1.9	2.1
	Never	97.9	97.6	96.3	95.3	94.8	94.6	94.4
	7 years	0.1	0.1	0.2	0.1	0.1	0.1	0.1
	8-9 years	0.1	0.1	0.1	0.1	0.1	0.0	0.0
	10-11 years	0.1	0.3	0.2	0.3	0.1	0.1	0.1
	12-13 years	0.1	0.6	0.8	0.8	0.5	0.5	0.2
	14-15 years			0.1	0.6	1.3	1.8	1.3
	16-17 years				0.1	0.6	1.4	1.7
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.4
Ritalin	No Answer	2.2	1.5	1.9	2.0	2.1	2.1	2.1
	Never	97.0	96.9	95.0	92.8	92.2	92.1	92.8
	7 years	0.2	0.1	0.1	0.2	0.2	0.1	0.2
	8-9 years	0.1	0.1	0.2	0.2	0.1	0.1	0.1
	10-11 years	0.2	0.4	0.5	0.4	0.4	0.2	0.2
	12-13 years	0.1	0.9	1.5	2.0	1.1	1.0	0.5
	14-15 years	0.0	0.0	0.8	2.2	3.0	2.5	1.7
	16-17 years			0.0	0.1	0.9	1.8	2.2
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.3

Female



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Tranquilizers	No Answer	1.8	1.4	1.9	2.1	2.0	2.2	2.2
	Never	94.6	91.8	88.0	84.3	82.9	83.5	85.4
	7 years	0.5	0.5	0.5	0.5	0.5	0.4	0.4
	8-9 years	0.5	0.4	0.5	0.3	0.2	0.1	0.1
	10-11 years	1.1	1.5	1.4	1.0	0.7	0.5	0.3
	12-13 years	1.3	4.0	4.7	4.5	3.2	2.2	1.5
	14-15 years	0.0	0.3	2.9	6.9	7.6	6.2	3.8
	16-17 years	0.0		0.0	0.3	2.8	4.9	5.7
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.1	0.8
Narcotics	No Answer	2.3	1.5	1.8	2.1	2.0	2.1	2.2
	Never	97.1	96.7	94.3	91.2	89.1	88.5	89.7
	7 years	0.2	0.1	0.2	0.2	0.3	0.2	0.3
	8-9 years	0.0	0.2	0.2	0.1	0.1	0.1	0.1
	10-11 years	0.2	0.4	0.5	0.5	0.2	0.2	0.2
	12-13 years	0.2	1.0	1.7	2.1	1.7	1.0	0.6
	14-15 years		0.1	1.3	3.5	5.0	4.1	2.5
	16-17 years				0.2	1.5	3.6	4.3
	Over 18 years	0.1	0.0	0.0	0.1	0.0	0.1	0.3
LSD Only	No Answer	2.2	1.4	1.7	1.9	1.9	2.0	2.0
	Never	97.5	97.7	96.6	95.7	95.3	95.1	95.1
	7 years	0.1	0.1	0.1	0.1	0.1	0.1	0.2
	8-9 years	0.0	0.1	0.1	0.1	0.0	0.0	
	10-11 years	0.1	0.3	0.2	0.2	0.1	0.1	0.1
	12-13 years	0.1	0.5	0.7	0.7	0.6	0.4	0.2
	14-15 years	0.0	0.1	0.6	1.3	1.2	1.2	0.8
	16-17 years	0.0		0.0	0.0	0.6	1.1	1.3
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.3
Other Psychedelics	No Answer	2.6	1.6	1.8	2.0	2.0	2.0	2.0
	Never	97.1	97.6	97.0	96.4	96.2	95.8	95.8
	7 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	8-9 years	0.1	0.1	0.1	0.0	0.1	0.0	0.1
	10-11 years	0.1	0.2	0.2	0.2	0.1	0.1	0.1
	12-13 years	0.1	0.4	0.5	0.5	0.4	0.3	0.1
	14-15 years		0.0	0.4	0.7	0.9	0.8	0.5
	16-17 years				0.1	0.3	0.9	1.0
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.3

Female



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Heroin	No Answer	2.0	1.2	1.7	1.9	2.0	2.0	2.0
	Never	97.7	97.8	97.1	96.6	96.5	96.6	96.9
	7 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	8-9 years	0.0	0.1	0.1	0.1	0.1	0.0	0.1
	10-11 years	0.1	0.2	0.1	0.2	0.1	0.1	0.1
	12-13 years	0.1	0.5	0.5	0.3	0.3	0.2	0.0
	14-15 years		0.0	0.4	0.7	0.7	0.4	0.2
	16-17 years		0.0	0.0	0.1	0.3	0.6	0.4
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.2
Steroids	No Answer	1.6	1.4	1.7	1.8	1.9	1.9	2.1
	Never	98.1	97.9	97.4	97.1	97.0	97.5	97.2
	7 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	8-9 years	0.0	0.1	0.1	0.1	0.1		0.0
	10-11 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	12-13 years	0.1	0.3	0.3	0.3	0.2	0.1	0.2
	14-15 years		0.1	0.3	0.4	0.4	0.2	0.1
	16-17 years			0.0	0.0	0.1	0.1	0.2
	Over 18 years	0.0	0.0	0.0	0.0	0.0		0.1
Injected Drugs	No Answer	1.8	1.6	1.9	2.1	2.0	2.3	2.2
	Never	97.5	97.3	96.5	96.5	96.5	96.5	96.7
	7 years	0.3	0.2	0.2	0.1	0.1	0.1	0.1
	8-9 years	0.1	0.1	0.1	0.1	0.0	0.0	0.1
	10-11 years	0.2	0.2	0.2	0.2	0.1	0.1	0.1
	12-13 years	0.1	0.6	0.5	0.3	0.3	0.2	0.1
	14-15 years	0.0	0.1	0.6	0.6	0.5	0.3	0.3
	16-17 years			0.0	0.1	0.3	0.4	0.3
	Over 18 years	0.0	0.0	0.0	0.1	0.0	0.0	0.2
MDMA	No Answer	2.6	1.5	1.8	2.0	1.9	2.1	2.1
	Never	97.1	97.4	96.1	94.9	94.3	93.6	94.3
	7 years	0.1	0.1	0.2	0.2	0.1	0.1	0.2
	8-9 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	10-11 years	0.1	0.2	0.2	0.2	0.1	0.1	0.0
	12-13 years	0.1	0.7	0.9	0.8	0.7	0.5	0.2
	14-15 years	0.0	0.0	0.8	1.7	1.8	1.3	0.9
	16-17 years		0.0	0.0	0.2	1.0	2.3	1.6
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.6

Female



**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Rohypnol	No Answer	2.7	1.6	1.7	1.9	1.9	1.9	2.1
	Never	97.0	97.7	97.2	96.6	96.6	96.7	97.1
	7 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	8-9 years	0.1	0.1	0.1	0.0	0.1	0.0	0.0
	10-11 years	0.1	0.2	0.2	0.2	0.1	0.1	0.1
	12-13 years	0.0	0.4	0.4	0.5	0.3	0.3	0.2
	14-15 years		0.0	0.3	0.6	0.6	0.5	0.1
	16-17 years			0.0	0.1	0.2	0.4	0.3
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.1
GHB	No Answer	2.8	1.8	1.8	1.9	2.0	2.1	2.1
	Never	97.0	97.8	97.6	97.4	97.3	97.4	97.4
	7 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	8-9 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	10-11 years	0.1	0.1	0.1	0.1	0.0	0.0	0.1
	12-13 years	0.0	0.2	0.1	0.2	0.2	0.1	0.1
	14-15 years			0.1	0.3	0.2	0.2	0.0
	16-17 years			0.0		0.1	0.1	0.2
	Over 18 years	0.0	0.0	0.0	0.0	0.0	0.0	0.1

Female

Afternoons

Recreational SKILLS LESSONS Culture & Values KNOWLEDGE
Recreational SKILLS LESSONS Culture & Values KNOWLEDGE

R O C K

in Indiana


Working toward a healthier future



Afternoons R.O.C.K. in Indiana delivers evidence-based drug abuse prevention programs in the after-school hours.


Youth benefit from structured recreation as well as academic and social skills development in a safe and supervised environment.

To start an Afternoons ROCK in Indiana program in your area, contact the Indiana Prevention Resource Center at 800-346-3077 or via the web at www.ROCK.indiana.edu






Monthly Use of Selected Drugs and Binge Drinking by Indiana and U.S.A. 12th Graders



Source: Indiana Prevention Resource Center at Indiana University, 2006
Johnston, et al., National Institute on Drug Abuse, 2005


Alcohol Use among Indiana Students 6th - 12th Graders, 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006




Monthly Alcohol Use among Indiana Students by Grade and Gender, 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006


Binge Drinking among Indiana Students by Grade and Gender, 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006




Cigarette Use among Indiana Students 6th - 12th Graders, 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006


Monthly Cigarette Use among Indiana Students by Grade and Gender, 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006




Smokeless Tobacco Use among Indiana Students 6th - 12th Graders, 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006


Monthly Smokeless Tobacco Use among Indiana Students by Grade and Gender, 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006




Marijuana Use among Indiana Students 6th - 12th Graders, 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006


Monthly Marijuana Use among Indiana Students by Grade and Gender, 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006



Inhalant Use among Indiana Students 6th - 12th Graders, 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006

**OVER 8,000 RETAIL OUTLET INSPECTIONS
CONDUCTED IN 2005.**

OVER 87% DID NOT SELL TOBACCO TO MINORS.



Tobacco Retailer
Inspection Program

To report a vendor suspected of selling tobacco products to youth, contact the Indiana State Excise Police at 1.866.2STOPEM (1.866.278.6736).

Major Robin Poindexter
TRIP Project Director
Indiana State Excise Police
Phone: 1.317.233.0838
Web: www.trip.indiana.edu


Desiree D. Goetze, MPH, CHES, CPP
TRIP Operations Coordinator
Indiana Prevention Resource Center
Phone: 812-855-1237
Email: trip@indiana.edu



Compliance Checks Change Indiana




Trends of Annual Alcohol Use Among Indiana Students: 1991-2006



Source: Indiana Prevention Resource Center at Indiana University, 2006


Trends of Monthly Alcohol Use Among Indiana Students: 1991- 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006




Trend of Binge Drinking among Indiana Students: 1991-2006



Source: Indiana Prevention Resource Center at Indiana University, 2006


Trend of Binge Drinking by Indiana and U.S.A. 12th Graders: 1975-2006



Source: Indiana Prevention Resource Center at Indiana University, 2006
Johnston, et al., National Institute on Drug Abuse, 2005




Trends of Annual Cigarette Use Among Indiana Students: 1991- 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006


Trends of Monthly Cigarette Use Among Indiana Students: 1991-2006



Source: Indiana Prevention Resource Center at Indiana University, 2006




Trend of Monthly Cigarette Use by Indiana and U.S.A. 12th Graders: 1975-2006



Source: Indiana Prevention Resource Center at Indiana University, 2006
Johnston, et al., National Institute on Drug Abuse, 2005


Trends of Annual Smokeless Tobacco Use Among Indiana Students: 1991- 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006




Trends of Monthly Smokeless Tobacco Use Among Indiana Students: 1991- 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006


Trends of Annual Marijuana Use Among Indiana Students: 1991- 2006



Source: Indiana Prevention Resource Center at Indiana University, 2006




Trend of Monthly Marijuana Use among Indiana Students: 1991-2006



Source: Indiana Prevention Resource Center at Indiana University , 2006


Trend of Monthly Marijuana Use by Indiana and U.S.A. 12th Graders: 1975-2006



Source: Indiana Prevention Resource Center at Indiana University, 2006
Johnston, et al., National Institute on Drug Abuse, 2005




Trend of Annual Inhalent Use among Indiana Students: 1991-2006




Source: Indiana Prevention Resource Center at Indiana University, 2006

Trend of Monthly Inhalant Use among Indiana Students: 1991-2006



Source: Indiana Prevention Resource Center at Indiana University, 2006



Understanding your community is the first step to addressing public health problems. PREV-STAT community analysis reports offer market segmentation data, risk and protective factors and other insights into behavior that allows for precision prevention planning and grant writing.

Contact: For your community profile, visit
www.drugs.indiana.edu or call 800-346-3077.

II. PREVALENCE RATES AND FREQUENCIES

2. REGIONAL DATA

- Prevalence Rates by Grade:
Lifetime, Annual, Monthly, Daily and Special Use



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	11.9	11.4	11.7	8.7 ▼	12.8	10.6 ▼	16.1 ▲	9.9 ▼	14.5 ▲
Cigars	5.4	5.0	5.2	3.7 ▼	6.4 ▲	4.7 ▼	6.6 ▲	4.7	6.6 ▲
Pipe	1.8	2.6 ▲	1.4	1.6	2.1	1.4 ▼	2.2	1.6	1.9
Smokeless Tobacco	2.7	2.1	2.0	1.6 ▼	3.7 ▲	1.7 ▼	3.9 ▲	3.6 ▲	3.7 ▲
Alcohol	25.2	30.4 ▲	22.8 ▼	19.5 ▼	26.2	21.8 ▼	25.7	29.0 ▲	28.8 ▲
Marijuana	2.9	3.1	2.3	1.5 ▼	2.5	3.9 ▲	3.3	1.3 ▼	2.6
Cocaine	0.6	0.7	0.7	0.6	0.7 ▲	0.6	0.7	0.4	0.4
Crack	0.8	0.9	0.7	0.9	1.2 ▲	0.6	1.1	0.6	0.5
Inhalants	6.5	8.7 ▲	5.4	5.1	6.6	5.5 ▼	8.6 ▲	5.7	7.3
Amphetamines	0.5	0.6	0.3	0.3	0.8	0.5	1.0 ▲	0.5	0.4
Methamphetamines	0.5	0.8	0.3	0.3	0.6	0.4	0.8	0.3	0.4
Ritalin	0.9	1.3	0.9	0.6	0.8	0.8	1.7 ▲	0.6	0.7
Tranquilizers	3.9	4.5	3.8	3.6	4.1	3.7	3.6	3.3	4.2
Narcotics	0.8	0.9	0.3 ▼	0.8	0.8	0.7	1.7 ▲	0.4	0.8
Psychedelics [#]	0.6	0.9	0.3	0.4	0.8	0.7	0.7	0.5	0.5
LSD	0.5	0.6	0.2	0.3	0.6	0.5	0.6	0.2	0.3
Others	0.4	0.7	0.1	0.3	0.5	0.5	0.3	0.4	0.2
Heroin	0.5	0.7	0.4	0.5	0.6	0.5	0.6	0.2	0.3
Steroids	0.8	0.9	0.3 ▼	0.5	1.1	0.6	1.1	0.6	0.8
Injected Drugs	0.9	1.2	0.9	0.5	1.3	0.8	0.9	1.0	0.4 ▼
MDMA	0.4	0.6	0.4	0.3	0.4	0.4	0.5	0.1	0.2
Rohypnol	0.4	0.6	0.1	0.3	0.5	0.4	0.4	0.3	0.2
GHB	0.3	0.4	0.1	0.3	0.4	0.3	0.3	0.1	0.2

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	6.2	6.1	6.0	5.4	6.6	5.5	8.2 ▲	4.8 ▼	7.1
Cigars	3.4	3.5	3.7	2.3	3.6	3.1	4.1	2.9	4.0
Pipe	1.1	1.1	0.8	1.1	1.4	0.9	1.4	1.3	1.0
Smokeless Tobacco	1.9	1.7	1.6	1.3	2.2	1.4 ▼	2.9 ▲	2.4	2.7 ▲
Alcohol	17.3	22.7 ▲	15.4 ▼	13.0 ▼	17.3	15.1 ▼	16.8	19.4	19.7 ▲
Marijuana	2.3	2.5	2.0	1.1 ▼	1.7	3.0 ▲	3.2 ▲	0.8 ▼	2.1
Cocaine	0.4	0.6	0.5	0.4	0.4	0.4	0.6	0.1	0.2
Crack	0.5	0.6	0.5	0.6	0.6	0.4	0.9 ▲	0.4	0.3
Inhalants	4.5	6.0 ▲	3.5	3.2	4.9	3.7 ▼	5.1	3.9	5.6 ▲
Amphetamines	0.4	0.4	0.2	0.3	0.5	0.4	0.7	0.5	0.3
Methamphetamines	0.3	0.5	0.3	0.3	0.3	0.2	0.5	0.2	0.2
Ritalin	0.5	0.7	0.5	0.3	0.4	0.5	0.9 ▲	0.2	0.5
Tranquilizers	2.7	3.5 ▲	2.5	2.5	2.6	2.7	2.7	2.1	2.8
Narcotics	0.5	0.7	0.2	0.5	0.6	0.4	0.9 ▲	0.3	0.4
Psychedelics [#]	0.4	0.5	0.2	0.2	0.5	0.5	0.3	0.3	0.3
LSD	0.3	0.4	0.1	0.2	0.4	0.4	0.3	0.2	0.2
Others	0.2	0.3	0.2	0.1	0.3	0.2	0.2	0.2	0.1
Heroin	0.3	0.5	0.3	0.3	0.4	0.3	0.4	0.2	0.2
Steroids	0.5	0.6	0.2	0.4	0.8	0.4	0.8	0.4	0.5
Injected Drugs	0.5	0.8	0.5	0.4	0.8	0.5	0.6	0.7	0.3
MDMA	0.3	0.4	0.3	0.2	0.3	0.2	0.5	0.1	0.1
Rohypnol	0.2	0.3	0.1	0.3	0.4	0.2	0.3	0.2	0.1
GHB	0.1	0.3	0.1	0.2	0.2	0.2	0.2	0.1	0.0

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	3.3	3.5	3.1	2.1 ▼	3.8	3.1	4.5 ▲	2.5	3.7
Cigars	1.8	1.8	2.0	1.0 ▼	1.9	1.8	2.3	1.3	1.9
Pipe	0.7	1.0	0.5	0.3	0.8	0.5	0.8	0.8	0.6
Smokeless Tobacco	1.1	0.8	0.9	0.9	1.2	0.8	1.7 ▲	1.5	1.3
Alcohol	7.6	9.7 ▲	7.0	5.2 ▼	6.6	6.8	7.5	8.7	8.9 ▲
Marijuana	1.4	1.7	0.6 ▼	0.5	1.0	1.9 ▲	2.0 ▲	0.7 ▼	1.1
Cocaine	0.2	0.3	0.3	0.4	0.2	0.2	0.3	0.2	0.1
Crack	0.3	0.3	0.2	0.3	0.3	0.3	0.5	0.2	0.1
Inhalants	2.9	3.6	2.5	2.1	2.9	2.4	3.5	2.7	3.7 ▲
Amphetamines	0.2	0.2	0.1	0.3	0.3	0.2	0.5	0.2	0.2
Methamphetamines	0.2	0.3	0.2	0.2	0.1	0.2	0.3	0.1	0.3
Ritalin	0.3	0.4	0.3	0.2	0.2	0.2	0.4	0.2	0.3
Tranquilizers	1.5	2.0	1.1	1.2	1.2	1.3	1.7	1.5	2.0
Narcotics	0.3	0.3	0.2	0.3	0.4	0.3	0.6	0.3	0.3
Psychedelics [#]	0.2	0.3	0.2	0.2	0.3	0.3	0.3	0.2	0.2
LSD	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.1	0.1
Others	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
Heroin	0.2	0.3	0.2	0.2	0.4	0.2	0.2	0.1	0.1
Steroids	0.3	0.3	0.1	0.2	0.6 ▲	0.2	0.5	0.2	0.3
Injected Drugs	0.3	0.3	0.3	0.1	0.6 ▲	0.3	0.5	0.3	0.3
MDMA	0.1	0.2	0.1	0.2	0.2	0.2	0.3	0.1	0.1
Rohypnol	0.1	0.1	0.0	0.2	0.1	0.1	0.1	0.2	0.1
GHB	0.1	0.1	0.0	0.1	0.0	0.1	0.0	0.1	0.0

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
CIGARETTES									
Daily Use	1.4	1.6	1.2	0.8	1.5	1.4	1.9	1.1	1.3
1/2+ Pack per Day	0.7	0.9	0.5	0.3	0.8	1.0	0.6	0.5	0.6
CIGARS									
	0.2	0.1	0.3	0.0	0.5 ▲	0.2	0.3	0.2	0.2
SMOKELESS TOBACCO									
	0.2	0.1	0.1	0.0	0.2	0.2	0.1	0.3	0.4 ▲
ALCOHOL									
Daily Use	0.3	0.4	0.4	0.1	0.2	0.3	0.5	0.5	0.4
Binge Drinking	4.2	4.7	4.2	2.7 ▼	4.0	4.1	4.7	4.4	4.5
MARIJUANA									
	0.3	0.3	0.2	0.0	0.3	0.5 ▲	0.5	0.2	0.0

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p<.05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p<.05$).



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 7th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	19.7	21.6 ▲	21.5 ▲	16.2 ▼	18.9	12.9 ▼	30.9 ▲	19.8	25.6 ▲
Cigars	8.3	8.9	8.5	6.5 ▼	8.2	5.2 ▼	12.5 ▲	9.3	12.1 ▲
Pipe	2.9	2.8	3.2	2.3	2.9	1.8 ▼	4.6 ▲	3.0	4.5 ▲
Smokeless Tobacco	5.0	3.5 ▼	3.9 ▼	2.9 ▼	5.6	3.2 ▼	7.1 ▲	6.3 ▲	10.1 ▲
Alcohol	33.3	40.5 ▲	35.8 ▲	30.3 ▼	30.9 ▼	25.2 ▼	37.6 ▲	35.1	41.0 ▲
Marijuana	6.8	9.2 ▲	6.2	4.1 ▼	5.6 ▼	4.8 ▼	13.0 ▲	5.4	8.7 ▲
Cocaine	1.5	2.1 ▲	2.1 ▲	1.0	1.1	1.0 ▼	2.1	1.0	2.0
Crack	1.5	2.1	1.5	1.0	1.6	1.0 ▼	2.3 ▲	1.1	2.2 ▲
Inhalants	9.0	12.2 ▲	8.7	7.2	7.3 ▼	6.9 ▼	12.2 ▲	9.0	10.7 ▲
Amphetamines	1.4	1.5	1.7	0.6 ▼	1.4	1.0 ▼	2.2 ▲	1.3	1.9
Methamphetamines	1.2	1.6	1.1	0.6	1.1	0.8 ▼	1.5	1.0	1.8 ▲
Ritalin	1.8	1.9	2.1	1.5	1.6	1.2 ▼	3.1 ▲	1.4	2.8 ▲
Tranquilizers	6.0	7.5 ▲	6.5	4.3 ▼	5.7	4.3 ▼	7.8 ▲	5.9	7.9 ▲
Narcotics	2.0	2.5	1.8	1.8 ▼	1.7	1.3 ▼	3.4 ▲	1.4	3.2 ▲
Psychedelics [#]	1.5	2.1 ▲	1.4	0.6 ▼	1.3	1.0 ▼	2.4 ▲	1.2	2.3 ▲
LSD	1.1	1.8 ▲	0.9	0.1	0.9	0.9	1.7 ▲	0.8	1.6 ▲
Others	0.8	1.1	0.7	0.5	0.8	0.5 ▼	1.0	0.8	1.4 ▲
Heroin	1.1	1.4	1.2	0.5	0.8	0.8 ▼	1.4	1.3	1.7 ▲
Steroids	1.2	1.5	1.4	0.8	0.8	0.7 ▼	1.6	0.9	2.0 ▲
Injected Drugs	1.3	1.9 ▲	1.1	0.5	1.1	1.2	1.3	1.1	1.6
MDMA	1.2	2.1 ▲	1.2	0.1	0.8	0.9	1.5	0.6 ▼	2.1 ▲
Rohypnol	0.8	1.1	0.7	0.3	0.7	0.5	0.9	0.5	1.4 ▲
GHB	0.5	0.9 ▲	0.4	0.0	0.4	0.3	0.4	0.3	1.2 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 7th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	11.7	12.7	12.2	9.7	10.8	7.8	19.0	11.3	15.9
Cigars	5.8	6.4	5.7	4.8	5.7	3.6	9.7	5.9	7.9
Pipe	2.1	2.2	2.6	1.3	1.9	1.2	3.3	2.3	3.4
Smokeless Tobacco	3.9	2.8	▼	3.1	2.0	▼	4.1	2.5	▼
Alcohol	25.5	32.3	▲	26.2	23.3	24.6	18.6	29.4	▲
Marijuana	5.6	7.8	▲	5.2	3.0	▼	4.6	3.9	▼
Cocaine	1.2	1.8	▲	1.7	▲	0.7	0.8	0.7	1.7
Crack	1.2	1.5	1.3	0.9	1.1	0.7	▼	2.1	▲
Inhalants	6.3	8.7	▲	6.7	5.0	5.0	▼	4.7	▼
Amphetamines	1.1	1.4	1.3	0.7	1.0	0.8	▼	1.7	▲
Methamphetamines	0.8	1.3	▲	0.8	0.3	0.5	0.5	▼	1.3
Ritalin	1.3	1.6	1.6	1.1	1.2	0.7	▼	1.8	0.9
Tranquilizers	4.6	5.2	5.3	3.6	4.5	▼	3.2	▼	5.9
Narcotics	1.4	2.0	▲	1.4	1.4	1.2	0.8	▼	2.1
Psychedelics [#]	1.1	1.7	▲	1.0	0.5	0.8	0.7	▼	1.5
LSD	0.8	1.5	▲	0.7	0.1	0.5	0.5	▼	1.0
Others	0.6	0.8	0.7	0.4	0.5	0.3	▼	0.7	0.5
Heroin	0.8	1.2	0.9	0.3	0.7	0.5	▼	1.1	0.8
Steroids	0.9	1.2	1.3	0.7	0.7	0.4	1.2	0.6	1.6
Injected Drugs	1.0	1.7	▲	1.0	0.4	0.8	0.7	▼	0.9
MDMA	0.9	1.7	▲	0.8	0.1	0.5	0.6	1.1	0.5
Rohypnol	0.6	0.9	▲	0.6	0.2	0.5	0.3	▼	0.7
GHB	0.4	0.8	▲	0.4	0.0	0.3	0.2	0.2	0.4

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 7th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	6.7	7.2	6.9	5.1	6.0	4.3 ▼	12.0 ▲	7.2	9.0 ▲
Cigars	3.2	2.7	3.5	2.6	3.3	1.7 ▼	5.7 ▲	3.7	5.2 ▲
Pipe	1.2	1.2	1.5	0.6	1.2	0.6 ▼	2.0 ▲	1.2	2.4 ▲
Smokeless Tobacco	2.3	1.4 ▼	1.9	0.9 ▼	2.5	1.2 ▼	3.6 ▲	2.9	5.3 ▲
Alcohol	12.9	17.2 ▲	12.7	10.1 ▼	11.6	8.6 ▼	16.6 ▲	14.4 ▲	16.8 ▲
Marijuana	3.6	5.2 ▲	3.1	1.4 ▼	2.9	2.5 ▼	7.4 ▲	2.9	4.6 ▲
Cocaine	0.8	1.1	1.1	0.6	0.5	0.5 ▼	1.0	0.4	1.2 ▲
Crack	0.8	1.3 ▲	0.9	0.6	0.8	0.3 ▼	1.0	0.5	1.3 ▲
Inhalants	3.6	4.7 ▲	4.4	3.0	2.7 ▼	2.3 ▼	5.3 ▲	3.3	5.1 ▲
Amphetamines	0.8	1.0	0.8	0.7	0.8	0.6 ▼	1.0	0.7	1.3 ▲
Methamphetamines	0.6	0.9	0.5	0.2	0.5	0.3 ▼	0.8	0.4	1.1 ▲
Ritalin	0.8	0.9	0.8	0.7	1.0	0.5 ▼	1.0	0.8	1.3 ▲
Tranquilizers	2.8	3.1	3.2	2.5	2.9	1.8 ▼	3.8 ▲	2.6	4.1 ▲
Narcotics	0.9	1.3 ▲	0.8	0.9	0.8	0.5 ▼	1.0	0.7	1.3 ▲
Psychedelics [#]	0.7	1.2	0.7	0.4	0.8	0.4 ▼	0.8	0.6	1.0
LSD	0.5	1.0 ▲	0.6	0.1	0.4	0.3 ▼	0.6	0.5	0.8
Others	0.4	0.6	0.3	0.3	0.5	0.3	0.3	0.4	0.6
Heroin	0.5	0.8	0.7	0.1	0.3	0.3	0.7	0.5	0.7
Steroids	0.6	0.8	0.9	0.6	0.5	0.3 ▼	0.8	0.2	0.9
Injected Drugs	0.6	1.0 ▲	0.8	0.3	0.5	0.4 ▼	0.7	0.5	0.9
MDMA	0.6	1.1 ▲	0.5	0.1	0.3	0.5	0.8	0.3	1.0 ▲
Rohypnol	0.3	0.5	0.4	0.1	0.3	0.2	0.4	0.2	0.6
GHB	0.3	0.6 ▲	0.3	0.0	0.2	0.2	0.3	0.2	0.6 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 7th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
CIGARETTES									
Daily Use	3.0	2.9	3.3	2.4	2.5	2.2 ▼	5.7 ▲	2.7	4.4 ▲
1/2+ Pack per Day	1.4	1.4	1.3	0.8	1.0	0.9 ▼	3.0 ▲	0.9	2.3 ▲
CIGARS	0.6	0.4	0.5	0.4	0.6	0.4	1.5 ▲	0.8	0.7
SMOKELESS TOBACCO									
	0.4	0.1	0.4	0.3	0.4	0.2 ▼	0.6	0.4	0.9 ▲
ALCOHOL									
Daily Use	0.8	1.0	0.6	0.6	0.7	0.6 ▼	1.0	1.0	1.6 ▲
Binge Drinking	7.0	9.9 ▲	6.9	4.2 ▼	5.4 ▼	4.5 ▼	10.1 ▲	6.9	10.3 ▲
MARIJUANA	0.7	1.2 ▲	0.5	0.3	0.5	0.5	1.3 ▲	0.5	1.0

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p<.05$).
 ▼ indicates that local prevalence rates are significantly lower than state rates ($p<.05$).



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 8th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	30.2	31.6	30.8	27.1 ▼	30.5	26.6 ▼	36.5 ▲	21.8 ▼	35.1 ▲
Cigars	14.6	14.5	14.9	12.2 ▼	14.3	12.7 ▼	17.5 ▲	10.4 ▼	18.5 ▲
Pipe	4.9	5.1	5.4	2.9 ▼	5.4	4.3	5.1	3.4 ▼	6.0 ▲
Smokeless Tobacco	8.3	4.6 ▼	8.0	6.7	11.6 ▲	5.8 ▼	7.8	6.9 ▼	13.5 ▲
Alcohol	46.5	51.4 ▲	46.3	40.1 ▼	45.0 ▼	41.5 ▼	48.7 ▲	46.9	52.9 ▲
Marijuana	15.6	19.3 ▲	14.5	12.3 ▼	14.1 ▼	16.2	20.0 ▲	7.8 ▼	15.6
Cocaine	2.5	3.3 ▲	2.8	1.9	2.5	2.3	3.2 ▲	1.1 ▼	2.9
Crack	2.2	2.6	2.6	1.8	2.8	2.0	2.6	1.2 ▼	2.2
Inhalants	10.4	11.9 ▲	11.1	8.1 ▼	9.8	8.4 ▼	12.9 ▲	9.2	12.3 ▲
Amphetamines	3.2	3.7	3.7	2.1	3.8	2.7	3.4	1.7 ▼	3.8
Methamphetamines	2.0	2.2	2.4	1.3	2.2	1.6	1.9	1.1 ▼	2.5 ▲
Ritalin	3.1	3.6	3.9 ▲	2.7	3.0	2.8	3.6	1.6 ▼	3.6
Tranquilizers	8.9	10.6 ▲	8.8	6.8 ▼	8.4	8.3	10.0	6.4 ▼	10.3 ▲
Narcotics	4.1	5.0 ▲	4.0	2.8 ▼	4.0	3.5 ▼	5.1 ▲	2.4 ▼	5.3 ▲
Psychedelics [#]	2.8	2.9	2.8	2.0	2.8	2.9	2.9	1.0 ▼	3.5 ▲
LSD	2.3	2.4	1.9	1.8	2.4	2.3	2.5	0.4 ▼	3.0 ▲
Others	1.3	1.4	1.7	0.7	1.1	1.4	1.2	0.9	1.7
Heroin	1.5	2.0 ▲	1.3	1.0	1.6	1.4	1.7	0.8 ▼	1.5
Steroids	1.5	1.4	2.2 ▲	1.7	1.6	1.5	1.5	0.6 ▼	1.7
Injected Drugs	1.7	1.8	2.2	1.3	2.1	1.5	1.5	1.0 ▼	1.8
MDMA	2.3	2.8	1.9	1.2 ▼	2.6	2.4	2.3	0.8 ▼	2.9 ▲
Rohypnol	1.2	1.6	1.1	0.7	1.4	1.0	1.4	0.6 ▼	1.5
GHB	0.8	0.9	0.7	0.6	0.8	0.7	0.7	0.5	0.9

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 8th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	19.1	20.0	20.1	17.6	18.8	16.9 ▼	23.8 ▲	12.5 ▼	22.4 ▲
Cigars	10.3	10.8	9.5	8.0 ▼	10.0	9.1 ▼	12.6 ▲	6.9 ▼	13.2 ▲
Pipe	3.5	3.4	4.0	2.0 ▼	3.8	3.2	3.6	2.2 ▼	4.3 ▲
Smokeless Tobacco	6.4	3.9 ▼	6.0	5.2	9.0 ▲	4.4 ▼	5.8	4.8 ▼	11.1 ▲
Alcohol	38.3	43.4 ▲	37.8	32.6 ▼	36.0 ▼	33.8 ▼	40.2	39.6	43.7 ▲
Marijuana	12.7	15.9 ▲	11.1 ▼	9.8 ▼	10.9 ▼	13.3	16.6 ▲	6.2 ▼	13.3
Cocaine	1.9	2.4	2.2	1.1	2.1	1.6	2.1	0.8 ▼	2.2
Crack	1.7	2.1	1.9	1.2	2.1	1.5	1.8	0.7 ▼	1.8
Inhalants	7.2	8.9 ▲	7.3	6.1	7.0	5.7 ▼	8.8 ▲	5.8 ▼	8.7 ▲
Amphetamines	2.5	2.7	2.8	1.5 ▼	3.4 ▲	2.0 ▼	2.7	1.5 ▼	3.0
Methamphetamines	1.4	1.7	1.8	1.0	1.9	1.2	1.1	0.6 ▼	1.9 ▲
Ritalin	2.3	2.7	2.7	2.0	2.0	2.1	2.6	1.4 ▼	2.8
Tranquilizers	6.8	8.3 ▲	6.7	5.1 ▼	6.9	6.2	7.4	5.1 ▼	8.0 ▲
Narcotics	3.0	4.0 ▲	3.0	1.7	3.0	2.5 ▼	3.4	1.7 ▼	3.8 ▲
Psychedelics [#]	2.1	2.5	2.1	1.4	2.1	2.0	2.2	0.8 ▼	2.6 ▲
LSD	1.7	2.1	1.5	1.1	1.8	1.6	1.7	0.5 ▼	2.2 ▲
Others	1.1	1.3	1.2	0.7	0.9	1.0	1.0	0.6	1.3
Heroin	1.0	1.5	0.9	0.5	1.3	1.0	1.1	0.5 ▼	1.1
Steroids	1.1	1.2	1.4	1.3	1.2	1.0	1.0	0.6 ▼	1.1
Injected Drugs	1.2	1.5	1.5	1.1	1.5	1.0	1.0	0.9	1.3
MDMA	1.7	2.2	1.5	1.0	1.8	1.7	1.6	0.6 ▼	2.2 ▲
Rohypnol	0.8	1.3 ▲	0.7	0.4	1.2	0.7	0.8	0.4 ▼	1.0
GHB	0.5	0.7	0.5	0.3	0.7	0.4	0.5	0.5	0.6

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 8th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	11.6	12.3	11.9	11.1	11.5	10.5 ▼	15.3 ▲	6.4 ▼	13.0
Cigars	5.9	5.5	5.9	4.3 ▼	5.7	5.5	7.4 ▲	3.7 ▼	7.6 ▲
Pipe	2.0	2.0	2.4	1.5	2.3	2.0	1.9	1.0 ▼	2.4
Smokeless Tobacco	3.8	2.2 ▼	3.6	2.3 ▼	5.2 ▲	2.8 ▼	3.4	2.2 ▼	6.9 ▲
Alcohol	20.9	24.3 ▲	19.6	16.8 ▼	19.9	18.0 ▼	21.5	21.1	25.6 ▲
Marijuana	8.2	10.5 ▲	6.5 ▼	7.0	7.0 ▼	8.6	10.3 ▲	4.1 ▼	8.6
Cocaine	1.1	1.9 ▲	1.5	0.5	1.2	0.9	1.1	0.5 ▼	1.3
Crack	1.0	1.4	1.2	0.6	1.2	0.8	0.9	0.6	1.1
Inhalants	4.1	5.2 ▲	4.5	3.3	3.7	3.2 ▼	4.6	3.3	5.0 ▲
Amphetamines	1.4	1.6	1.9	0.9	1.8	1.2	1.4	0.9	1.7
Methamphetamines	0.9	1.1	1.3	0.4	1.3 ▲	0.7	0.8	0.5	1.0
Ritalin	1.3	1.5	1.7	1.4	1.2	1.2	1.1	0.9	1.6
Tranquilizers	4.1	4.9	4.3	3.0	4.3	3.8	4.0	3.2 ▼	4.9 ▲
Narcotics	1.8	2.4 ▲	1.9	1.3	1.9	1.5	2.1	1.1 ▼	2.1
Psychedelics [#]	1.2	1.6	1.3	0.8	1.1 ▼	1.1	1.3	0.5 ▼	1.5
LSD	0.9	1.3	0.8	0.7	1.0	0.9	0.9	0.4 ▼	1.3
Others	0.6	0.8	1.0	0.4	0.5	0.6	0.7	0.3	0.6
Heroin	0.7	1.1 ▲	0.7	0.4	0.9	0.6	0.8	0.3	0.7
Steroids	0.7	0.8	0.9	1.0	0.7	0.7	0.6	0.4	0.6
Injected Drugs	0.8	1.0	1.2	0.3	1.2	0.7	0.8	0.4	0.9
MDMA	1.0	1.3	1.0	0.5	1.0	0.9	1.0	0.3	1.3
Rohypnol	0.5	0.8	0.6	0.2	0.7	0.5	0.6	0.2	0.5
GHB	0.4	0.5	0.4	0.1	0.3	0.3	0.3	0.3	0.4

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

- ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
- ▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 8th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
CIGARETTES									
Daily Use	5.8	5.3	6.4	4.7	5.7	5.3	8.2 ▲	3.2 ▼	6.5
1/2+ Pack per Day	2.9	2.4	3.2	2.0	2.7	2.7	4.0 ▲	1.8 ▼	3.4
CIGARS									
	1.3	1.0	1.5	0.9	1.4	1.1	1.6	0.5 ▼	1.8 ▲
SMOKELESS TOBACCO									
	1.0	0.2	0.6	0.2	1.3	0.8	0.8	0.5 ▼	2.2 ▲
ALCOHOL									
Daily Use	1.6	1.9	1.6	0.9	1.5	1.4	1.8	0.9 ▼	2.3 ▲
Binge Drinking	11.0	13.0 ▲	11.0	8.3 ▼	11.6	9.6 ▼	10.9	10.1	13.0 ▲
MARIJUANA									
	2.0	2.3	1.5	0.7 ▼	1.7	2.2	3.1 ▲	0.8 ▼	1.9

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p<.05$).
 ▼ indicates that local prevalence rates are significantly lower than state rates ($p<.05$).



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 9th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	36.3	37.6	38.4	31.8 ▼	36.3	30.1 ▼	42.0 ▲	37.7	43.2 ▲
Cigars	20.4	20.8	21.6	15.1 ▼	21.1	18.1 ▼	23.2 ▲	21.5	23.1 ▲
Pipe	6.9	6.3	6.8	5.6	7.0	6.5	8.3 ▲	5.8	8.0
Smokeless Tobacco	11.6	7.7 ▼	11.0	10.1	15.1 ▲	9.0 ▼	10.2	17.5 ▲	19.3 ▲
Alcohol	55.5	59.1 ▲	53.5	50.4 ▼	55.7	51.1 ▼	58.8 ▲	59.9 ▲	60.0 ▲
Marijuana	22.5	25.6 ▲	20.5	18.9 ▼	22.1	20.5 ▼	28.6 ▲	16.3 ▼	24.3
Cocaine	4.4	4.6	3.9	4.6	4.5	4.6	4.9	2.4 ▼	4.6
Crack	3.0	2.8	2.3	2.8	2.9	3.2	3.2	2.2	3.7
Inhalants	11.4	11.8	10.3	8.3 ▼	11.6	10.5	14.0 ▲	12.1	12.0
Amphetamines	6.8	6.5	6.5	4.4 ▼	7.6	6.6	8.4 ▲	5.5	7.1
Methamphetamines	3.0	2.6	2.6	3.0	3.5	2.8	2.7	2.9	3.8
Ritalin	5.4	5.3	5.6	4.5	6.0	5.6	5.4	3.9	5.3
Tranquilizers	12.4	13.4	10.6 ▼	8.6 ▼	13.4	11.6	14.4 ▲	11.5	14.0
Narcotics	7.4	6.7	6.8	6.1	7.5	7.5	8.8 ▲	5.7	8.3
Psychedelics [#]	4.2	4.1	3.2 ▼	3.5	4.2	5.0 ▲	4.1	3.7	4.1
LSD	3.4	3.0	2.7	3.3	3.4	4.0	3.4	3.2	3.4
Others	2.2	2.3	1.3	1.8	2.4	2.8 ▲	1.8	1.7	1.9
Heroin	1.9	1.8	1.4	2.0	1.9	2.3	1.7 ▼	2.0	1.8
Steroids	1.8	1.7	1.3	2.2	1.5	2.2	1.6 ▼	1.8	2.3
Injected Drugs	1.8	1.7	1.4	2.0	1.7	2.2	1.3	1.1	1.9
MDMA	3.7	4.3	3.0	2.4 ▼	3.7	4.4 ▲	3.4 ▼	2.1 ▼	3.7
Rohypnol	1.8	1.6	1.3	1.9	1.7	2.2	1.7	1.1	2.1
GHB	1.0	1.0	0.5	1.1	1.0	1.2	1.0	0.3	0.9

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

- ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
- ▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 9th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	25.0	25.0	26.4	20.1 ▼	24.8	21.3 ▼	27.8 ▲	29.4 ▲	30.5 ▲
Cigars	15.3	15.5	16.3	10.6 ▼	15.3	13.9 ▼	17.7 ▲	16.2 ▲	17.5 ▲
Pipe	5.1	5.1	5.1	3.6 ▼	5.2	5.1	5.8	3.3	6.1
Smokeless Tobacco	9.6	5.9 ▼	8.7	8.3	12.3 ▲	7.7 ▼	8.5	14.1 ▲	15.9 ▲
Alcohol	47.3	50.6 ▲	46.5	42.0 ▼	47.2	43.5 ▼	49.6	50.7	52.5 ▲
Marijuana	18.4	21.9 ▲	16.6	13.8	18.0	17.4	23.3 ▲	12.2 ▼	19.3
Cocaine	3.3	3.8	2.9	3.2	2.8	3.8	3.6	1.4 ▼	3.1
Crack	2.2	2.0	1.5	1.7	2.1	2.5	2.4	1.5	2.5
Inhalants	7.5	7.8	6.9	5.1 ▼	7.3	7.2	9.2 ▲	8.5	7.9
Amphetamines	5.1	5.2	5.2	3.2 ▼	5.2	5.1	6.3 ▲	3.9	5.2
Methamphetamines	2.2	2.0	1.9	2.0	2.3	2.2	1.9	1.8	2.8
Ritalin	3.8	3.8	3.8	3.6	4.1	3.9	3.7	3.1	4.0
Tranquilizers	9.6	10.5	7.8 ▼	6.3 ▼	10.0	8.9	11.8 ▲	8.9	10.9
Narcotics	5.3	5.2	5.0	3.8 ▼	5.3	5.7	5.9	3.9	5.9
Psychedelics [#]	3.1	3.1	2.0 ▼	3.0	2.9	4.0 ▲	3.0	2.2	2.8
LSD	2.4	2.3	1.6 ▼	2.6	2.3	3.0 ▲	2.5	2.0	2.4
Others	1.6	1.6	0.9 ▼	1.4	1.6	2.3 ▲	1.5	1.1	1.2
Heroin	1.4	1.3	1.1	1.5	1.3	2.0 ▲	1.3	1.1	1.2
Steroids	1.4	1.2	0.9	1.5	1.1	1.8 ▲	1.3	1.1	1.8
Injected Drugs	1.3	1.3	1.0	1.5	1.2	1.6	1.0	0.9	1.4
MDMA	2.7	3.1	2.0	1.8	2.3	3.6 ▲	2.4	1.3 ▼	2.4
Rohypnol	1.3	1.2	1.0	1.4 ▲	1.1	1.7 ▲	1.1	0.9	1.4
GHB	0.8	0.7	0.5	0.7	0.7	1.0	0.8	0.3	0.6

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 9th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	16.5	15.5	16.8	13.6 ▼	15.7	14.4 ▼	19.2 ▲	19.6 ▲	21.5 ▲
Cigars	8.8	9.0	8.5	6.0 ▼	9.1	7.7 ▼	10.9 ▲	9.5 ▲	10.5 ▲
Pipe	3.0	3.0	2.9	1.7 ▼	3.3	2.8	3.3	1.7	3.9 ▲
Smokeless Tobacco	5.7	3.2 ▼	5.5	5.8	7.4 ▲	4.3 ▼	4.8	9.0 ▲	9.9 ▲
Alcohol	28.5	31.0 ▲	26.3	25.9	28.9	26.0 ▼	29.4	32.0 ▲	31.4 ▲
Marijuana	11.4	14.0 ▲	9.5 ▼	8.6 ▼	11.1	10.7	15.0 ▲	6.9 ▼	12.0
Cocaine	1.7	2.5 ▲	1.6	1.3	1.4	1.8	1.5	0.7 ▼	1.8
Crack	1.1	1.3	0.8	1.2	1.2	1.2	0.7	0.7	1.4
Inhalants	3.6	3.8	3.0	2.6	3.8	3.3	4.3	4.0	4.0
Amphetamines	3.0	3.3	3.1	2.2	3.3	3.0	3.5	2.7	2.3
Methamphetamines	1.2	1.4	1.3	1.0	1.5	1.2	0.9	1.1	1.6
Ritalin	2.0	2.2	1.8	1.8	2.3	2.3	1.9	1.8	1.8
Tranquilizers	5.7	6.5	4.2 ▼	3.8 ▼	5.8	5.1	7.3 ▲	5.6	6.8
Narcotics	2.9	2.8	2.9	2.6	2.7	3.2	3.2	2.4	2.8
Psychedelics [#]	1.6	2.1	0.9 ▼	1.7	1.4	1.8	1.6	1.5	1.4
LSD	1.3	1.5	0.8	1.5	1.1	1.4	1.4	1.4	0.9
Others	0.8	1.2	0.4	0.7	0.9	1.0	0.6	0.7	0.7
Heroin	0.8	1.1	0.5	1.0	0.9	1.1	0.4	0.9	0.6
Steroids	0.9	0.9	0.4 ▼	1.0	0.8	1.1	0.8	1.0	0.9
Injected Drugs	0.8	1.0	0.5	0.9	0.7	1.0	0.4	0.5	0.9
MDMA	1.5	1.9	0.9	1.0	1.2	1.8	1.3	0.6 ▼	1.5
Rohypnol	0.8	1.0	0.5	0.7	0.7	0.9	0.8	0.4	0.7
GHB	0.5	0.5	0.2	0.6	0.4	0.6	0.5	0.3	0.3

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 9th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
CIGARETTES									
Daily Use	9.6	9.5	9.9	7.2 ▼	9.0	7.7 ▼	12.9 ▲	9.8	12.4 ▲
1/2+ Pack per Day	5.4	5.2	5.0	4.5	5.6	4.1 ▼	7.6 ▲	5.8	7.3 ▲
CIGARS									
	1.9	1.6	1.7	1.0 ▼	1.8	1.4 ▼	3.3 ▲	2.6	3.0 ▲
SMOKELESS TOBACCO									
	1.8	1.0 ▼	1.5	1.5	2.2	1.3 ▼	1.7	3.7 ▲	3.7 ▲
ALCOHOL									
Daily Use	2.7	3.7 ▲	2.2	1.9	2.4	2.0 ▼	2.6	4.4 ▲	3.2
Binge Drinking	15.6	16.9	15.4	14.3	16.4	13.8 ▼	15.8	16.9	17.7 ▲
MARIJUANA									
	3.2	4.0 ▲	2.9	2.6	3.4	2.4 ▼	4.5 ▲	2.3	3.2

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p<.05$).
 ▼ indicates that local prevalence rates are significantly lower than state rates ($p<.05$).



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 10th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	43.0	43.6	41.4	39.4 ▼	44.2	37.8 ▼	48.8 ▲	39.7 ▼	49.4 ▲
Cigars	26.8	25.7	24.9	24.0	28.1	23.7 ▼	28.1	25.9	32.3 ▲
Pipe	8.6	7.9	9.1	7.1	9.9 ▲	7.0 ▼	8.8	8.2	10.5 ▲
Smokeless Tobacco	15.4	9.6 ▼	11.6 ▼	13.9	20.2 ▲	11.3 ▼	16.1	19.2 ▲	21.4 ▲
Alcohol	62.3	66.7 ▲	57.3 ▼	59.7	64.2 ▲	55.8 ▼	62.2	63.8	69.9 ▲
Marijuana	30.1	34.1 ▲	26.1 ▼	24.9 ▼	30.6	29.7	34.6 ▲	20.4 ▼	34.5 ▲
Cocaine	5.9	6.7	6.1	4.6	5.7	5.4	7.1 ▲	2.6 ▼	7.5 ▲
Crack	3.4	2.8	3.5	2.3	3.8	3.5	3.9	1.8 ▼	4.0
Inhalants	11.1	12.0	11.6	7.9 ▼	11.9	8.5 ▼	12.3	9.3 ▼	14.1 ▲
Amphetamines	9.0	9.9	9.6	7.2 ▼	9.5	8.0 ▼	10.6 ▲	5.1 ▼	10.9 ▲
Methamphetamine	3.5	2.3 ▼	3.4	3.1	4.2	3.1	3.4	2.6 ▼	5.4 ▲
Ritalin	6.5	6.5	7.1	5.6	7.2	5.7	7.0	4.4 ▼	7.6 ▲
Tranquilizers	14.3	14.9	13.7	10.4 ▼	14.9	12.4 ▼	16.9 ▲	10.2 ▼	18.3 ▲
Narcotics	9.8	10.8	8.8	7.2 ▼	9.7	9.1	12.3 ▲	6.4 ▼	12.3 ▲
Psychedelics [#]	5.3	5.7	4.5	4.4	5.3	5.6	5.5	2.6 ▼	7.1 ▲
LSD	4.3	3.9	3.7	3.6	3.7	4.6	4.8	2.3 ▼	6.2 ▲
Others	2.7	3.1	2.4	2.0	3.1	2.8	2.3	1.2 ▼	3.5 ▲
Heroin	2.0	1.9	2.3	1.4	2.2	1.9	2.1	1.3 ▼	2.5
Steroids	2.2	2.2	2.3	1.7	2.5	1.9	2.0	1.6	2.8 ▲
Injected Drugs	2.0	2.0	2.1	1.3	2.1	1.9	1.8	1.4	2.6 ▲
MDMA	4.4	5.7 ▲	4.4	3.0 ▼	3.5 ▼	4.5	4.3	2.4 ▼	5.9 ▲
Rohypnol	1.9	1.6	2.1	1.1	2.0	1.6	2.2	1.0 ▼	2.8 ▲
GHB	1.1	0.9	1.3	0.8	1.1	1.1	1.2	0.6 ▼	1.7 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 10th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	30.0	29.6	29.0	27.0 ▼	32.4 ▲	25.6 ▼	32.9 ▲	30.1	34.3 ▲
Cigars	20.2	18.8	18.1 ▼	18.0	21.2	17.6 ▼	21.0	20.9	24.9 ▲
Pipe	6.3	5.9	5.9	6.3	7.0	5.2 ▼	6.3	6.1	7.7 ▲
Smokeless Tobacco	12.2	7.1 ▼	8.4 ▼	11.6	16.0 ▲	9.2 ▼	11.7	15.6 ▲	17.5 ▲
Alcohol	53.7	58.2 ▲	48.1 ▼	50.2 ▼	55.3	47.9 ▼	52.0	56.5	61.4 ▲
Marijuana	23.8	27.0 ▲	19.7 ▼	21.1 ▼	24.1	23.9	26.9 ▲	16.1 ▼	27.4 ▲
Cocaine	4.3	5.2	4.6	3.8	4.3	3.9	4.6	1.8 ▼	5.7 ▲
Crack	2.3	1.8	2.3	1.6	2.6	2.3	2.7	1.1 ▼	3.0 ▲
Inhalants	6.5	6.5	7.3	5.2	7.1	4.6 ▼	7.2	5.5	8.9 ▲
Amphetamines	6.8	7.2	6.9	5.9	6.9	6.0	8.0 ▲	3.7 ▼	8.8 ▲
Methamphetamines	2.4	1.6 ▼	2.5	2.1	2.8	2.0	2.2	1.5 ▼	3.7 ▲
Ritalin	4.5	4.5	5.0	4.3	4.9	4.1	4.7	2.7 ▼	5.1
Tranquilizers	10.7	11.8	10.0	7.9 ▼	11.3	9.2 ▼	12.5 ▲	7.7 ▼	13.6 ▲
Narcotics	7.0	7.2	6.3	5.1 ▼	6.2	6.5	9.4 ▲	5.0 ▼	9.1 ▲
Psychedelics [#]	3.7	4.4	3.2	3.4	3.7	3.8	3.3	1.8 ▼	5.0 ▲
LSD	2.9	2.9	2.5	2.7	2.4	3.0	2.8	1.5 ▼	4.2 ▲
Others	2.0	2.5	2.0	1.5	2.1	2.0	1.4	0.9 ▼	2.4
Heroin	1.3	1.3	1.4	0.9	1.6	1.2	1.4	0.8	1.6
Steroids	1.6	1.4	1.7	1.4	1.9	1.3	1.6	1.0	2.1 ▲
Injected Drugs	1.4	1.4	1.7	1.0	1.4	1.2	1.1	0.9	1.9 ▲
MDMA	3.0	4.1 ▲	3.4	2.3	2.4 ▼	2.7	2.8	1.8	4.2 ▲
Rohypnol	1.4	1.1	1.6	0.6	1.4	1.3	1.5	0.9	2.0 ▲
GHB	0.9	0.8	1.1	0.5	0.8	0.9	1.0	0.5	1.2

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 10th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	20.6	19.7	19.6	18.2	22.4 ▲	17.1 ▼	23.2 ▲	20.6	24.0 ▲
Cigars	11.6	10.1 ▼	11.1	11.4	11.9	9.5 ▼	12.7	12.5	14.3 ▲
Pipe	3.4	2.9	3.7	4.5	3.2	3.2	3.3	3.6	3.9
Smokeless Tobacco	7.8	4.0 ▼	5.2 ▼	7.4	10.6 ▲	5.9 ▼	7.7	10.2 ▲	11.5 ▲
Alcohol	33.0	34.8	29.0	31.1	34.1	28.8 ▼	31.2	38.6 ▲	38.0 ▲
Marijuana	14.6	16.8 ▲	11.9	13.0	14.2	14.2	18.3 ▲	9.0 ▼	17.2 ▲
Cocaine	2.1	2.4	2.5 ▲	1.8	1.9	1.8	2.6	1.1 ▼	2.8 ▲
Crack	1.2	0.9	1.4	1.0	1.2	1.3	1.7 ▲	0.4 ▼	1.3
Inhalants	3.0	2.7	4.4 ▲	2.3	3.2	1.9 ▼	3.3	2.3	3.9 ▲
Amphetamines	3.7	3.1	4.3	3.2	3.9	3.2	4.7 ▲	1.9 ▼	4.9 ▲
Methamphetamines	1.2	0.8	1.2	1.1	1.4	0.9	1.3	0.9	1.7 ▲
Ritalin	2.2	1.9	2.7	1.8	2.8 ▲	1.9	2.4	1.1 ▼	2.4
Tranquilizers	5.9	6.4	5.9	4.3 ▼	6.3	5.0 ▼	7.2 ▲	3.6 ▼	7.7 ▲
Narcotics	3.8	3.5	3.9	2.4 ▼	3.5	3.4	5.4 ▲	2.4 ▼	4.9 ▲
Psychedelics [#]	1.9	2.2	1.6	1.6	1.9	1.9	2.1	1.0 ▼	2.2
LSD	1.5	1.6	1.4	1.4	1.4	1.5	1.8	1.0	1.9
Others	1.0	1.2	0.9	0.8	1.2	1.0	1.1	0.5 ▼	1.2
Heroin	0.8	0.6	1.0	0.6	1.1	0.9	0.9	0.5	0.9
Steroids	1.1	0.8	1.3	1.0	1.4	0.9	1.2	0.6	1.3
Injected Drugs	0.9	0.9	1.2	0.8	1.0	0.9	0.8	0.5	0.9
MDMA	1.5	2.2 ▲	1.5	1.0	1.1	1.3	1.4	0.9 ▼	2.0 ▲
Rohypnol	0.8	0.7	1.1	0.3	0.8	0.8	0.9	0.6	0.8
GHB	0.5	0.4	0.8	0.3	0.5	0.6	0.6	0.3	0.5

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 10th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
CIGARETTES									
Daily Use	12.2	11.5	12.1	10.0 ▼	13.1	10.2 ▼	15.0 ▲	10.2 ▼	14.7 ▲
1/2+ Pack per Day	7.1	7.1	6.8	4.9 ▼	7.3	5.5 ▼	9.4 ▲	5.4 ▼	9.2 ▲
CIGARS									
	2.5	1.6 ▼	2.5	2.5	2.6	2.1	3.2 ▲	2.0	3.2 ▲
SMOKELESS TOBACCO									
	3.2	0.9 ▼	1.6 ▼	2.5 ▼	4.9 ▲	2.4 ▼	2.7	4.0	5.4 ▲
ALCOHOL									
Daily Use	3.6	3.3	3.2	2.7	4.0	2.9 ▼	3.7	4.3	4.2 ▲
Binge Drinking	19.9	21.1	17.4 ▼	19.7	19.7	17.3 ▼	19.3	23.9 ▲	22.7
MARIJUANA									
	4.5	5.2	3.7	5.3	3.8	4.4	6.8 ▲	2.2 ▼	4.9

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p<.05$).
 ▼ indicates that local prevalence rates are significantly lower than state rates ($p<.05$).



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 11th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	46.8	48.5	47.5	41.8 ▼	48.7	42.5 ▼	46.8	50.2	53.3 ▲
Cigars	30.7	30.7	31.6	28.0	31.3	30.2	29.2	31.0	34.7 ▲
Pipe	9.2	8.9	9.5	6.6 ▼	9.5	9.5	8.9	8.3	11.0 ▲
Smokeless Tobacco	17.2	13.6 ▼	16.2	15.2	19.0	14.1 ▼	16.4	23.7 ▲	27.7 ▲
Alcohol	66.7	70.4 ▲	64.3	62.5 ▼	69.8 ▲	62.1 ▼	65.6	71.4 ▲	71.3 ▲
Marijuana	34.4	40.2 ▲	30.7 ▼	28.9 ▼	34.4	33.9	36.8 ▲	26.5 ▼	36.8
Cocaine	7.8	9.6 ▲	6.7	6.0	8.3	7.6	8.4	4.3 ▼	8.1
Crack	3.8	3.8	3.0	3.1	4.1	3.4 ▼	4.4	3.1	4.9
Inhalants	10.1	10.9	11.0	7.8 ▼	10.0	8.7	11.6	8.8	12.3 ▲
Amphetamines	10.8	11.0	10.9	7.4 ▼	11.9	10.3	12.6 ▲	8.0 ▼	11.8
Methamphetamines	4.2	4.0	4.1	2.9	4.4	3.9	4.5	4.9	5.3
Ritalin	6.7	7.8	7.1	5.4	6.9	6.6	7.5	3.5 ▼	6.0
Tranquilizers	14.5	16.0	13.4	10.1 ▼	14.7	14.0	16.5 ▲	11.5 ▼	16.8 ▲
Narcotics	11.3	12.2	10.4	7.2 ▼	11.6	12.0	12.8	7.3 ▼	11.9
Psychedelics [#]	6.3	7.1 ▲	4.4 ▼	5.3	7.6 ▲	6.6	6.6	3.5 ▼	6.6
LSD	4.9	5.3	3.0 ▼	4.1	5.0	5.4	5.4	3.1 ▼	5.5
Others	3.8	4.7	2.5 ▼	3.7	4.6	3.7	3.9	1.7 ▼	3.5
Heroin	2.3	2.9	2.0	2.1	2.6	2.3	2.2	1.6	2.2
Steroids	1.9	2.3	1.7	2.2	1.9	1.7	2.2	1.6	1.9
Injected Drugs	2.1	2.0	1.6	2.2	2.0	2.1	2.3	2.2	2.6
MDMA	5.4	8.3 ▲	4.2	3.3 ▼	4.7	6.0	5.0	3.3 ▼	5.2
Rohypnol	2.0	2.3	1.9	1.5	2.2	1.6	2.2	1.3	2.5
GHB	1.2	1.7	1.5	0.8	1.1	1.2	1.4	0.6	1.1

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

- ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
- ▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 11th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	33.7	34.5	34.1	30.3 ▼	34.4	31.6 ▼	31.7	37.8 ▲	39.3 ▲
Cigars	23.1	22.2	23.5	22.6	22.5	23.4	21.5	22.5	27.7 ▲
Pipe	6.6	6.2	7.6	5.3	7.0	6.8	6.0	4.5 ▼	8.6 ▲
Smokeless Tobacco	13.1	9.9 ▼	12.5	12.7	15.0 ▲	10.9 ▼	11.5	18.2 ▲	21.4 ▲
Alcohol	57.0	61.9 ▲	52.9 ▼	54.3	58.0	53.6 ▼	54.2 ▼	61.8 ▲	63.2 ▲
Marijuana	26.5	31.1 ▲	23.7 ▼	21.7 ▼	26.1	26.5	29.3 ▲	19.0 ▼	27.9
Cocaine	5.3	6.6 ▲	4.6	4.0	5.4	5.6	5.5	2.6 ▼	6.2
Crack	2.4	2.4	2.0	1.9	2.2	2.5	2.8	1.6	3.7 ▲
Inhalants	5.2	5.7	5.8	3.7 ▼	4.8	4.4	6.2	4.1	6.9 ▲
Amphetamines	7.6	7.8	7.5	4.9 ▼	8.0	7.6	8.6	5.0 ▼	9.6 ▲
Methamphetamines	2.7	2.4	2.7	2.1	2.9	2.5	2.3	3.0	3.8 ▲
Ritalin	4.2	4.8	4.7	3.3	4.1	4.1	4.6	2.1 ▼	4.3
Tranquilizers	10.0	11.5 ▲	9.5	7.2 ▼	10.1	9.6	11.3	7.1 ▼	11.8
Narcotics	7.6	8.1	7.0	5.2 ▼	7.8	8.5	8.0	4.5 ▼	8.7
Psychedelics [#]	4.2	4.3	2.8 ▼	3.4	5.1 ▲	4.8	4.0	1.9 ▼	4.6
LSD	3.1	3.2	1.7 ▼	2.1	2.9	3.7 ▲	3.3	1.6 ▼	3.9
Others	2.6	2.9	1.9	2.5 ▼	3.5 ▲	2.6	2.7	0.7 ▼	2.5
Heroin	1.6	1.7	1.1	1.5	1.7	1.8	1.7	0.8	1.7
Steroids	1.3	1.4	1.3	1.5	1.1	1.0	1.5	1.0	1.5
Injected Drugs	1.3	1.4	0.9	1.6	1.1	1.2	1.4	0.8	1.9
MDMA	3.7	5.7 ▲	2.3 ▼	2.0 ▼	3.3	4.6 ▲	2.9	1.4 ▼	3.8
Rohypnol	1.2	1.5	1.0	1.3	1.4	0.9	1.5	0.3	1.8
GHB	0.8	1.1	1.0	0.7	0.7	0.6	1.0	0.2	1.0

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 11th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	23.8	24.6	25.3	20.0 ▼	24.1	22.6 ▼	22.5	25.4	27.4 ▲
Cigars	12.8	11.5	14.0	11.3	12.3	12.7	12.9	10.5 ▼	17.4 ▲
Pipe	3.5	3.6	3.4	2.6	4.0	3.6	3.2	1.9 ▼	4.6
Smokeless Tobacco	8.0	5.7 ▼	8.2	7.2	9.5 ▲	6.2	6.3 ▼	11.3 ▲	14.3 ▲
Alcohol	35.8	39.6 ▲	33.4	32.4 ▼	35.6	33.9	31.6 ▼	40.2 ▲	42.5 ▲
Marijuana	15.7	17.5 ▲	13.3 ▼	13.2 ▼	16.2	15.6	18.0 ▲	8.7 ▼	17.9
Cocaine	2.7	3.2	2.9	2.5	2.4	2.6	2.5	1.0 ▼	3.9 ▲
Crack	1.3	1.5	1.7	1.4	1.0	1.1	1.2	0.6	2.1 ▲
Inhalants	2.3	2.8	2.4	1.9	2.5	1.9	2.5	1.0 ▼	2.8
Amphetamines	3.9	4.1	4.1	2.7	3.9	3.8	4.4	1.9 ▼	4.7
Methamphetamines	1.5	1.4	1.8	1.2	1.5	1.3	1.5	1.4	2.2
Ritalin	2.0	2.5	2.1	1.9	1.6	1.9	2.5	0.6	2.2
Tranquilizers	5.6	6.2	5.3	4.1 ▼	6.3	5.4	6.1	3.6 ▼	6.3
Narcotics	4.2	4.4	3.2	2.6 ▼	4.2	4.4	4.9	2.7 ▼	5.6 ▲
Psychedelics [#]	2.1	2.8 ▲	1.2 ▼	2.0	2.0	2.4	2.4	1.0 ▼	2.0
LSD	1.6	2.0	0.7 ▼	1.4	1.4	2.0	2.1	1.0	1.5
Others	1.3	1.9	0.8	1.4	1.3	1.4	1.4	0.3	1.3
Heroin	1.0	1.1	0.9	0.9	0.9	1.2	0.9	0.2	1.1
Steroids	0.8	0.9	1.0	1.2	0.6	0.7 ▼	1.0	0.5	1.0
Injected Drugs	0.8	0.8	0.8	1.1	0.5	0.8	0.9	0.7	1.5 ▲
MDMA	1.8	3.4	1.1	1.3	1.4	2.2	1.5	0.6	1.7
Rohypnol	0.7	1.0	0.7	0.7	0.8	0.4	0.9	0.3	1.0
GHB	0.6	0.7	0.7	0.7	0.6	0.5	0.5	0.2	0.5

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 11th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
CIGARETTES									
Daily Use	14.8	15.7	15.8	12.4 ▼	14.4	13.2 ▼	15.7	15.6	17.7 ▲
1/2+ Pack per Day	8.5	8.8	9.1	5.9 ▼	8.8	7.2 ▼	9.1	8.8	10.9 ▲
CIGARS									
	2.6	2.4	2.6	1.9	2.4	2.1	3.5 ▲	2.0	3.9 ▲
SMOKELESS TOBACCO									
	3.4	2.3 ▼	3.5	3.1	3.7	2.3 ▼	2.9	6.2 ▲	7.3 ▲
ALCOHOL									
Daily Use	3.5	3.6	3.7	1.8 ▼	3.1	2.6 ▼	3.8	5.8 ▲	4.8 ▲
Binge Drinking	21.8	25.5 ▲	21.5	17.6 ▼	20.9	19.8 ▼	20.2 ▼	24.1	26.3 ▲
MARIJUANA									
	4.9	5.5	4.4	3.4 ▼	5.2	4.8	6.1	3.4 ▼	5.1

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p<.05$).
 ▼ indicates that local prevalence rates are significantly lower than state rates ($p<.05$).



**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs
by Indiana 12th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	50.9	49.8	48.2	49.6	49.5	46.3 ▼	51.9	55.7 ▲	56.2 ▲
Cigars	37.7	37.1	38.5	42.9 ▲	38.4	34.4 ▼	34.6 ▼	39.7	40.6 ▲
Pipe	10.7	11.4	11.2	11.2	13.4 ▲	10.3	7.7 ▼	10.9	10.5
Smokeless Tobacco	20.4	15.8 ▼	18.9	22.7	21.7	15.7 ▼	18.9	27.9 ▲	24.6 ▲
Alcohol	70.2	71.9	64.5 ▼	70.8	69.4	64.9 ▼	69.7	75.4 ▲	75.5 ▲
Marijuana	37.1	39.9 ▲	30.6 ▼	39.1	35.3	37.4	37.8	34.7	39.6 ▲
Cocaine	8.6	10.3 ▲	7.7	9.7	9.6	8.4	8.0	6.6 ▼	8.5
Crack	4.0	4.0	4.6	4.4	5.2 ▲	4.0	4.1	2.8 ▼	3.5
Inhalants	10.1	10.3	12.8 ▲	10.0	11.2 ▲	8.3 ▼	10.2	11.0	8.9
Amphetamines	11.3	11.5	11.0	11.9	12.8	9.7 ▼	12.1	10.4	12.0
Methamphetamine	5.0	4.2	5.5	5.2	6.4 ▲	4.5	3.8 ▼	5.4	5.3
Ritalin	7.0	8.0	8.2	7.7	7.5	6.6	7.0	5.8	6.2
Tranquilizers	14.5	15.3	13.1	12.8	15.3	13.2	14.8	12.7	17.3 ▲
Narcotics	12.0	13.2	9.7 ▼	10.9	12.4	11.8	12.9	10.2	12.9
Psychedelics [#]	7.0	8.0 ▲	6.1	9.1 ▲	8.4	7.8	5.0 ▼	4.9 ▼	7.3
LSD	5.3	5.2	4.0	6.6 ▲	6.4	6.5 ▲	3.9 ▼	3.6 ▼	5.8
Others	4.6	6.0	3.6	5.5	6.1 ▲	5.0	2.6 ▼	3.2 ▼	4.7
Heroin	2.3	3.2 ▲	3.6 ▲	2.1	3.5 ▲	2.3	1.7	1.0 ▼	1.7
Steroids	2.3	3.1	1.6	2.5	2.9	2.4	2.1	2.0	1.9
Injected Drugs	2.2	2.7	2.2	2.1	2.9	2.7	1.8	0.9 ▼	1.9
MDMA	5.6	8.6 ▲	5.0	5.5	5.2	6.3	4.4 ▼	3.3 ▼	5.8
Rohypnol	1.7	2.3	1.6	1.2	2.3	2.2	1.1	0.9 ▼	1.4
GHB	1.3	1.5	1.0	1.0 ▼	1.9	2.1 ▲	0.6 ▼	0.5 ▼	0.9

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p<.05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p<.05$).



**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs
by Indiana 12th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	36.9	36.3	36.3	35.7	37.4	32.8 ▼	35.1	41.4 ▲	41.0 ▲
Cigars	29.1	29.5	29.7	33.8	29.3	26.5 ▼	27.4	29.0	31.6
Pipe	7.8	9.1	8.4	8.4	10.1 ▲	7.7	6.1 ▼	6.2 ▼	7.4
Smokeless Tobacco	15.4	11.8 ▼	15.4	17.6	17.3 ▲	11.2 ▼	13.7	20.6 ▲	18.5 ▲
Alcohol	61.5	63.7	55.7 ▼	63.3	61.7	54.7 ▼	59.9	67.3 ▲	67.2 ▲
Marijuana	27.5	30.3 ▲	23.1 ▼	29.4	27.1	27.1	28.8	24.0 ▼	29.3 ▲
Cocaine	6.1	7.5 ▲	5.9	7.2	6.8	5.4	5.1	4.1 ▼	7.1
Crack	2.5	2.6	3.1	2.7	3.0	2.4	2.7	1.2 ▼	2.7 ▲
Inhalants	4.6	4.7	7.4 ▲	4.3	5.6	3.7	4.0	4.7	4.2
Amphetamines	7.5	7.1	7.8	6.9	8.5	6.5	7.7	7.1	8.2
Methamphetamines	3.1	2.7	3.2	3.2	4.0	2.8	2.2	3.5	3.4
Ritalin	3.8	4.4	4.8	3.8	3.7	4.0	3.9	3.1	3.5
Tranquilizers	9.8	11.0	8.6	7.7	9.4	9.1	10.2	8.2	12.2 ▲
Narcotics	7.5	8.7	6.0	5.9	8.2	7.1	7.4	6.3	9.0 ▲
Psychedelics [#]	4.5	5.7 ▲	3.8	5.5	5.4	4.4	2.8 ▼	3.0 ▼	5.4
LSD	3.1	3.4	1.8 ▼	3.8	3.5	3.6	2.0 ▼	1.7 ▼	4.3 ▲
Others	3.2	4.8 ▲	2.8	3.1 ▼	3.9	3.1	1.7 ▼	2.5	3.4
Heroin	1.5	2.3 ▲	2.2	1.3	2.1	1.6	0.9	0.5 ▼	1.2
Steroids	1.5	2.4 ▲	1.1	1.2	1.7	1.4	1.1	1.2 ▼	1.5
Injected Drugs	1.2	2.0 ▲	0.9	1.5	1.7	1.5	0.7	0.5 ▼	0.9
MDMA	3.5	6.4 ▲	3.0	4.0	2.8	3.7	2.7	1.9 ▼	3.6
Rohypnol	1.0	1.8 ▲	0.5	0.4	1.4	1.5 ▲	0.4 ▼	0.7	0.8
GHB	0.7	1.4 ▲	0.5	0.7	1.1	1.1	0.4	0.1	0.4

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 12th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	26.9	26.5	26.3	25.7	26.8	23.2 ▼	26.3	29.7 ▲	30.8 ▲
Cigars	16.6	16.3	17.5	18.4	17.3	13.8 ▼	17.3	15.5	18.6 ▲
Pipe	4.6	5.4	5.2	5.8	6.4 ▲	4.5	3.3 ▼	3.1 ▼	4.1
Smokeless Tobacco	9.8	6.9 ▼	9.3	10.5	10.8	7.2 ▼	8.7	13.4 ▲	12.6 ▲
Alcohol	42.2	43.8	37.2 ▼	44.0	41.0	37.2 ▼	40.2	47.4 ▲	47.8 ▲
Marijuana	17.2	19.9 ▲	12.6	19.8	16.3	16.3	18.6	14.5 ▼	19.2
Cocaine	3.1	4.1 ▲	3.3	3.8	3.5	2.9	2.2	1.7 ▼	3.3
Crack	1.4	1.8	1.7	1.3	1.6	1.5	1.1	0.6 ▼	1.3
Inhalants	2.2	2.9	2.4	2.1	3.2 ▲	2.0	1.6	1.5	1.9
Amphetamines	4.1	4.1	4.1	4.1	4.3	3.5	4.6	3.5	4.7
Methamphetamines	1.5	1.8	1.5	1.3	2.1	1.4	1.3	1.7	1.3
Ritalin	1.8	1.9	2.9 ▲	2.1	1.4	2.0	1.6	1.2	1.7
Tranquilizers	5.3	6.1	5.2	4.0	4.9	5.1	5.4	4.2 ▼	6.7 ▲
Narcotics	4.0	4.4	3.0	3.1	4.1	3.7	4.2	3.4	4.9
Psychedelics [#]	2.1	2.8	1.8	2.1	2.6	2.1	1.3 ▼	1.1 ▼	2.5
LSD	1.3	1.4	0.9	1.3	1.8	1.7	0.9	0.7 ▼	1.6
Others	1.5	2.4 ▲	1.6	1.2	1.7	1.5	0.8	0.9	1.7
Heroin	1.0	1.7 ▲	1.2	1.0	1.5	1.0	0.4 ▼	0.4 ▼	0.8
Steroids	1.0	1.4	0.5	0.9	1.3	1.0	0.7	0.7	1.0
Injected Drugs	0.9	1.5 ▲	0.8	1.0	1.3	1.2	0.5	0.3	0.8
MDMA	1.7	3.5 ▲	1.3	1.9	1.5	1.7	1.1	0.9 ▼	1.3
Rohypnol	0.7	1.0	0.5	0.3	0.8	1.0	0.3	0.4	0.6
GHB	0.5	0.8	0.5	0.4	0.7	0.7	0.3	0.1	0.3

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs
by Indiana 12th Grade Students: 2006**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
CIGARETTES									
Daily Use	16.9	17.9	16.1	16.4	17.3	15.2 ▼	17.3	16.6	18.7
1/2+ Pack per Day	10.5	10.5	10.0	10.6	11.2	8.8 ▼	11.7	9.5	12.3 ▲
CIGARS									
	3.1	2.9	3.1	4.0	3.8	2.5	3.5	2.9	3.2
SMOKELESS TOBACCO									
	4.4	2.4 ▼	3.3	5.9	5.4	2.5 ▼	3.4	7.3 ▲	6.1 ▲
ALCOHOL									
Daily Use	5.6	5.6	4.4	7.2	5.9	3.7 ▼	5.1	7.6 ▲	6.4
Binge Drinking	27.3	27.4	22.8 ▼	29.6	25.6	23.8 ▼	26.9	31.7 ▲	31.4 ▲
MARIJUANA									
	6.1	7.5 ▲	4.1 ▼	5.5	6.3	5.7	7.5 ▲	4.6 ▼	6.4

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p<.05$).
 ▼ indicates that local prevalence rates are significantly lower than state rates ($p<.05$).



III. RISK AND PROTECTIVE FACTORS

• Perceptions, Grades 6-12	182
• After-School and Extra-Curricular Activities, Grades 6-12	188
• Consequences of Use, Grades 6-12	200
• Safety, Grades 6-12	206



**Perceived Risk of Selected Substance Use Behavior
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	2.6	2.7	2.6	2.6	2.7	2.6	3.0
	No Risk	8.6	7.1	6.5	5.7	5.1	4.5	4.1
	Slight Risk	14.2	11.6	11.8	10.7	9.3	8.5	7.7
	Moderate Risk	30.3	30.4	29.4	28.0	26.7	25.6	24.5
	Great Risk	44.3	48.2	49.6	53.0	56.2	58.7	60.6
Smoke marijuana occasionally	No Answer	2.9	3.0	2.6	2.8	2.8	2.8	3.1
	No Risk	8.8	8.1	10.4	13.0	14.7	16.4	17.0
	Slight Risk	11.1	12.8	17.0	20.9	24.4	26.5	26.9
	Moderate Risk	34.1	33.7	33.1	31.3	29.2	27.3	27.3
	Great Risk	43.2	42.4	36.9	32.1	29.0	27.0	25.7
Smoke marijuana regularly	No Answer	3.1	3.0	2.8	2.9	2.9	2.9	3.2
	No Risk	8.3	7.0	7.7	8.0	8.3	8.6	8.4
	Slight Risk	4.4	4.5	6.8	9.0	10.8	12.5	13.2
	Moderate Risk	14.2	14.2	16.8	19.7	22.4	23.8	24.2
	Great Risk	70.0	71.3	65.8	60.3	55.6	52.2	51.0
Use cocaine occasionally	No Answer	3.1	3.2	2.9	3.1	3.0	3.0	3.3
	No Risk	8.4	6.5	5.9	4.8	4.6	4.2	3.9
	Slight Risk	8.9	7.8	7.4	7.0	5.8	6.1	5.3
	Moderate Risk	33.7	32.8	31.0	28.9	26.6	24.3	22.7
	Great Risk	45.9	49.7	52.9	56.2	59.9	62.4	64.9
Use cocaine regularly	No Answer	4.8	4.5	4.7	4.6	4.5	4.5	4.6
	No Risk	8.2	6.5	5.8	4.7	4.3	3.8	3.3
	Slight Risk	3.5	2.5	2.6	2.3	2.0	1.9	1.6
	Moderate Risk	11.8	9.5	8.9	8.5	7.2	6.7	6.3
	Great Risk	71.6	76.9	78.0	80.0	82.1	83.1	84.2
Occasionally consume 1-2 alcoholic drinks	No Answer	3.6	3.7	3.5	3.5	3.5	3.1	3.4
	No Risk	21.2	22.0	26.5	31.1	32.0	35.6	37.3
	Slight Risk	39.8	39.5	39.5	38.1	38.6	37.8	36.9
	Moderate Risk	19.6	19.5	17.1	14.7	14.3	12.6	11.6
	Great Risk	15.8	15.3	13.4	12.6	11.6	10.9	10.8
Binge drink weekly	No Answer	3.0	3.0	2.9	2.9	3.0	2.8	3.2
	No Risk	9.6	8.5	8.9	9.2	8.8	8.9	9.3
	Slight Risk	12.1	12.6	15.2	16.9	17.5	19.3	20.1
	Moderate Risk	30.0	29.4	31.4	31.5	32.6	31.9	31.9
	Great Risk	45.3	46.5	41.6	39.4	38.1	37.1	35.4

General

**Perceived Risk of Selected Substance Use Behavior
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	3.2	3.5	3.4	3.3	3.5	3.6	3.9
	No Risk	10.1	8.5	8.0	7.4	6.8	6.1	5.4
	Slight Risk	14.2	11.4	11.7	11.5	9.9	9.3	8.6
	Moderate Risk	31.2	30.9	30.1	28.5	28.4	26.9	26.4
	Great Risk	41.3	45.7	46.8	49.3	51.5	54.1	55.8
Smoke marijuana occasionally	No Answer	3.5	3.6	3.3	3.5	3.6	3.6	3.9
	No Risk	10.4	9.7	12.4	16.2	18.5	20.8	22.7
	Slight Risk	10.9	13.0	17.5	21.8	25.6	27.7	28.0
	Moderate Risk	32.6	32.2	30.6	27.7	25.7	22.7	23.1
	Great Risk	42.6	41.5	36.1	30.7	26.6	25.2	22.4
Smoke marijuana regularly	No Answer	3.6	3.7	3.5	3.7	3.7	3.7	3.9
	No Risk	9.9	8.6	9.6	10.7	11.0	11.5	11.5
	Slight Risk	4.5	4.7	7.4	10.0	12.6	14.6	16.0
	Moderate Risk	14.1	14.7	17.5	20.9	23.4	24.5	25.4
	Great Risk	67.9	68.3	62.1	54.7	49.3	45.6	43.2
Use cocaine occasionally	No Answer	3.6	3.8	3.5	3.8	3.8	3.9	4.0
	No Risk	9.9	7.9	7.4	6.4	6.4	5.9	5.2
	Slight Risk	8.6	6.9	6.9	7.0	6.2	6.9	6.6
	Moderate Risk	31.8	31.1	28.5	26.4	25.2	23.4	22.7
	Great Risk	46.1	50.3	53.6	56.4	58.5	59.9	61.4
Use cocaine regularly	No Answer	5.0	4.9	5.1	5.0	5.0	5.0	5.2
	No Risk	9.6	7.9	7.3	6.3	6.1	5.2	4.4
	Slight Risk	3.4	2.4	2.6	2.4	2.3	2.5	1.9
	Moderate Risk	11.8	8.9	8.6	8.6	7.5	7.1	7.3
	Great Risk	70.2	75.9	76.4	77.6	79.1	80.3	81.2
Occasionally consume 1-2 alcoholic drinks	No Answer	4.2	4.2	4.0	4.1	4.1	4.0	4.1
	No Risk	23.2	23.6	28.2	33.5	36.1	41.0	44.3
	Slight Risk	39.2	38.8	38.2	35.8	35.4	33.1	32.1
	Moderate Risk	17.8	17.5	14.9	12.9	12.1	10.2	8.5
	Great Risk	15.7	15.9	14.6	13.7	12.3	11.8	11.1
Binge drink weekly	No Answer	3.6	3.6	3.6	3.8	3.7	3.9	4.0
	No Risk	11.3	10.0	10.6	12.1	11.7	12.2	13.4
	Slight Risk	12.5	12.9	15.8	17.5	19.7	22.1	22.9
	Moderate Risk	30.1	29.4	31.4	30.4	31.6	29.9	30.4
	Great Risk	42.5	44.1	38.7	36.2	33.2	31.9	29.3

Male



**Perceived Risk of Selected Substance Use Behavior
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	2.0	2.0	1.8	1.9	1.9	1.7	1.9
	No Risk	7.0	5.6	5.1	4.0	3.5	3.0	2.8
	Slight Risk	14.3	11.8	11.9	10.0	8.8	7.8	6.8
	Moderate Risk	29.4	29.9	28.8	27.5	25.4	24.5	22.8
	Great Risk	47.3	50.8	52.4	56.6	60.4	63.0	65.6
Smoke marijuana occasionally	No Answer	2.3	2.3	2.0	2.0	2.0	1.9	2.0
	No Risk	7.1	6.4	8.3	10.0	11.1	12.4	11.4
	Slight Risk	11.1	12.7	16.5	20.0	23.3	25.6	26.0
	Moderate Risk	35.7	35.2	35.6	34.6	32.4	31.3	31.5
	Great Risk	43.7	43.3	37.7	33.4	31.1	28.8	29.1
Smoke marijuana regularly	No Answer	2.6	2.3	2.1	2.1	2.1	2.1	2.3
	No Risk	6.7	5.4	5.8	5.5	5.8	5.9	5.2
	Slight Risk	4.3	4.2	6.3	8.1	9.2	10.4	10.5
	Moderate Risk	14.2	13.7	16.2	18.6	21.4	23.2	23.1
	Great Risk	72.2	74.4	69.6	65.7	61.4	58.4	58.8
Use cocaine occasionally	No Answer	2.6	2.5	2.2	2.3	2.2	2.1	2.2
	No Risk	6.8	5.1	4.3	3.3	2.9	2.6	2.5
	Slight Risk	9.3	8.6	7.8	7.0	5.5	5.4	4.1
	Moderate Risk	35.6	34.6	33.4	31.2	28.0	25.1	22.7
	Great Risk	45.7	49.2	52.3	56.2	61.3	64.7	68.5
Use cocaine regularly	No Answer	4.7	4.1	4.2	4.1	3.9	4.1	3.8
	No Risk	6.7	5.0	4.2	3.1	2.6	2.4	2.1
	Slight Risk	3.6	2.6	2.5	2.1	1.7	1.5	1.3
	Moderate Risk	11.8	10.1	9.3	8.4	7.0	6.3	5.4
	Great Risk	73.2	78.1	79.8	82.4	84.8	85.8	87.4
Occasionally consume 1-2 alcoholic drinks	No Answer	3.1	3.2	3.0	2.8	2.8	2.3	2.5
	No Risk	19.2	20.4	24.7	28.9	28.3	30.7	30.5
	Slight Risk	40.4	40.3	40.8	40.3	41.5	42.1	41.8
	Moderate Risk	21.4	21.5	19.2	16.4	16.3	14.9	14.7
	Great Risk	15.9	14.6	12.3	11.5	11.0	10.1	10.5
Binge drink weekly	No Answer	2.4	2.4	2.1	2.1	2.2	1.9	2.1
	No Risk	7.9	6.9	7.2	6.5	6.1	5.8	5.2
	Slight Risk	11.7	12.3	14.6	16.3	15.5	16.9	17.4
	Moderate Risk	29.9	29.5	31.4	32.6	33.4	33.6	33.6
	Great Risk	48.1	48.8	44.7	42.5	42.7	41.9	41.6

Female

**Perceived Peer Approval of Using Selected Substances
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	4.7	4.0	3.6	3.5	3.5	3.5	3.8
	Strongly Approve	4.9	4.4	4.2	3.9	3.7	3.7	3.8
	Approve	1.8	3.4	5.0	6.8	8.5	9.2	9.0
	Do Not Know	12.7	14.1	17.2	17.8	18.2	17.8	18.5
	Disapprove	15.6	15.6	18.2	19.7	20.5	21.3	22.7
	Strongly Disapprove	60.2	58.6	51.8	48.3	45.6	44.5	42.2
Smoke marijuana occasionally	No Answer	4.8	4.3	3.7	3.4	3.6	3.5	3.8
	Strongly Approve	5.1	4.8	5.5	6.2	6.5	6.8	7.1
	Approve	1.7	3.7	7.4	10.8	13.8	15.6	15.3
	Do Not Know	9.8	11.3	14.8	16.2	16.7	16.9	17.8
	Disapprove	11.5	11.3	13.4	15.1	16.4	16.6	17.1
	Strongly Disapprove	67.2	64.7	55.2	48.3	43.1	40.6	38.9
Smoke marijuana regularly	No Answer	4.9	4.4	3.8	3.6	3.8	3.6	3.9
	Strongly Approve	5.1	4.8	5.6	6.2	6.3	6.7	6.7
	Approve	1.2	2.4	4.8	6.5	8.4	9.2	8.5
	Do Not Know	9.1	10.8	13.7	14.9	15.3	15.1	15.2
	Disapprove	7.7	7.8	10.4	12.5	13.7	14.6	15.9
	Strongly Disapprove	72.0	69.8	61.7	56.3	52.5	50.8	49.7
Use cocaine occasionally	No Answer	5.0	4.6	4.0	3.6	3.7	3.8	3.9
	Strongly Approve	4.8	3.9	3.6	3.1	2.8	2.8	2.6
	Approve	0.8	1.3	1.9	2.1	2.1	2.4	2.4
	Do Not Know	9.2	10.3	12.0	12.8	12.1	11.6	11.4
	Disapprove	10.7	10.3	11.9	12.7	12.7	12.2	11.9
	Strongly Disapprove	69.5	69.6	66.7	65.7	66.6	67.1	67.8
Use cocaine regularly	No Answer	5.4	5.2	4.8	4.4	4.6	4.8	4.6
	Strongly Approve	4.8	4.0	3.7	3.1	2.7	2.8	2.5
	Approve	0.7	1.1	1.5	1.5	1.6	1.6	1.5
	Do Not Know	8.8	9.9	11.5	12.3	11.5	11.0	10.7
	Disapprove	6.9	6.8	8.3	9.3	9.3	8.7	8.7
	Strongly Disapprove	73.4	73.1	70.3	69.4	70.3	71.2	72.1
Occasionally consume 1-2 alcoholic drinks	No Answer	5.2	4.6	4.1	3.7	3.8	3.7	3.9
	Strongly Approve	5.5	5.9	7.5	10.0	11.5	12.4	14.8
	Approve	6.8	10.6	17.4	24.9	28.8	32.2	34.0
	Do Not Know	17.7	19.0	23.0	22.7	22.2	21.3	19.9
	Disapprove	15.8	15.6	14.5	13.4	12.5	11.6	10.7
	Strongly Disapprove	49.1	44.3	33.4	25.3	21.3	18.7	16.7
Binge drink weekly	No Answer	5.1	4.5	4.1	3.7	3.6	3.7	3.9
	Strongly Approve	5.3	5.3	6.4	8.3	9.2	10.0	11.3
	Approve	2.2	4.6	8.3	12.9	16.9	19.4	21.0
	Do Not Know	12.1	14.0	19.0	21.1	20.7	20.4	20.4
	Disapprove	10.9	11.5	14.3	14.9	14.8	15.3	14.8
	Strongly Disapprove	64.3	60.1	47.9	39.1	34.7	31.2	28.5

General



**Perceived Peer Approval of Using Selected Substances
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	5.5	4.9	4.7	4.7	4.4	4.6	4.8
	Strongly Approve	5.9	5.4	5.3	5.2	4.9	4.9	5.2
	Approve	2.1	3.6	4.5	6.7	8.8	9.7	10.0
	Do Not Know	14.7	16.8	19.8	21.5	21.6	20.7	22.3
	Disapprove	17.2	16.4	18.7	20.1	21.2	21.3	22.7
	Strongly Disapprove	54.6	52.9	47.0	41.8	39.1	38.8	35.0
Smoke marijuana occasionally	No Answer	5.7	5.2	4.9	4.6	4.5	4.6	4.8
	Strongly Approve	6.1	6.0	6.6	8.1	8.3	9.0	9.6
	Approve	2.0	3.9	7.1	10.5	14.1	15.6	16.1
	Do Not Know	11.8	13.8	17.8	19.8	20.0	19.6	20.7
	Disapprove	12.3	12.0	13.4	15.6	16.4	15.5	17.0
	Strongly Disapprove	62.1	59.1	50.3	41.4	36.7	35.7	31.8
Smoke marijuana regularly	No Answer	5.9	5.4	5.0	4.7	4.8	4.5	5.0
	Strongly Approve	6.1	6.1	6.7	8.2	8.4	9.2	9.2
	Approve	1.6	2.6	4.7	6.8	9.0	9.9	9.9
	Do Not Know	10.8	13.2	16.6	18.6	18.8	18.0	18.0
	Disapprove	8.4	8.4	11.0	13.1	13.9	14.2	16.4
	Strongly Disapprove	67.1	64.4	56.0	48.6	45.0	44.2	41.5
Use cocaine occasionally	No Answer	6.1	5.4	5.1	4.7	4.6	4.8	4.9
	Strongly Approve	5.7	5.0	4.6	4.4	4.0	4.1	3.6
	Approve	1.1	1.3	1.9	2.3	2.6	2.9	3.0
	Do Not Know	10.9	12.6	14.7	16.4	15.5	14.7	14.9
	Disapprove	11.4	11.1	12.0	13.1	13.1	13.0	12.6
	Strongly Disapprove	64.9	64.6	61.7	59.2	60.3	60.4	61.0
Use cocaine regularly	No Answer	6.3	6.0	5.7	5.4	5.2	5.7	5.4
	Strongly Approve	5.8	5.1	4.8	4.4	3.9	4.0	3.4
	Approve	0.8	1.1	1.5	1.8	2.1	1.9	2.0
	Do Not Know	10.5	12.1	14.3	16.1	15.0	14.1	14.0
	Disapprove	7.8	7.5	8.7	9.8	9.8	9.5	9.2
	Strongly Disapprove	68.8	68.2	65.0	62.6	64.0	64.8	66.0
Occasionally consume 1-2 alcoholic drinks	No Answer	6.1	5.3	5.1	4.8	4.6	4.8	4.9
	Strongly Approve	6.3	7.0	8.2	11.3	13.6	14.9	18.3
	Approve	7.3	9.8	15.8	22.3	26.8	29.8	31.5
	Do Not Know	19.5	20.8	25.0	25.0	23.4	21.8	21.2
	Disapprove	15.8	15.5	13.7	12.5	11.2	10.1	9.1
	Strongly Disapprove	45.0	41.6	32.2	24.0	20.5	18.6	14.9
Binge drink weekly	No Answer	6.1	5.5	5.3	4.9	4.7	4.8	5.1
	Strongly Approve	6.3	6.5	7.6	10.3	12.2	13.4	15.8
	Approve	2.6	4.6	8.0	12.6	17.5	19.9	21.4
	Do Not Know	13.9	16.3	21.1	23.7	22.5	21.7	21.7
	Disapprove	12.3	11.9	14.3	13.9	13.4	13.4	13.1
	Strongly Disapprove	58.7	55.2	43.7	34.6	29.8	26.9	22.9

Male

**Perceived Peer Approval of Using Selected Substances
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	3.9	3.1	2.5	2.3	2.5	2.5	2.5
	Strongly Approve	3.9	3.3	3.1	2.6	2.6	2.6	2.3
	Approve	1.6	3.2	5.5	6.9	8.3	8.7	8.1
	Do Not Know	10.7	11.3	14.5	14.3	15.1	15.2	14.8
	Disapprove	14.0	14.7	17.8	19.4	19.9	21.3	22.8
	Strongly Disapprove	66.0	64.5	56.7	54.5	51.6	49.7	49.5
Smoke marijuana occasionally	No Answer	3.9	3.5	2.5	2.3	2.6	2.6	2.5
	Strongly Approve	4.0	3.5	4.4	4.5	4.7	4.8	4.6
	Approve	1.4	3.4	7.8	11.1	13.5	15.6	14.4
	Do Not Know	7.7	8.8	11.8	12.7	13.8	14.3	15.0
	Disapprove	10.7	10.5	13.3	14.6	16.3	17.6	17.4
	Strongly Disapprove	72.3	70.3	60.3	54.9	49.0	45.1	46.0
Smoke marijuana regularly	No Answer	3.8	3.5	2.6	2.5	2.9	2.8	2.6
	Strongly Approve	4.0	3.5	4.4	4.3	4.4	4.4	4.2
	Approve	0.8	2.2	4.8	6.3	7.8	8.5	7.0
	Do Not Know	7.4	8.3	10.8	11.5	12.1	12.4	12.5
	Disapprove	6.9	7.3	9.8	11.8	13.4	14.9	15.7
	Strongly Disapprove	77.1	75.4	67.6	63.6	59.4	56.9	58.0
Use cocaine occasionally	No Answer	4.0	3.7	2.8	2.6	2.8	2.8	2.6
	Strongly Approve	3.7	2.8	2.5	1.9	1.7	1.6	1.5
	Approve	0.6	1.3	1.9	1.9	1.8	2.0	1.8
	Do Not Know	7.4	8.0	9.1	9.5	9.0	8.7	8.0
	Disapprove	10.1	9.6	11.8	12.4	12.3	11.5	11.3
	Strongly Disapprove	74.2	74.6	71.8	71.9	72.4	73.4	74.7
Use cocaine regularly	No Answer	4.5	4.4	3.8	3.4	4.1	3.9	3.5
	Strongly Approve	3.7	2.8	2.6	1.8	1.6	1.6	1.5
	Approve	0.6	1.0	1.5	1.3	1.1	1.3	0.9
	Do Not Know	7.2	7.6	8.6	8.7	8.3	8.2	7.4
	Disapprove	5.9	6.0	8.0	8.9	8.8	7.8	8.3
	Strongly Disapprove	78.1	78.1	75.6	75.9	76.2	77.1	78.4
Occasionally consume 1-2 alcoholic drinks	No Answer	4.1	3.9	3.1	2.7	2.9	2.8	2.7
	Strongly Approve	4.5	4.8	6.8	8.7	9.5	10.1	11.4
	Approve	6.4	11.4	19.1	27.3	30.7	34.3	36.5
	Do Not Know	15.9	17.1	21.1	20.5	21.1	20.9	18.8
	Disapprove	15.7	15.7	15.4	14.2	13.7	13.0	12.3
	Strongly Disapprove	53.3	47.1	34.6	26.6	22.1	19.0	18.3
Binge drink weekly	No Answer	4.1	3.6	2.7	2.5	2.6	2.6	2.5
	Strongly Approve	4.3	4.0	5.2	6.4	6.5	6.9	6.8
	Approve	1.8	4.5	8.6	13.1	16.4	18.9	20.7
	Do Not Know	10.3	11.7	16.9	18.6	19.1	19.2	19.2
	Disapprove	9.5	11.2	14.3	15.8	16.1	17.0	16.7
	Strongly Disapprove	70.0	65.0	52.2	43.5	39.4	35.3	34.1

Female



**Number of Days per School Year Spent in Various Afterschool Activities
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
School sports team	No Answer	6.1	6.1	5.9	6.2	6.1	6.3	6.3
	None	40.2	36.6	39.1	41.1	46.1	49.6	52.8
	1-9 Days	17.3	12.3	10.1	6.5	4.9	3.8	3.3
	10-29 Days	9.9	7.3	5.9	3.3	2.4	2.0	2.1
	30-59 Days	9.8	11.7	10.9	8.0	7.0	6.1	5.8
	60-89 Days	7.2	10.1	9.7	10.3	10.0	9.7	9.7
	90-129 Days	5.3	8.8	9.9	12.5	11.8	11.7	10.5
	130-180 Days	4.1	7.1	8.5	12.1	11.7	10.8	9.5
School arts programs	No Answer	6.8	6.7	6.5	6.8	6.8	6.9	7.0
	None	64.5	61.7	59.9	60.6	61.4	61.3	62.6
	1-9 Days	17.6	17.8	17.6	14.3	12.5	11.0	9.7
	10-29 Days	4.5	5.7	6.3	5.8	5.9	5.3	5.4
	30-59 Days	2.6	3.0	3.7	4.1	4.2	4.6	4.5
	60-89 Days	1.3	1.8	2.3	3.2	3.5	3.9	3.9
	90-129 Days	1.0	1.1	1.3	2.3	2.7	3.2	3.4
	130-180 Days	1.7	2.3	2.5	2.9	3.1	3.8	3.5
School club/ intramurals	No Answer	7.7	7.5	7.1	7.3	7.2	7.4	7.2
	None	64.4	65.1	63.8	61.3	57.8	53.0	52.8
	1-9 Days	14.7	13.2	13.8	14.2	14.7	14.5	14.6
	10-29 Days	6.3	6.5	6.9	7.9	9.0	11.0	11.5
	30-59 Days	3.2	3.5	3.7	4.1	5.1	6.5	6.4
	60-89 Days	1.4	2.0	2.0	2.2	2.9	3.7	3.8
	90-129 Days	1.0	1.1	1.3	1.6	1.6	2.0	2.1
	130-180 Days	1.3	1.2	1.4	1.3	1.6	1.9	1.6
Youth centers (YMCA, etc)	No Answer	8.1	7.6	7.2	7.2	7.1	7.2	7.2
	None	72.7	74.5	74.7	77.5	77.7	77.9	79.1
	1-9 Days	10.2	8.8	9.2	7.7	7.2	6.8	6.2
	10-29 Days	3.7	3.6	3.4	3.0	3.0	3.2	3.0
	30-59 Days	2.1	2.2	2.2	2.1	2.2	2.0	2.1
	60-89 Days	1.1	1.2	1.3	1.1	1.3	1.3	1.0
	90-129 Days	0.9	0.9	0.8	0.7	0.7	0.5	0.5
	130-180 Days	1.3	1.2	1.2	0.8	0.8	0.9	0.7
Extended day programs	No Answer	9.2	8.3	7.8	7.7	7.6	7.7	7.6
	None	70.5	73.6	74.4	78.7	80.5	81.2	82.6
	1-9 Days	14.1	12.5	12.0	9.1	7.9	6.9	5.8
	10-29 Days	3.0	2.6	2.9	2.1	1.8	1.8	1.5
	30-59 Days	1.2	1.2	1.2	0.9	0.8	1.1	0.8
	60-89 Days	0.7	0.6	0.6	0.6	0.5	0.5	0.5
	90-129 Days	0.5	0.4	0.4	0.4	0.4	0.2	0.3
	130-180 Days	0.8	0.8	0.7	0.6	0.5	0.6	0.8

Note: The number of school days in a year is estimated as 180 days.

General



**Number of Days per School Year Spent in Various Afterschool Activities
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Supervised youth activities (4H, scouts)	No Answer	9.5	9.1	8.9	8.7	8.6	8.6	8.4
	None	60.3	63.6	66.5	70.0	71.4	72.2	73.7
	1-9 Days	11.3	8.7	8.1	6.9	6.9	6.4	6.2
	10-29 Days	6.5	6.0	5.2	4.9	4.6	4.8	4.7
	30-59 Days	5.1	5.2	4.7	3.8	3.7	3.7	2.9
	60-89 Days	3.0	3.0	2.8	2.7	2.2	2.0	1.9
	90-129 Days	2.0	2.1	1.7	1.4	1.2	0.9	1.1
Afterschool classes (art, sport, etc)	130-180 Days	2.2	2.2	2.1	1.5	1.4	1.5	1.1
	No Answer	8.8	8.0	7.9	7.7	7.7	7.8	7.5
	None	46.7	48.3	50.8	53.4	56.1	57.9	62.0
	1-9 Days	18.0	15.6	14.4	12.1	11.3	10.3	9.3
	10-29 Days	7.9	7.7	7.3	6.7	6.6	6.3	5.6
	30-59 Days	6.1	6.3	6.0	5.9	5.5	5.2	4.6
	60-89 Days	4.2	4.6	4.4	4.5	4.0	3.9	3.3
Socialization without adult supervision	90-129 Days	3.2	3.6	3.7	3.9	3.5	3.2	2.6
	130-180 Days	5.1	5.9	5.6	5.8	5.4	5.5	5.1
	No Answer	7.5	7.3	6.8	6.5	6.5	6.8	6.5
	None	37.6	26.1	17.9	13.8	10.9	8.5	8.5
	1-9 Days	23.0	21.3	18.2	14.8	11.7	8.3	7.8
	10-29 Days	11.6	13.9	14.8	14.2	12.5	11.1	11.0
	30-59 Days	6.9	9.5	12.0	14.2	14.4	14.6	14.6
At home without adult supervision	60-89 Days	4.3	7.6	9.5	12.4	13.6	15.1	15.5
	90-129 Days	3.5	5.8	8.8	11.1	14.1	16.3	16.4
	130-180 Days	5.6	8.5	11.8	13.0	16.2	19.2	19.6
	No Answer	8.6	8.4	7.8	7.4	7.1	7.3	7.2
	None	28.9	20.6	16.1	12.4	11.1	9.7	9.7
	1-9 Days	28.9	24.3	20.1	16.6	14.3	11.9	11.1
	10-29 Days	13.4	15.3	16.2	16.0	15.5	15.5	14.5
At home with adult supervision	30-59 Days	7.4	10.8	12.6	15.3	15.5	16.1	16.5
	60-89 Days	4.3	7.0	9.6	11.9	13.4	14.1	14.0
	90-129 Days	3.3	5.6	7.4	9.5	11.0	12.0	12.3
	130-180 Days	5.2	8.0	10.2	10.9	12.2	13.4	14.8
	No Answer	9.4	8.9	8.4	7.9	7.7	7.7	7.5
	None	13.5	11.5	11.5	10.5	10.1	10.0	11.4
	1-9 Days	13.2	12.2	12.1	10.6	10.0	9.4	9.6

Note: The number of school days in a year is estimated as 180 days.

General



**Number of Days per School Year Spent in Various Afterschool Activities
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Work for pay outside home	No Answer	8.6	8.2	7.5	7.4	7.3	7.1	7.0
	None	55.0	49.8	46.8	48.1	40.4	29.4	25.2
	1-9 Days	18.2	18.0	16.7	12.9	9.2	6.2	5.1
	10-29 Days	8.7	10.5	11.5	10.5	8.9	7.2	6.6
	30-59 Days	4.2	6.1	7.9	8.6	10.2	10.6	10.1
	60-89 Days	2.0	3.1	4.2	5.5	9.1	13.0	13.0
	90-129 Days	1.3	1.8	2.4	3.4	7.5	13.2	15.0
Work for pay inside a home	130-180 Days	2.0	2.4	2.9	3.6	7.4	13.2	18.0
	No Answer	8.5	8.1	7.5	7.5	7.5	7.6	7.5
	None	44.0	41.9	42.8	46.4	50.9	54.8	59.7
	1-9 Days	22.0	20.2	18.9	16.7	15.1	14.3	12.7
	10-29 Days	11.2	12.0	12.2	10.9	10.0	8.9	7.9
	30-59 Days	5.5	7.3	7.5	7.4	6.6	6.0	4.8
	60-89 Days	2.9	3.7	3.9	4.0	3.5	3.0	2.7
Volunteer outside the home	90-129 Days	1.9	2.5	2.6	2.6	2.3	2.2	1.8
	130-180 Days	4.1	4.2	4.5	4.4	4.0	3.3	2.9
	No Answer	8.6	8.2	7.4	7.6	7.3	7.4	7.2
	None	55.4	54.3	54.2	54.3	53.6	50.0	48.3
	1-9 Days	20.0	19.0	18.6	17.2	16.6	16.3	16.2
	10-29 Days	7.7	8.5	9.1	9.8	10.6	12.1	13.1
	30-59 Days	3.7	4.5	4.9	5.3	6.1	6.9	7.7
Volunteer inside a home	60-89 Days	1.8	2.2	2.5	2.6	2.7	3.4	3.4
	90-129 Days	1.2	1.3	1.4	1.5	1.5	1.9	1.9
	130-180 Days	1.6	1.8	1.8	1.6	1.6	1.9	2.1
	No Answer	8.3	8.1	7.4	7.4	7.3	7.3	7.1
	None	45.8	44.8	46.4	48.2	49.1	49.9	51.3
	1-9 Days	22.0	20.3	18.8	16.3	15.1	14.0	13.5
	10-29 Days	9.7	10.3	10.2	9.6	9.3	9.1	8.9

Note: The number of school days in a year is estimated as 180 days.

**Number of Days per School Year Spent in Various Afterschool Activities
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
School sports team	No Answer	7.2	7.4	7.4	7.8	7.5	8.2	8.2
	None	40.4	35.9	36.7	38.8	43.5	45.3	48.0
	1-9 Days	15.9	11.4	9.4	5.8	4.9	3.7	3.3
	10-29 Days	9.6	6.8	5.9	3.3	2.6	2.2	2.3
	30-59 Days	10.0	11.8	11.7	8.6	7.5	6.4	6.3
	60-89 Days	7.4	10.3	9.7	10.5	9.7	10.2	10.4
	90-129 Days	5.1	8.6	9.9	12.1	11.2	11.5	11.0
	130-180 Days	4.5	7.7	9.3	12.9	13.1	12.5	10.5
School arts programs	No Answer	7.9	8.1	8.2	8.5	8.4	9.0	8.8
	None	70.9	68.9	66.8	68.6	68.8	66.8	67.5
	1-9 Days	13.8	13.4	13.3	9.3	8.8	8.4	7.3
	10-29 Days	2.9	3.8	4.5	4.3	4.3	4.4	4.4
	30-59 Days	1.7	2.0	2.7	3.1	3.3	3.6	3.9
	60-89 Days	0.8	1.1	1.5	2.3	2.4	3.0	2.9
	90-129 Days	0.7	0.8	0.9	1.6	1.7	2.1	2.3
	130-180 Days	1.4	2.0	2.2	2.3	2.4	2.8	2.9
School club/ intramurals	No Answer	8.6	8.8	8.7	9.1	8.8	9.5	9.1
	None	68.2	69.8	68.2	66.9	64.0	57.4	57.6
	1-9 Days	11.9	10.3	11.0	10.0	10.9	11.9	11.7
	10-29 Days	5.6	5.0	5.7	6.7	7.6	9.9	10.5
	30-59 Days	2.8	2.7	2.7	3.2	4.1	5.4	5.3
	60-89 Days	1.1	1.5	1.4	1.6	2.1	2.9	2.8
	90-129 Days	0.7	0.7	0.9	1.1	1.1	1.3	1.4
	130-180 Days	1.1	1.1	1.3	1.3	1.5	1.7	1.6
Youth centers (YMCA, etc)	No Answer	9.0	8.9	8.8	9.0	8.7	9.0	9.1
	None	74.7	75.9	75.0	77.7	77.3	76.6	77.9
	1-9 Days	8.5	7.2	7.7	6.0	6.3	6.5	5.5
	10-29 Days	3.1	3.0	3.0	2.8	2.8	3.1	3.0
	30-59 Days	1.9	1.9	2.2	2.0	2.0	2.2	2.0
	60-89 Days	1.0	1.0	1.1	1.0	1.2	1.2	1.1
	90-129 Days	0.7	0.8	0.8	0.6	0.6	0.4	0.5
	130-180 Days	1.1	1.2	1.4	1.0	1.0	1.1	0.9
Extended day programs	No Answer	10.0	9.5	9.2	9.3	9.2	9.2	9.4
	None	73.4	75.7	76.2	79.4	80.9	81.2	82.6
	1-9 Days	11.4	9.8	9.6	7.1	6.1	5.4	4.7
	10-29 Days	2.5	2.2	2.4	1.7	1.6	1.6	1.2
	30-59 Days	1.0	1.1	1.0	0.9	0.7	1.0	0.6
	60-89 Days	0.6	0.6	0.6	0.5	0.5	0.5	0.5
	90-129 Days	0.4	0.3	0.3	0.4	0.3	0.2	0.2
	130-180 Days	0.7	0.8	0.8	0.8	0.6	0.9	0.8

Note: The number of school days in a year is estimated as 180 days.

Male



**Number of Days per School Year Spent in Various Afterschool Activities
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Supervised youth activities (4H, scouts)	No Answer	9.9	9.7	9.9	9.9	9.8	9.6	9.8
	None	61.7	66.0	68.0	72.0	73.0	73.5	73.9
	1-9 Days	9.3	6.8	6.8	5.2	5.6	5.6	5.2
	10-29 Days	6.0	5.3	4.5	4.1	3.9	4.2	4.6
	30-59 Days	5.2	5.0	4.7	3.5	3.4	3.5	2.6
	60-89 Days	3.4	3.0	2.5	2.4	1.9	1.4	1.8
	90-129 Days	2.0	1.8	1.3	1.3	1.1	0.7	0.8
Afterschool classes (art, sport, etc)	130-180 Days	2.4	2.4	2.2	1.7	1.4	1.5	1.3
	No Answer	9.6	9.4	9.2	9.6	9.3	9.5	9.6
	None	55.4	57.5	60.0	62.3	64.1	63.9	66.5
	1-9 Days	14.8	12.4	11.4	9.3	8.8	8.7	7.6
	10-29 Days	6.8	6.4	6.2	5.3	5.6	5.2	4.9
	30-59 Days	4.8	4.9	4.5	4.5	4.1	4.1	3.6
	60-89 Days	3.0	3.1	2.8	2.8	2.5	2.7	2.0
Socialization without adult supervision	90-129 Days	2.0	2.1	2.0	2.4	1.8	1.8	1.4
	130-180 Days	3.7	4.2	3.9	3.8	3.9	4.1	4.4
	No Answer	8.7	8.8	8.6	8.4	8.2	8.7	8.5
	None	39.9	30.5	22.3	18.1	14.6	11.0	11.5
	1-9 Days	21.4	19.4	17.1	13.9	10.8	7.3	6.6
	10-29 Days	10.6	13.2	13.8	13.5	12.1	10.9	9.9
	30-59 Days	6.5	8.6	11.4	13.6	14.4	14.3	13.6
At home without adult supervision	60-89 Days	4.0	6.8	8.5	10.9	12.2	13.8	14.4
	90-129 Days	3.3	4.9	7.5	9.3	11.8	14.8	14.9
	130-180 Days	5.6	7.9	10.9	12.4	15.9	19.1	20.6
	No Answer	9.5	10.0	9.4	9.3	8.9	9.1	9.3
	None	30.5	24.2	19.8	16.4	14.5	12.2	12.3
	1-9 Days	26.9	22.6	18.4	15.3	13.5	10.5	9.5
	10-29 Days	13.4	14.4	15.4	15.2	15.2	15.4	13.5
At home with adult supervision	30-59 Days	7.1	9.9	11.9	14.2	14.5	16.2	16.0
	60-89 Days	4.0	6.3	8.8	11.1	12.3	13.1	13.8
	90-129 Days	3.0	4.8	6.4	8.0	9.3	10.3	10.9
	130-180 Days	5.6	7.8	9.8	10.6	11.8	13.3	14.7
	No Answer	10.1	10.0	9.9	9.7	9.4	9.5	9.3
	None	16.9	15.4	15.4	14.9	14.1	13.2	14.5
	1-9 Days	12.4	11.6	11.5	10.3	9.9	9.1	9.0

Note: The number of school days in a year is estimated as 180 days.

Male

**Number of Days per School Year Spent in Various Afterschool Activities
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Work for pay outside home	No Answer	9.4	9.6	9.1	9.1	8.9	8.9	8.8
	None	51.7	46.9	42.7	44.9	37.6	28.8	25.8
	1-9 Days	18.0	16.9	15.6	11.7	8.5	5.8	4.9
	10-29 Days	10.0	10.7	11.7	10.8	9.0	7.9	6.8
	30-59 Days	4.6	6.7	8.8	9.1	11.0	10.9	10.5
	60-89 Days	2.5	3.5	5.1	6.0	9.4	13.2	11.9
	90-129 Days	1.5	2.4	2.9	3.6	7.1	11.1	13.3
Work for pay inside a home	130-180 Days	2.3	3.3	4.2	4.8	8.5	13.4	17.9
	No Answer	9.5	9.6	9.1	9.2	9.3	9.6	9.5
	None	49.4	48.4	49.8	53.5	57.8	60.5	65.6
	1-9 Days	19.4	18.0	16.5	14.3	13.0	12.4	10.0
	10-29 Days	9.9	9.5	9.8	8.2	7.7	6.5	5.9
	30-59 Days	4.5	5.6	5.9	5.8	4.6	4.5	3.1
	60-89 Days	2.3	2.8	2.9	2.6	2.4	2.1	1.9
Volunteer outside the home	90-129 Days	1.3	1.9	1.8	2.1	1.4	1.5	1.2
	130-180 Days	3.6	4.1	4.2	4.2	3.9	2.9	2.9
	No Answer	9.5	9.6	8.9	9.4	8.9	9.3	9.1
	None	57.2	57.0	56.8	56.9	56.9	53.9	53.6
	1-9 Days	18.5	16.9	16.2	14.8	14.4	14.0	13.5
	10-29 Days	7.5	7.4	8.3	8.7	9.3	10.7	11.0
	30-59 Days	3.4	4.0	4.3	4.7	5.2	6.0	6.1
Volunteer inside a home	60-89 Days	1.5	1.9	2.3	2.3	2.3	2.5	2.8
	90-129 Days	0.9	1.3	1.2	1.3	1.2	1.3	1.5
	130-180 Days	1.6	1.9	2.0	1.9	1.9	2.4	2.3
	No Answer	9.4	9.6	9.1	9.4	9.0	9.3	9.1
	None	52.5	52.2	53.9	55.1	55.8	55.5	56.9
	1-9 Days	18.7	17.3	16.1	13.7	12.7	11.1	10.6
	10-29 Days	8.2	7.9	7.8	7.7	7.4	7.8	7.5

Note: The number of school days in a year is estimated as 180 days.



**Number of Days per School Year Spent in Various Afterschool Activities
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
School sports team	No Answer	5.0	4.7	4.4	4.6	4.7	4.6	4.1
	None	40.0	37.2	41.5	43.2	48.5	53.5	57.6
	1-9 Days	18.8	13.2	10.8	7.1	5.0	3.9	3.3
	10-29 Days	10.2	7.8	5.9	3.3	2.4	1.7	1.9
	30-59 Days	9.6	11.7	10.1	7.4	6.5	5.8	5.3
	60-89 Days	7.1	9.9	9.7	10.2	10.3	9.3	9.1
	90-129 Days	5.5	8.9	9.9	12.8	12.3	11.8	10.1
	130-180 Days	3.8	6.6	7.6	11.4	10.4	9.3	8.6
School arts programs	No Answer	5.7	5.3	4.7	5.2	5.2	5.0	4.9
	None	58.2	54.3	52.9	53.2	54.7	56.4	58.0
	1-9 Days	21.5	22.2	21.9	19.0	15.9	13.3	12.1
	10-29 Days	6.0	7.6	8.2	7.2	7.4	6.2	6.4
	30-59 Days	3.4	4.0	4.7	5.1	5.1	5.5	5.2
	60-89 Days	1.8	2.5	3.0	4.0	4.5	4.7	4.8
	90-129 Days	1.4	1.6	1.6	2.9	3.6	4.2	4.4
	130-180 Days	2.0	2.6	2.9	3.4	3.7	4.7	4.1
School club/ intramurals	No Answer	6.8	6.2	5.4	5.6	5.6	5.5	5.0
	None	60.6	60.3	59.4	56.2	52.2	49.1	48.3
	1-9 Days	17.5	16.1	16.7	18.1	18.2	16.9	17.4
	10-29 Days	7.1	7.9	8.0	9.0	10.4	11.9	12.6
	30-59 Days	3.6	4.3	4.6	5.0	6.1	7.6	7.5
	60-89 Days	1.7	2.5	2.6	2.7	3.7	4.3	4.8
	90-129 Days	1.2	1.5	1.8	2.0	2.1	2.6	2.8
	130-180 Days	1.5	1.3	1.4	1.4	1.8	2.0	1.5
Youth centers (YMCA, etc)	No Answer	7.1	6.2	5.4	5.4	5.5	5.6	5.1
	None	70.9	73.2	74.5	77.3	78.1	79.2	80.6
	1-9 Days	11.8	10.4	10.7	9.4	8.1	7.1	6.9
	10-29 Days	4.3	4.2	3.8	3.1	3.1	3.3	3.1
	30-59 Days	2.2	2.6	2.2	2.1	2.3	1.9	2.2
	60-89 Days	1.2	1.3	1.5	1.2	1.5	1.4	1.0
	90-129 Days	1.0	1.0	0.9	0.7	0.8	0.7	0.5
	130-180 Days	1.4	1.1	1.0	0.7	0.7	0.7	0.6
Extended day programs	No Answer	8.3	7.0	6.3	6.2	6.1	6.2	5.7
	None	67.5	71.4	72.7	78.0	80.2	81.3	83.0
	1-9 Days	16.9	15.2	14.4	11.0	9.6	8.1	7.0
	10-29 Days	3.5	3.0	3.5	2.5	1.9	2.0	1.7
	30-59 Days	1.5	1.4	1.4	1.0	0.9	1.2	0.9
	60-89 Days	0.9	0.7	0.6	0.6	0.6	0.6	0.4
	90-129 Days	0.6	0.5	0.4	0.4	0.4	0.3	0.4
	130-180 Days	0.8	0.7	0.6	0.4	0.3	0.4	0.8

Note: The number of school days in a year is estimated as 180 days.

Female

**Number of Days per School Year Spent in Various Afterschool Activities
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Supervised youth activities (4H, scouts)	No Answer	9.2	8.5	7.8	7.6	7.4	7.5	6.8
	None	58.8	61.2	65.0	68.1	70.0	71.2	73.7
	1-9 Days	13.4	10.6	9.4	8.5	8.1	7.1	7.2
	10-29 Days	7.0	6.8	6.0	5.6	5.3	5.2	4.8
	30-59 Days	4.9	5.4	4.7	4.1	4.0	3.8	3.2
	60-89 Days	2.7	3.1	3.2	3.1	2.5	2.6	2.1
	90-129 Days	2.0	2.3	2.1	1.5	1.3	1.1	1.4
Afterschool classes (art, sport, etc)	130-180 Days	2.0	2.1	1.9	1.4	1.4	1.4	1.0
	No Answer	7.8	6.6	6.4	6.0	6.1	6.1	5.3
	None	38.0	38.9	41.7	45.0	48.8	52.6	57.8
	1-9 Days	21.1	18.9	17.3	14.8	13.5	11.7	11.0
	10-29 Days	9.1	9.0	8.4	7.9	7.6	7.2	6.3
	30-59 Days	7.5	7.7	7.5	7.3	6.7	6.2	5.6
	60-89 Days	5.4	6.2	6.1	6.1	5.5	5.0	4.5
Socialization without adult supervision	90-129 Days	4.5	5.1	5.3	5.2	4.9	4.4	3.7
	130-180 Days	6.5	7.6	7.3	7.7	6.8	6.8	5.8
	No Answer	6.3	5.8	5.0	4.7	4.9	5.0	4.4
	None	35.4	21.7	13.5	9.7	7.5	6.1	5.6
	1-9 Days	24.7	23.2	19.3	15.7	12.4	9.2	9.0
	10-29 Days	12.6	14.6	15.8	14.9	12.9	11.4	12.1
	30-59 Days	7.2	10.5	12.7	14.7	14.4	14.9	15.7
At home without adult supervision	60-89 Days	4.6	8.4	10.6	13.8	15.0	16.4	16.6
	90-129 Days	3.6	6.9	10.2	12.7	16.4	17.8	17.9
	130-180 Days	5.7	9.0	12.8	13.6	16.6	19.2	18.7
	No Answer	7.7	6.7	6.2	5.7	5.3	5.6	5.1
	None	27.3	16.9	12.4	8.7	8.1	7.6	7.0
	1-9 Days	31.0	26.0	21.8	17.9	15.0	13.2	12.7
	10-29 Days	13.5	16.2	17.1	16.8	15.9	15.6	15.5
At home with adult supervision	30-59 Days	7.7	11.8	13.3	16.2	16.3	16.1	17.1
	60-89 Days	4.5	7.7	10.4	12.6	14.4	15.0	14.1
	90-129 Days	3.6	6.5	8.4	10.9	12.5	13.5	13.7
	130-180 Days	4.7	8.1	10.5	11.2	12.4	13.5	14.8
	No Answer	8.7	7.6	6.9	6.2	6.1	6.1	5.6
	None	10.1	7.6	7.6	6.4	6.5	7.1	8.3
	1-9 Days	14.1	12.7	12.8	10.8	10.2	9.7	10.2

Note: The number of school days in a year is estimated as 180 days.

Female



**Number of Days per School Year Spent in Various Afterschool Activities
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Work for pay outside home	No Answer	7.8	6.7	5.9	5.8	5.7	5.4	5.0
	None	58.3	52.8	50.9	51.2	42.9	30.0	24.6
	1-9 Days	18.4	19.0	17.9	14.2	10.0	6.6	5.4
	10-29 Days	7.4	10.3	11.4	10.2	8.9	6.5	6.4
	30-59 Days	3.7	5.6	7.1	8.1	9.5	10.4	9.8
	60-89 Days	1.6	2.7	3.4	5.1	8.8	13.0	14.0
	90-129 Days	1.1	1.3	1.9	3.2	7.9	15.1	16.6
Work for pay inside a home	130-180 Days	1.7	1.6	1.6	2.4	6.3	13.0	18.2
	No Answer	7.4	6.7	5.9	5.9	5.8	5.6	5.2
	None	38.6	35.3	35.8	39.8	44.7	49.7	54.2
	1-9 Days	24.7	22.3	21.3	19.0	17.1	16.1	15.4
	10-29 Days	12.5	14.5	14.6	13.4	12.2	11.1	9.8
	30-59 Days	6.4	9.0	9.2	9.0	8.5	7.3	6.4
	60-89 Days	3.5	4.7	4.9	5.2	4.4	3.8	3.5
Volunteer outside the home	90-129 Days	2.4	3.2	3.5	3.1	3.2	2.8	2.4
	130-180 Days	4.5	4.3	4.9	4.7	4.2	3.6	3.0
	No Answer	7.7	6.8	5.8	5.9	5.7	5.6	5.1
	None	53.7	51.6	51.6	52.0	50.5	46.6	43.2
	1-9 Days	21.6	21.2	21.1	19.3	18.5	18.4	18.9
	10-29 Days	8.0	9.7	9.9	10.8	11.9	13.5	15.2
	30-59 Days	4.0	5.1	5.6	5.9	6.9	7.8	9.2
Volunteer inside a home	60-89 Days	2.0	2.5	2.7	3.0	3.2	4.3	4.1
	90-129 Days	1.4	1.4	1.6	1.8	1.8	2.4	2.3
	130-180 Days	1.5	1.6	1.7	1.3	1.4	1.5	2.0
	No Answer	7.2	6.5	5.7	5.6	5.6	5.4	4.8
	None	39.1	37.2	39.0	41.7	42.9	44.8	46.0
	1-9 Days	25.2	23.3	21.6	18.9	17.3	16.6	16.4
	10-29 Days	11.2	12.8	12.6	11.3	11.0	10.4	10.3

Note: The number of school days in a year is estimated as 180 days.

Female



**Participation in Some Form of Camp
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No participation in camps	71.6	74.1	77.7	81.2	81.1	81.2	81.7
Afternoons R.O.C.K. in Indiana	6.3	5.6	3.9	1.9	1.1	0.7	0.6
Youth Leadership Programs	10.0	12.3	11.7	9.4	9.4	9.5	8.2
S.A.D.D., S.T.A.N.D. or other prevention programs	10.5	6.6	5.4	6.1	6.6	6.5	7.2
Participated in 2+ types of camps	1.5	1.4	1.3	1.3	1.8	2.1	2.3

General

**Participation in Some Form of Camp
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No participation in camps	74.0	77.3	81.4	85.6	86.0	85.9	86.4
Afternoons R.O.C.K. in Indiana	6.2	5.5	3.9	2.1	1.1	0.9	0.7
Youth Leadership Programs	8.9	10.4	9.4	7.6	8.0	8.2	7.1
S.A.D.D., S.T.A.N.D. or other prevention programs	9.8	5.8	4.4	3.9	3.8	3.8	4.3
Participated in 2+ types of camps	1.2	1.0	0.9	0.8	1.1	1.2	1.4

Male

**Participation in Some Form of Camp
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No participation in camps	69.2	70.8	74.0	77.2	76.7	77.0	77.0
Afternoons R.O.C.K. in Indiana	6.4	5.6	4.0	1.7	1.0	0.6	0.4
Youth Leadership Programs	11.2	14.3	13.9	11.1	10.6	10.6	9.3
S.A.D.D., S.T.A.N.D. or other prevention programs	11.3	7.4	6.4	8.1	9.1	8.9	10.1
Participated in 2+ types of camps	1.9	1.8	1.6	1.9	2.5	2.9	3.2

Female



**Participation in Organized Family Events
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No Answer	4.1	3.2	3.0	2.7	2.5	2.5	2.4
Never	14.4	15.7	19.0	19.4	19.9	20.8	22.0
Once per Week	38.1	41.5	43.6	46.1	47.1	49.6	49.3
Twice per Week	20.3	20.0	18.3	18.0	16.9	15.5	15.1
3+ Times per Week	23.1	19.7	16.1	13.9	13.6	11.5	11.2

General

**Participation in Organized Family Events
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No Answer	4.8	3.7	3.5	3.1	3.1	3.0	3.1
Never	14.6	15.7	19.8	20.9	22.0	23.2	25.5
Once per Week	37.8	42.1	43.6	45.8	46.5	49.3	48.8
Twice per Week	20.4	19.3	18.2	17.1	15.9	14.5	13.3
3+ Times per Week	22.4	19.3	14.9	13.1	12.4	10.1	9.3

Male

**Participation in Organized Family Events
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No Answer	3.4	2.6	2.4	2.2	1.9	2.1	1.6
Never	14.1	15.6	18.2	17.9	18.0	18.7	18.6
Once per Week	38.4	40.9	43.7	46.5	47.7	50.0	49.9
Twice per Week	20.2	20.7	18.5	18.8	17.8	16.4	17.0
3+ Times per Week	23.8	20.1	17.2	14.6	14.7	12.9	13.0

Female

ADDERALL®
By: Christian H. Deneck, MPH and Jennifer E. Kelle, MPH

THE DRUG
Adderall®, a combination of Dex® Amphetamine, is a Schedule II stimulant used in the treatment of Attention Deficit Hyperactivity Disorder (ADHD). ADHD is defined as narcolepsy.

At least 10 percent of Indiana teenagers abuse Adderall® compared to 3.8 percent of teenagers nationwide. Between 1998 and 2005 we saw a steady increase in annual use among 12th graders reported in annual use of 4.7 percent in 1998, 5.3 percent in 10th grade and 5.3 percent in 12th grade (3.8 percent in 2005).

ACQUIRED
Drugs such as Adderall® are legally prescribed for children and adults with ADHD/ADD and are easily available for non-medical use. The cost for illicit use of the drug can range between 2 to 20 dollars for a 20 milligram tablet. Adolescents may acquire the drug from peers, friends or family members who have legal prescriptions. In some cases they may steal from

Indiana Annual Use Among Indiana Adolescents: 1998-2005

THE LAW
While the use of Adderall® itself is not illegal when taken as prescribed, use without a prescription is a violation of federal law.

New Factlines:

Print or order information on drugs from our Factline series. Each Factline includes information about the drug as well as Indiana specific data.

- Adderall
- Alcohol
- Benzodiazapines
- Cocaine
- Ecstasy
- Heroin
- Inhalants
- Ketamine
- Methamphetamine
- Non-Alcoholic Beer
- OxyContin
- Ritalin
- Steroids
- Tobacco

Corresponding training material available at www.drugs.indiana.edu



**Annual Prevalence of ATOD Use-Related Incidents
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had a hangover	No Answer	2.9	3.0	2.4	2.9	2.9	3.0	3.1
	Never	92.0	86.8	78.9	71.1	65.7	61.5	57.3
	Once	2.7	4.5	7.6	9.3	10.8	11.5	12.1
	Twice	1.1	2.3	4.2	5.4	6.8	7.7	7.8
	3-5 times	0.7	1.5	3.3	5.5	6.4	7.8	9.2
	6-9 times	0.2	0.6	1.2	1.8	2.4	2.9	3.4
	10+ times	0.4	1.3	2.4	3.9	5.0	5.6	7.2
Poor on school test	No Answer	3.1	3.2	2.8	3.3	3.4	3.3	3.5
	Never	89.0	86.8	84.2	83.0	83.0	84.5	84.3
	Once	3.0	3.2	3.9	3.8	3.6	3.3	3.5
	Twice	1.7	2.0	2.4	2.2	2.4	2.1	2.2
	3-5 times	1.5	2.1	2.7	2.9	2.9	2.9	2.5
	6-9 times	0.5	0.7	1.2	1.1	1.4	1.2	1.2
	10+ times	1.1	1.9	2.8	3.7	3.3	2.7	2.8
Had trouble with police	No Answer	3.1	3.1	2.8	3.3	3.4	3.5	3.5
	Never	92.5	91.1	88.7	87.8	87.9	88.3	87.4
	Once	2.3	2.5	3.8	4.1	4.3	4.7	5.6
	Twice	0.8	1.1	1.7	1.8	1.7	1.4	1.5
	3-5 times	0.8	1.1	1.4	1.6	1.2	1.1	1.1
	6-9 times	0.2	0.3	0.6	0.5	0.5	0.3	0.2
	10+ times	0.4	0.7	1.0	0.9	0.9	0.6	0.6
Had trouble with parents	No Answer	3.3	3.3	2.9	3.4	3.5	3.4	3.4
	Never	86.5	84.5	80.3	77.6	76.3	76.2	77.0
	Once	3.4	4.1	5.9	7.2	8.4	9.0	9.5
	Twice	1.5	1.8	2.6	2.9	3.6	4.2	3.5
	3-5 times	1.4	1.7	2.4	2.9	2.9	3.1	3.0
	6-9 times	0.6	0.7	1.1	1.2	1.1	1.0	1.0
	10+ times	3.3	3.8	4.8	4.8	4.1	3.0	2.6
Damaged property	No Answer	3.2	3.4	3.0	3.3	3.4	3.5	3.6
	Never	93.9	91.9	89.9	89.0	89.1	90.6	90.6
	Once	1.3	2.0	2.6	2.6	2.5	1.9	1.8
	Twice	0.6	0.8	1.4	1.6	1.5	1.2	1.1
	3-5 times	0.4	0.7	1.4	1.4	1.4	1.3	1.1
	6-9 times	0.1	0.3	0.6	0.7	0.6	0.5	0.6
	10+ times	0.4	0.8	1.2	1.4	1.3	1.1	1.3
Got into a fight	No Answer	3.2	3.4	3.0	3.5	3.6	3.6	3.9
	Never	86.0	83.6	78.8	76.7	75.6	76.1	76.2
	Once	3.4	4.3	5.7	6.4	7.1	7.2	6.9
	Twice	1.9	2.2	3.2	3.6	3.9	4.4	4.4
	3-5 times	2.0	2.2	3.4	3.8	4.0	4.1	4.3
	6-9 times	0.9	1.0	1.4	1.6	1.8	1.5	1.6
	10+ times	2.5	3.2	4.4	4.4	4.0	3.1	2.6

General

**Annual Prevalence of ATOD Use-Related Incidents
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had nausea or vomited	No Answer	3.5	3.5	3.0	3.4	3.5	3.5	3.8
	Never	90.2	86.7	81.7	76.0	71.0	67.6	63.1
	Once	2.6	4.5	6.7	8.7	11.1	12.0	13.1
	Twice	1.2	1.8	2.9	4.2	5.0	5.7	6.9
	3-5 times	1.2	1.6	2.3	3.7	4.7	6.0	7.1
	6-9 times	0.4	0.7	1.2	1.5	1.7	2.3	2.6
	10+ times	0.9	1.3	2.1	2.6	3.1	2.9	3.3
Driven under the influence	No Answer	3.6	3.5	3.3	3.8	3.6	3.8	3.8
	Never	94.4	93.7	92.3	90.7	86.5	80.5	76.0
	Once	1.0	1.2	1.7	2.0	3.9	5.9	6.1
	Twice	0.4	0.4	0.8	1.0	1.7	2.4	3.2
	3-5 times	0.2	0.4	0.7	0.9	1.8	2.7	3.9
	6-9 times	0.1	0.2	0.3	0.4	0.7	1.3	1.6
	10+ times	0.4	0.6	1.0	1.2	1.8	3.2	5.3
Rode with a drunk driver	No Answer	3.6	3.6	3.1	3.5	3.4	3.5	3.5
	Never	84.2	82.1	77.7	75.5	73.0	72.6	70.4
	Once	5.0	4.9	6.1	6.5	7.6	7.8	8.2
	Twice	2.1	2.8	3.5	4.0	4.4	4.5	4.7
	3-5 times	2.0	2.5	3.8	4.0	4.7	5.1	5.2
	6-9 times	0.5	0.8	1.5	1.6	2.1	1.9	2.2
	10+ times	2.5	3.3	4.4	4.9	4.9	4.5	5.8
Missed school	No Answer	3.5	3.5	3.1	3.5	3.4	3.5	3.5
	Never	88.5	88.0	86.2	85.7	85.1	85.4	85.2
	Once	1.8	2.3	2.8	2.8	3.2	2.9	3.0
	Twice	1.4	1.3	1.8	1.9	1.8	2.0	2.0
	3-5 times	2.2	2.0	2.5	2.4	2.6	2.5	2.3
	6-9 times	1.0	1.0	1.3	1.2	1.3	1.1	1.4
	10+ times	1.5	1.8	2.3	2.4	2.6	2.6	2.8
Had a memory loss	No Answer	3.4	3.4	3.1	3.4	3.4	3.4	3.5
	Never	92.2	89.4	85.9	81.2	77.9	76.2	74.8
	Once	2.1	3.4	4.4	5.5	6.2	6.9	6.7
	Twice	0.8	1.3	2.1	3.0	3.6	4.1	4.2
	3-5 times	0.6	1.1	1.9	2.8	3.7	4.0	4.6
	6-9 times	0.3	0.5	0.8	1.0	1.6	1.6	1.8
	10+ times	0.6	1.1	1.9	3.1	3.7	3.8	4.4



**Annual Prevalence of ATOD Use-Related Incidents
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had a hangover	No Answer	3.7	3.9	3.1	3.7	4.0	4.2	3.9
	Never	90.6	86.2	79.2	71.5	66.2	62.0	56.3
	Once	3.0	4.2	6.7	8.2	9.7	10.4	10.6
	Twice	1.2	2.3	3.7	4.9	5.8	6.8	6.9
	3-5 times	0.8	1.3	3.4	5.0	5.8	6.9	9.0
	6-9 times	0.1	0.6	1.2	1.6	2.5	2.7	3.6
	10+ times	0.5	1.5	2.8	5.0	6.0	7.1	9.6
Poor on school test	No Answer	3.9	4.0	3.5	4.3	4.4	4.6	4.4
	Never	87.5	85.6	83.8	82.3	82.0	82.5	81.8
	Once	3.2	3.2	3.8	3.4	3.2	3.2	3.5
	Twice	1.9	1.8	2.2	2.0	2.3	2.0	2.5
	3-5 times	1.6	2.3	2.6	2.6	2.8	3.1	2.9
	6-9 times	0.5	0.7	1.0	1.1	1.4	1.2	1.3
	10+ times	1.4	2.3	3.1	4.2	3.9	3.5	3.6
Had trouble with police	No Answer	3.9	3.9	3.5	4.2	4.6	4.8	4.3
	Never	90.1	89.1	86.8	85.3	85.4	85.8	84.2
	Once	2.9	2.9	4.0	4.3	4.5	4.9	6.4
	Twice	1.1	1.4	2.0	2.2	2.1	1.7	2.1
	3-5 times	1.1	1.3	1.7	1.9	1.6	1.5	1.7
	6-9 times	0.3	0.5	0.7	0.8	0.6	0.3	0.3
	10+ times	0.6	1.0	1.3	1.3	1.3	1.0	0.9
Had trouble with parents	No Answer	4.2	4.1	3.6	4.5	4.6	4.6	4.4
	Never	84.4	83.0	80.1	77.4	75.2	75.2	74.7
	Once	3.4	4.1	5.4	6.3	8.1	8.5	9.1
	Twice	1.6	1.7	2.5	2.8	3.8	4.0	3.9
	3-5 times	1.5	1.8	2.3	2.7	2.8	3.2	3.6
	6-9 times	0.7	0.7	1.0	1.0	1.1	1.0	1.3
	10+ times	4.1	4.5	5.1	5.3	4.4	3.5	3.1
Damaged property	No Answer	4.1	4.4	3.7	4.3	4.5	4.8	4.4
	Never	92.1	89.8	88.1	86.0	85.5	86.7	86.7
	Once	1.7	2.4	2.8	2.9	3.1	2.3	2.5
	Twice	0.8	1.1	1.6	1.9	1.9	1.7	1.8
	3-5 times	0.5	0.9	1.4	1.8	1.8	2.0	1.7
	6-9 times	0.2	0.4	0.8	1.1	1.0	0.7	0.9
	10+ times	0.6	1.0	1.7	2.0	2.1	1.8	2.0
Got into a fight	No Answer	4.0	4.2	3.7	4.6	4.6	4.9	4.7
	Never	83.8	82.4	78.5	76.5	75.0	74.8	73.5
	Once	3.7	4.1	5.3	5.7	6.4	6.6	7.2
	Twice	2.1	2.3	3.1	3.0	3.7	4.3	4.8
	3-5 times	2.2	2.2	3.3	3.9	3.8	4.1	4.6
	6-9 times	1.1	0.9	1.5	1.4	1.8	1.7	1.8
	10+ times	3.2	3.8	4.7	4.9	4.7	3.7	3.4

Male

**Annual Prevalence of ATOD Use-Related Incidents
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had nausea or vomited	No Answer	4.3	4.4	3.7	4.5	4.5	4.7	4.6
	Never	88.7	86.0	82.6	77.9	73.0	68.4	62.9
	Once	2.9	4.4	5.9	7.3	9.3	10.8	11.9
	Twice	1.3	1.6	2.5	3.3	4.3	5.2	6.8
	3-5 times	1.3	1.5	2.0	3.2	4.2	5.3	7.0
	6-9 times	0.4	0.6	1.1	1.2	1.5	2.2	2.6
	10+ times	1.2	1.4	2.1	2.7	3.2	3.5	4.2
Driven under the influence	No Answer	4.5	4.3	4.0	4.7	4.6	5.2	4.7
	Never	93.2	92.7	91.3	89.0	83.8	78.1	72.0
	Once	1.0	1.2	1.8	2.3	4.0	5.2	5.5
	Twice	0.5	0.4	0.8	1.0	1.7	2.4	3.4
	3-5 times	0.2	0.4	0.6	1.0	2.4	3.0	4.3
	6-9 times	0.1	0.1	0.3	0.5	0.7	1.4	2.1
	10+ times	0.5	0.9	1.2	1.5	2.7	4.6	8.1
Rode with a drunk driver	No Answer	4.3	4.4	3.7	4.5	4.5	4.8	4.5
	Never	83.5	82.2	78.8	77.5	74.2	73.7	69.4
	Once	4.7	4.4	5.6	4.9	6.2	5.7	6.9
	Twice	2.3	2.5	3.0	3.3	3.7	3.9	4.3
	3-5 times	2.1	2.2	3.4	3.3	4.0	4.6	4.9
	6-9 times	0.5	0.8	1.1	1.3	1.9	1.9	2.6
	10+ times	2.8	3.4	4.4	5.2	5.5	5.3	7.6
Missed school	No Answer	4.2	4.3	3.7	4.5	4.5	4.8	4.4
	Never	86.8	86.8	85.7	84.6	83.7	83.6	82.3
	Once	2.1	2.1	2.8	2.7	3.2	2.8	3.2
	Twice	1.4	1.5	1.6	2.1	1.9	2.1	2.2
	3-5 times	2.6	2.2	2.5	2.2	2.8	2.8	2.9
	6-9 times	1.0	1.1	1.2	1.2	1.1	0.9	1.7
	10+ times	2.0	2.0	2.5	2.8	2.8	2.9	3.3
Had a memory loss	No Answer	4.2	4.3	3.8	4.5	4.5	4.8	4.5
	Never	90.8	89.0	86.0	81.7	78.6	75.7	72.8
	Once	2.3	3.0	3.8	4.5	5.3	6.0	6.2
	Twice	1.0	1.2	1.9	2.5	2.9	3.6	3.9
	3-5 times	0.7	1.0	1.6	2.3	3.0	4.0	5.2
	6-9 times	0.3	0.4	0.7	0.8	1.4	1.6	1.9
	10+ times	0.8	1.1	2.1	3.6	4.3	4.3	5.4

Male



**Annual Prevalence of ATOD Use-Related Incidents
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had a hangover	No Answer	2.2	2.1	1.7	2.1	2.0	1.8	2.1
	Never	93.3	87.5	78.6	70.8	65.3	61.2	58.4
	Once	2.4	4.8	8.4	10.3	11.8	12.5	13.5
	Twice	0.9	2.1	4.6	5.9	7.8	8.5	8.7
	3-5 times	0.6	1.7	3.3	6.0	6.9	8.5	9.3
	6-9 times	0.2	0.5	1.2	2.0	2.3	3.1	3.1
	10+ times	0.3	1.2	2.1	2.9	4.0	4.4	4.8
Poor on school test	No Answer	2.4	2.4	2.1	2.3	2.4	2.1	2.3
	Never	90.5	88.0	84.6	83.6	84.0	86.3	87.1
	Once	2.8	3.2	4.1	4.2	3.8	3.3	3.5
	Twice	1.6	2.3	2.6	2.4	2.5	2.3	1.8
	3-5 times	1.4	2.0	2.8	3.2	3.1	2.7	2.2
	6-9 times	0.4	0.7	1.4	1.1	1.5	1.3	1.2
	10+ times	0.8	1.5	2.4	3.2	2.8	2.0	1.9
Had trouble with police	No Answer	2.3	2.4	2.1	2.5	2.3	2.4	2.4
	Never	94.8	93.3	90.6	90.0	90.2	90.6	90.8
	Once	1.7	2.0	3.6	3.9	4.2	4.6	4.8
	Twice	0.6	0.8	1.4	1.5	1.5	1.1	0.9
	3-5 times	0.4	0.8	1.1	1.2	0.9	0.7	0.6
	6-9 times	0.1	0.2	0.5	0.3	0.4	0.3	0.0
	10+ times	0.1	0.4	0.7	0.6	0.5	0.3	0.3
Had trouble with parents	No Answer	2.5	2.5	2.3	2.4	2.5	2.4	2.3
	Never	88.5	86.1	80.5	77.9	77.5	77.2	79.4
	Once	3.4	4.0	6.4	7.9	8.7	9.4	9.9
	Twice	1.3	1.9	2.7	2.9	3.4	4.3	3.1
	3-5 times	1.3	1.6	2.5	3.1	3.0	3.1	2.5
	6-9 times	0.6	0.7	1.2	1.4	1.1	1.0	0.8
	10+ times	2.4	3.1	4.5	4.4	3.8	2.6	2.0
Damaged property	No Answer	2.4	2.5	2.2	2.3	2.4	2.3	2.5
	Never	95.8	94.1	91.8	91.8	92.4	94.1	94.6
	Once	0.9	1.5	2.5	2.4	2.0	1.5	1.1
	Twice	0.3	0.6	1.2	1.3	1.2	0.7	0.5
	3-5 times	0.3	0.6	1.3	1.0	1.0	0.7	0.6
	6-9 times	0.1	0.3	0.4	0.3	0.3	0.2	0.2
	10+ times	0.3	0.5	0.7	0.9	0.7	0.5	0.4
Got into a fight	No Answer	2.4	2.6	2.2	2.4	2.6	2.4	2.8
	Never	88.2	84.9	79.1	76.9	76.2	77.3	79.0
	Once	3.2	4.5	6.1	7.1	7.8	7.7	6.6
	Twice	1.7	2.2	3.4	4.2	4.1	4.5	4.1
	3-5 times	1.8	2.1	3.6	3.7	4.1	4.1	4.1
	6-9 times	0.8	1.1	1.4	1.9	1.8	1.4	1.5
	10+ times	1.9	2.6	4.2	3.9	3.5	2.6	1.9

Female

**Annual Prevalence of ATOD Use-Related Incidents
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had nausea or vomited	No Answer	2.7	2.7	2.3	2.4	2.5	2.4	2.8
	Never	91.6	87.4	80.9	74.3	69.3	66.9	63.6
	Once	2.4	4.5	7.6	10.0	12.7	13.1	14.3
	Twice	1.2	2.0	3.3	5.0	5.7	6.2	7.0
	3-5 times	1.0	1.6	2.7	4.1	5.1	6.6	7.2
	6-9 times	0.4	0.7	1.3	1.7	1.9	2.5	2.6
	10+ times	0.7	1.1	2.0	2.4	2.9	2.3	2.5
Driven under the influence	No Answer	2.7	2.7	2.5	2.9	2.7	2.6	2.7
	Never	95.5	94.9	93.3	92.3	88.9	82.7	80.1
	Once	0.9	1.1	1.6	1.8	3.8	6.6	6.8
	Twice	0.3	0.3	0.8	0.9	1.6	2.4	3.1
	3-5 times	0.2	0.4	0.7	0.9	1.3	2.5	3.5
	6-9 times	0.1	0.2	0.3	0.3	0.7	1.2	1.1
	10+ times	0.2	0.4	0.7	0.8	1.0	2.0	2.6
Rode with a drunk driver	No Answer	2.9	2.8	2.4	2.6	2.3	2.3	2.3
	Never	85.1	81.9	76.6	73.6	72.1	71.8	71.7
	Once	5.3	5.4	6.7	8.0	8.8	9.6	9.4
	Twice	2.0	3.1	4.0	4.6	5.0	5.0	5.1
	3-5 times	2.0	2.7	4.3	4.7	5.2	5.5	5.5
	6-9 times	0.5	0.8	1.8	1.9	2.2	1.9	1.9
	10+ times	2.1	3.3	4.3	4.6	4.3	3.9	4.1
Missed school	No Answer	2.8	2.8	2.4	2.6	2.4	2.4	2.3
	Never	90.3	89.3	86.7	86.8	86.4	87.1	88.3
	Once	1.5	2.4	2.9	3.0	3.1	2.9	2.6
	Twice	1.4	1.1	1.9	1.8	1.8	1.8	1.7
	3-5 times	1.9	1.8	2.6	2.5	2.3	2.3	1.7
	6-9 times	1.0	1.0	1.4	1.2	1.4	1.2	1.2
	10+ times	1.1	1.6	2.0	2.1	2.5	2.3	2.2
Had a memory loss	No Answer	2.5	2.6	2.2	2.4	2.3	2.1	2.3
	Never	93.5	89.8	85.8	80.8	77.4	76.7	77.0
	Once	1.9	3.7	4.9	6.4	7.0	7.7	7.2
	Twice	0.7	1.3	2.3	3.4	4.1	4.7	4.5
	3-5 times	0.6	1.1	2.3	3.4	4.2	4.0	4.1
	6-9 times	0.2	0.5	0.8	1.2	1.8	1.5	1.7
	10+ times	0.5	1.0	1.7	2.6	3.3	3.3	3.3

Female



**Perceived Safety in Neighborhood and in School
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Out alone during the day	No Answer	3.3	3.4	3.1	3.4	3.6	3.7	4.1
	Very Safe	54.5	62.5	66.4	71.6	74.2	77.8	79.5
	Somewhat Safe	31.1	27.0	24.4	20.5	18.1	15.1	13.2
	Somewhat Unsafe	7.5	5.0	3.9	2.8	2.5	1.8	1.6
	Very Unsafe	3.5	2.1	2.1	1.7	1.7	1.5	1.5
Out alone after dark	No Answer	3.4	3.6	3.2	3.5	3.7	3.9	4.2
	Very Safe	20.9	27.3	33.2	39.5	43.0	49.1	51.0
	Somewhat Safe	29.9	33.8	33.6	32.5	31.0	29.0	28.9
	Somewhat Unsafe	23.2	21.1	18.4	15.6	14.8	12.2	10.9
	Very Unsafe	22.5	14.3	11.6	8.8	7.6	5.8	5.0
Inside home alone during the day	No Answer	3.5	3.7	3.4	3.7	4.0	3.9	4.4
	Very Safe	69.4	76.2	79.8	81.9	82.9	84.4	85.0
	Somewhat Safe	19.9	15.9	13.3	11.4	10.3	9.2	8.2
	Somewhat Unsafe	4.7	2.7	2.1	1.5	1.6	1.2	1.0
	Very Unsafe	2.4	1.5	1.5	1.5	1.3	1.3	1.4
Inside home alone after dark	No Answer	3.6	3.6	3.3	3.5	3.8	3.8	4.3
	Very Safe	41.8	50.5	57.6	62.0	63.4	67.2	68.5
	Somewhat Safe	27.1	27.0	24.4	22.9	21.5	19.5	19.1
	Somewhat Unsafe	15.8	12.0	9.5	7.5	7.7	6.2	5.3
	Very Unsafe	11.8	6.8	5.2	4.2	3.6	3.3	2.8
Traveling to/from school	No Answer	3.4	3.7	3.3	3.6	3.8	3.9	4.3
	Very Safe	64.1	68.9	70.1	73.4	74.0	77.5	77.8
	Somewhat Safe	22.3	20.5	20.0	18.1	17.5	15.1	14.2
	Somewhat Unsafe	6.2	4.2	4.1	2.8	2.7	1.8	1.9
	Very Unsafe	3.9	2.7	2.5	2.1	1.9	1.7	1.7
Staying in class	No Answer	3.4	3.6	3.4	3.6	3.8	4.0	4.3
	Very Safe	83.6	80.5	76.5	74.2	72.6	75.4	76.8
	Somewhat Safe	9.8	12.1	15.2	17.2	18.1	15.6	14.4
	Somewhat Unsafe	1.6	1.8	2.3	2.4	2.7	2.5	2.1
	Very Unsafe	1.5	2.0	2.6	2.6	2.7	2.5	2.4
Alone in school hallways	No Answer	3.5	3.7	3.4	3.8	4.0	4.1	4.4
	Very Safe	71.3	69.9	70.7	68.0	68.6	72.8	74.9
	Somewhat Safe	18.4	18.9	18.3	20.4	19.9	16.5	15.2
	Somewhat Unsafe	4.3	4.8	4.3	4.7	4.3	3.5	2.9
	Very Unsafe	2.5	2.7	3.3	3.1	3.3	3.1	2.6
Staying for afterschool activities	No Answer	4.3	4.1	3.8	4.0	4.2	4.3	4.6
	Very Safe	71.4	72.2	71.6	70.7	70.2	73.5	75.2
	Somewhat Safe	17.6	17.1	17.8	18.5	18.9	16.2	15.1
	Somewhat Unsafe	4.1	4.0	3.3	3.7	3.5	3.0	2.3
	Very Unsafe	2.6	2.6	3.4	3.1	3.1	3.0	2.7

General

**Perceived Safety in Neighborhood and in School
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Out alone during the day	No Answer	4.1	4.2	4.3	4.3	4.5	5.0	5.4
	Very Safe	60.2	68.0	72.4	78.1	80.5	83.0	83.3
	Somewhat Safe	27.4	22.5	18.4	14.0	11.8	9.3	8.3
	Somewhat Unsafe	5.3	3.6	2.7	1.8	1.4	1.0	1.1
	Very Unsafe	2.9	1.8	2.2	1.9	1.8	1.6	1.9
Out alone after dark	No Answer	4.2	4.3	4.4	4.4	4.7	5.1	5.4
	Very Safe	27.6	36.1	44.6	54.3	59.7	66.8	68.4
	Somewhat Safe	31.8	34.2	31.8	28.6	25.3	21.1	19.4
	Somewhat Unsafe	21.1	16.2	12.5	8.5	7.0	4.3	4.0
	Very Unsafe	15.3	9.2	6.7	4.2	3.3	2.7	2.7
Inside home alone during the day	No Answer	4.2	4.5	4.4	4.6	5.1	5.2	5.6
	Very Safe	74.6	80.0	83.1	85.3	86.6	87.2	87.1
	Somewhat Safe	15.7	12.3	9.6	7.5	5.9	5.6	4.9
	Somewhat Unsafe	3.3	1.8	1.3	0.9	1.0	0.6	0.7
	Very Unsafe	2.1	1.4	1.6	1.7	1.5	1.4	1.7
Inside home alone after dark	No Answer	4.3	4.3	4.4	4.4	4.8	5.1	5.6
	Very Safe	50.7	59.8	68.5	74.2	77.4	80.9	81.1
	Somewhat Safe	25.1	23.7	18.7	15.6	13.0	10.7	9.7
	Somewhat Unsafe	12.1	7.9	5.2	3.3	2.8	1.5	1.6
	Very Unsafe	7.9	4.2	3.3	2.4	2.1	1.8	2.0
Traveling to/from school	No Answer	4.2	4.6	4.4	4.7	4.9	5.1	5.7
	Very Safe	67.4	71.9	73.9	77.0	78.0	80.6	80.1
	Somewhat Safe	20.3	17.8	16.1	13.6	12.7	10.7	10.4
	Somewhat Unsafe	4.7	3.1	3.0	2.2	2.1	1.6	1.7
	Very Unsafe	3.4	2.6	2.7	2.5	2.3	1.9	2.1
Staying in class	No Answer	4.3	4.3	4.5	4.7	4.9	5.3	5.6
	Very Safe	81.9	79.2	75.0	73.4	72.7	75.7	76.9
	Somewhat Safe	10.1	11.9	14.5	15.7	16.2	12.9	12.2
	Somewhat Unsafe	1.7	1.9	2.4	2.5	2.5	2.6	2.0
	Very Unsafe	1.9	2.6	3.6	3.7	3.6	3.5	3.3
Alone in school hallways	No Answer	4.4	4.5	4.5	4.9	5.0	5.4	5.7
	Very Safe	71.7	69.9	71.2	69.1	70.7	74.6	76.2
	Somewhat Safe	17.2	17.7	16.1	17.8	16.6	13.2	12.0
	Somewhat Unsafe	3.8	4.6	4.1	4.3	3.6	2.8	2.5
	Very Unsafe	2.9	3.4	4.2	3.9	4.2	3.9	3.6
Staying for afterschool activities	No Answer	5.5	5.0	5.0	5.2	5.4	5.7	6.1
	Very Safe	71.3	72.5	72.1	71.7	71.9	75.5	76.6
	Somewhat Safe	16.7	16.0	15.8	16.1	15.7	12.5	11.5
	Somewhat Unsafe	3.4	3.4	2.8	3.1	2.8	2.5	2.0
	Very Unsafe	3.1	3.0	4.2	3.9	4.2	3.9	3.8

Male



**Perceived Safety in Neighborhood and in School
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Out alone during the day	No Answer	2.5	2.6	1.9	2.4	2.7	2.5	2.7
	Very Safe	48.9	56.9	60.5	65.6	68.6	73.3	76.0
	Somewhat Safe	34.9	31.5	30.5	26.7	23.8	20.4	18.0
	Somewhat Unsafe	9.7	6.5	5.2	3.7	3.4	2.5	2.2
	Very Unsafe	4.1	2.4	1.9	1.5	1.5	1.3	1.1
Out alone after dark	No Answer	2.7	2.8	2.0	2.6	2.7	2.6	2.8
	Very Safe	14.3	18.3	21.7	25.5	27.7	33.2	33.9
	Somewhat Safe	28.0	33.5	35.5	36.3	36.3	36.2	38.3
	Somewhat Unsafe	25.4	26.0	24.3	22.4	21.9	19.4	17.8
	Very Unsafe	29.6	19.4	16.5	13.1	11.4	8.6	7.2
Inside home alone during the day	No Answer	2.8	3.0	2.3	2.8	2.9	2.8	2.9
	Very Safe	64.3	72.3	76.6	78.8	79.7	81.9	83.3
	Somewhat Safe	24.1	19.6	17.0	15.0	14.2	12.5	11.5
	Somewhat Unsafe	6.2	3.5	2.9	2.1	2.1	1.7	1.3
	Very Unsafe	2.7	1.6	1.3	1.3	1.0	1.1	1.0
Inside home alone after dark	No Answer	2.8	2.9	2.1	2.6	2.9	2.6	2.8
	Very Safe	33.0	41.0	46.8	50.4	50.7	54.8	56.2
	Somewhat Safe	29.2	30.5	30.0	29.7	29.3	27.6	28.4
	Somewhat Unsafe	19.5	16.2	13.9	11.3	12.2	10.4	9.0
	Very Unsafe	15.6	9.5	7.1	5.9	5.0	4.6	3.6
Traveling to/from school	No Answer	2.6	2.8	2.2	2.5	2.7	2.6	2.8
	Very Safe	61.0	66.0	66.5	70.1	70.5	74.8	75.9
	Somewhat Safe	24.3	23.2	23.9	22.3	21.9	19.1	18.0
	Somewhat Unsafe	7.8	5.3	5.2	3.4	3.3	2.0	2.0
	Very Unsafe	4.3	2.7	2.3	1.7	1.5	1.5	1.3
Staying in class	No Answer	2.4	2.8	2.3	2.6	2.8	2.8	2.8
	Very Safe	85.4	81.9	78.0	74.9	72.7	75.3	77.0
	Somewhat Safe	9.5	12.4	15.9	18.5	19.9	17.9	16.7
	Somewhat Unsafe	1.5	1.7	2.1	2.3	2.7	2.4	2.2
	Very Unsafe	1.1	1.2	1.7	1.7	1.8	1.6	1.4
Alone in school hallways	No Answer	2.5	3.0	2.3	2.6	2.9	2.8	2.8
	Very Safe	70.8	69.9	70.3	67.1	66.7	71.3	74.0
	Somewhat Safe	19.7	20.1	20.6	22.8	23.0	19.5	18.5
	Somewhat Unsafe	4.8	5.1	4.5	5.1	5.0	4.0	3.2
	Very Unsafe	2.2	2.0	2.4	2.4	2.4	2.4	1.5
Staying for afterschool activities	No Answer	3.1	3.2	2.5	2.8	3.2	3.0	2.9
	Very Safe	71.4	71.9	71.2	69.9	68.7	71.8	74.2
	Somewhat Safe	18.5	18.3	19.9	20.7	21.8	19.6	18.7
	Somewhat Unsafe	4.8	4.5	3.8	4.2	4.3	3.4	2.6
	Very Unsafe	2.2	2.1	2.7	2.3	2.1	2.2	1.5

Female

**Annual Prevalence of Weapon Use and Violent Behavior
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Physical fight	No Answer	3.6	3.6	3.6	3.9	4.0	4.3	4.7
	Never	60.5	60.8	57.9	63.1	66.8	70.5	74.5
	1-2 times	23.5	23.2	25.3	22.8	20.2	18.3	15.3
	3-9 times	8.1	8.2	8.9	6.9	6.2	4.8	3.9
	10+ times	4.3	4.2	4.4	3.2	2.7	2.1	1.7
Shouting argument	No Answer	3.9	4.0	3.9	4.2	4.3	4.5	4.8
	Never	46.0	41.5	34.7	34.6	34.5	36.5	39.4
	1-2 times	26.4	26.4	26.5	26.5	26.2	26.8	26.5
	3-9 times	12.7	14.7	18.3	18.6	19.2	18.6	17.9
	10+ times	11.0	13.4	16.6	16.1	15.8	13.6	11.5
Carried gun for hunting	No Answer	3.8	3.8	3.8	4.2	4.3	4.5	4.7
	Never	72.6	70.6	69.3	71.3	72.5	74.7	74.7
	1-2 times	9.9	10.2	10.3	8.8	8.1	7.3	6.8
	3-9 times	6.1	6.6	7.0	6.5	6.1	5.7	5.5
	10+ times	7.6	8.7	9.5	9.2	9.0	7.8	8.3
Carried gun for other reasons	No Answer	3.8	3.9	3.9	4.1	4.3	4.5	4.8
	Never	91.5	90.3	88.3	88.6	88.5	89.2	87.9
	1-2 times	2.5	3.0	4.0	3.4	3.3	3.0	3.1
	3-9 times	0.9	1.2	1.7	1.6	1.5	1.3	1.5
	10+ times	1.2	1.5	2.1	2.3	2.4	2.0	2.7
Carried weapon for hunting	No Answer	3.9	3.9	4.0	4.2	4.5	4.6	4.8
	Never	82.0	80.5	79.1	80.4	80.7	82.1	82.2
	1-2 times	7.1	7.2	7.2	6.1	5.4	4.6	4.2
	3-9 times	3.1	3.7	4.0	3.7	3.6	3.4	3.2
	10+ times	3.8	4.7	5.7	5.7	5.8	5.3	5.5
Carried weapon for other reasons	No Answer	3.9	4.0	4.0	4.3	4.5	4.5	4.9
	Never	87.6	85.6	81.9	82.2	81.0	82.6	82.4
	1-2 times	4.2	5.2	6.5	5.8	5.8	4.8	4.6
	3-9 times	1.8	2.2	3.0	3.0	3.0	2.6	2.5
	10+ times	2.4	3.0	4.5	4.8	5.7	5.4	5.7
Carried gun to school	No Answer	3.7	3.9	3.9	4.2	4.4	4.5	4.8
	Never	95.6	94.9	94.7	94.3	93.8	93.6	93.1
	1-2 times	0.3	0.5	0.7	0.7	0.6	0.7	0.9
	3-9 times	0.1	0.2	0.2	0.2	0.3	0.4	0.3
	10+ times	0.3	0.4	0.5	0.6	0.8	0.8	0.9
Carried weapon to school	No Answer	3.8	4.0	3.9	4.2	4.3	4.5	4.9
	Never	93.4	92.0	90.2	89.4	88.6	88.9	88.6
	1-2 times	2.0	2.7	3.9	3.6	3.6	3.1	2.6
	3-9 times	0.3	0.6	0.9	1.0	1.3	1.2	1.1
	10+ times	0.3	0.6	1.1	1.8	2.2	2.3	2.8

General



**Annual Prevalence of Weapon Use and Violent Behavior
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Threatened by weapon on school property	No Answer	4.0	4.1	4.0	4.4	4.5	4.7	4.9
	Never	89.5	88.5	87.8	87.8	88.5	90.0	90.7
	1-2 times	5.1	5.6	6.1	5.6	4.8	3.7	2.9
	3-9 times	0.8	1.1	1.2	1.2	1.2	0.9	0.7
Property stolen while at school	10+ times	0.7	0.8	0.8	1.0	1.0	0.7	0.8
	No Answer	4.3	4.3	4.1	4.4	4.6	4.7	4.9
	Never	67.3	62.1	61.3	60.5	63.8	68.0	71.2
	1-2 times	20.7	24.7	25.1	26.0	24.4	21.3	19.2
Missed school: Felt unsafe on school property	3-9 times	5.5	6.3	7.0	6.9	5.5	4.7	3.7
	10+ times	2.2	2.6	2.5	2.2	1.7	1.3	1.1
	No Answer	4.0	4.2	4.1	4.4	4.5	4.6	4.8
	Never	92.0	91.0	90.8	90.6	90.3	90.9	91.0
Missed school: Felt unsafe on way to/from school	1-2 times	3.1	3.4	3.6	3.5	3.5	3.0	2.6
	3-9 times	0.6	0.7	0.8	0.7	0.9	0.7	0.7
	10+ times	0.4	0.7	0.7	0.8	0.8	0.7	0.9
	No Answer	4.0	4.2	4.1	4.4	4.5	4.7	4.9
	Never	93.6	93.3	93.2	93.0	92.7	93.1	92.6
	1-2 times	1.8	1.6	1.7	1.5	1.5	1.2	1.3
	3-9 times	0.3	0.4	0.5	0.4	0.6	0.3	0.3
	10+ times	0.3	0.5	0.6	0.8	0.7	0.6	0.8

General

**Annual Prevalence of Weapon Use and Violent Behavior
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Physical fight	No Answer	4.5	4.4	4.7	5.1	5.1	5.7	6.2
	Never	45.9	48.8	46.7	53.1	57.0	60.5	65.3
	1-2 times	31.8	30.1	31.1	27.9	25.4	24.0	20.6
	3-9 times	11.6	10.9	11.6	9.3	8.5	6.6	5.5
	10+ times	6.2	5.8	5.9	4.6	4.0	3.3	2.6
Shouting argument	No Answer	4.9	4.9	5.1	5.4	5.5	5.9	6.4
	Never	41.0	38.9	33.3	34.0	34.4	35.0	36.8
	1-2 times	27.7	27.6	27.1	26.6	25.7	27.0	26.1
	3-9 times	13.8	15.0	18.1	17.8	18.9	18.6	18.7
	10+ times	12.6	13.6	16.4	16.2	15.6	13.5	12.0
Carried gun for hunting	No Answer	4.8	4.9	5.1	5.4	5.5	5.9	6.3
	Never	56.9	54.1	52.2	53.9	55.6	59.2	59.6
	1-2 times	14.4	14.6	14.1	12.6	11.6	10.7	9.7
	3-9 times	10.1	10.7	11.4	11.0	9.9	9.5	9.2
	10+ times	13.7	15.8	17.2	17.1	17.4	14.8	15.3
Carried gun for other reasons	No Answer	5.0	5.0	5.2	5.3	5.6	6.1	6.4
	Never	87.3	85.6	81.9	82.0	82.0	82.7	81.0
	1-2 times	4.0	4.7	6.4	5.5	5.3	5.2	5.2
	3-9 times	1.6	2.0	2.8	2.9	2.6	2.3	2.6
	10+ times	2.1	2.7	3.7	4.3	4.5	3.8	4.9
Carried weapon for hunting	No Answer	5.1	5.0	5.3	5.4	5.7	6.1	6.3
	Never	71.8	69.6	67.0	68.3	68.6	70.6	71.1
	1-2 times	11.1	10.7	10.6	9.3	8.1	7.3	6.6
	3-9 times	5.1	6.1	6.7	6.5	6.4	6.0	5.6
	10+ times	6.9	8.7	10.3	10.5	11.3	10.1	10.3
Carried weapon for other reasons	No Answer	5.0	5.1	5.5	5.6	5.8	5.9	6.4
	Never	82.0	79.4	73.8	73.2	71.6	73.5	72.8
	1-2 times	6.0	7.0	8.6	8.0	7.9	7.1	7.0
	3-9 times	2.9	3.4	4.6	4.7	4.8	4.1	4.0
	10+ times	4.1	5.1	7.6	8.5	9.9	9.4	9.7
Carried gun to school	No Answer	4.7	4.9	5.2	5.4	5.6	5.8	6.4
	Never	94.3	93.3	92.6	92.1	91.4	90.9	90.2
	1-2 times	0.5	0.8	1.1	1.2	1.1	1.2	1.4
	3-9 times	0.1	0.3	0.3	0.4	0.5	0.7	0.5
	10+ times	0.4	0.7	0.8	1.0	1.4	1.4	1.5
Carried weapon to school	No Answer	4.9	5.0	5.2	5.3	5.5	5.8	6.4
	Never	90.8	89.3	86.4	85.0	83.4	83.9	83.0
	1-2 times	3.3	3.9	5.7	5.2	5.4	4.7	4.2
	3-9 times	0.5	0.8	1.2	1.6	2.1	1.7	1.7
	10+ times	0.5	1.0	1.5	2.9	3.6	3.9	4.8

Male



**Annual Prevalence of Weapon Use and Violent Behavior
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Threatened by weapon on school property	No Answer	5.0	5.1	5.3	5.5	5.7	6.2	6.4
	Never	86.9	85.9	84.2	84.0	84.7	85.9	87.0
	1-2 times	6.2	6.5	7.6	7.0	6.2	5.0	4.2
	3-9 times	1.1	1.3	1.7	1.8	1.8	1.5	1.1
Property stolen while at school	10+ times	0.9	1.2	1.2	1.7	1.6	1.3	1.3
	No Answer	5.3	5.3	5.4	5.5	5.7	6.2	6.5
	Never	64.8	61.0	59.4	57.8	61.6	64.4	66.9
	1-2 times	20.7	23.4	24.2	25.0	23.7	21.2	20.1
Missed school: Felt unsafe on school property	3-9 times	6.3	7.1	7.7	8.3	6.4	6.1	4.9
	10+ times	2.8	3.2	3.2	3.3	2.6	2.1	1.5
	No Answer	4.9	5.3	5.3	5.6	5.6	6.0	6.4
	Never	91.6	90.3	90.5	89.9	89.6	89.9	89.3
Missed school: Felt unsafe on way to/from school	1-2 times	2.6	2.9	2.5	2.7	2.7	2.3	2.0
	3-9 times	0.5	0.7	0.7	0.6	0.8	0.7	0.9
	10+ times	0.4	0.8	0.9	1.2	1.2	1.1	1.3
	No Answer	5.0	5.3	5.4	5.7	5.7	6.1	6.6
	Never	93.0	92.3	92.2	91.5	91.3	91.2	90.7
	1-2 times	1.3	1.4	1.2	1.3	1.3	1.3	1.1
	3-9 times	0.3	0.4	0.4	0.4	0.6	0.3	0.4
	10+ times	0.4	0.7	0.8	1.2	1.1	1.1	1.3

**Annual Prevalence of Weapon Use and Violent Behavior
by Indiana Female 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Physical fight	No Answer	2.6	2.8	2.5	2.8	3.0	3.1	3.0
	Never	75.0	73.0	69.1	72.5	76.0	79.6	83.8
	1-2 times	15.3	16.2	19.4	18.2	15.4	13.2	10.2
	3-9 times	4.7	5.5	6.1	4.7	4.1	3.1	2.2
	10+ times	2.4	2.5	2.9	1.8	1.5	1.0	0.8
Shouting argument	No Answer	2.9	3.1	2.6	3.0	3.2	3.2	3.0
	Never	51.0	44.1	36.1	35.2	34.7	37.9	42.1
	1-2 times	25.1	25.2	26.0	26.4	26.7	26.6	26.9
	3-9 times	11.6	14.4	18.5	19.4	19.5	18.6	17.0
	10+ times	9.3	13.2	16.8	16.0	15.9	13.7	11.0
Carried gun for hunting	No Answer	2.7	2.8	2.5	3.0	3.2	3.1	2.9
	Never	88.1	87.4	86.4	87.6	87.9	88.9	89.7
	1-2 times	5.5	5.8	6.6	5.2	4.9	4.3	4.1
	3-9 times	2.2	2.5	2.7	2.4	2.5	2.2	1.9
	10+ times	1.5	1.5	1.8	1.8	1.5	1.4	1.4
Carried gun for other reasons	No Answer	2.6	2.8	2.5	2.9	3.2	3.1	3.1
	Never	95.7	95.3	94.8	94.9	94.5	95.2	95.0
	1-2 times	1.1	1.2	1.7	1.3	1.5	1.0	1.1
	3-9 times	0.2	0.4	0.5	0.4	0.4	0.3	0.3
	10+ times	0.4	0.3	0.4	0.5	0.4	0.4	0.5
Carried weapon for hunting	No Answer	2.8	2.9	2.6	2.9	3.3	3.2	3.1
	Never	92.0	91.6	91.2	91.7	91.8	92.6	93.4
	1-2 times	3.3	3.5	3.8	3.2	2.8	2.2	1.9
	3-9 times	1.1	1.2	1.4	1.1	1.2	1.0	0.9
	10+ times	0.8	0.8	1.1	1.1	0.9	1.0	0.8
Carried weapon for other reasons	No Answer	2.8	2.9	2.5	3.0	3.3	3.2	3.1
	Never	93.3	91.9	90.1	90.6	89.7	91.0	92.0
	1-2 times	2.5	3.4	4.5	3.7	3.9	2.8	2.3
	3-9 times	0.7	1.0	1.4	1.4	1.4	1.3	1.0
	10+ times	0.7	0.8	1.5	1.3	1.8	1.7	1.7
Carried gun to school	No Answer	2.7	2.9	2.5	3.0	3.2	3.2	3.1
	Never	96.9	96.6	96.8	96.4	96.1	96.1	96.2
	1-2 times	0.1	0.2	0.4	0.3	0.3	0.3	0.3
	3-9 times	0.0	0.1	0.1	0.1	0.1	0.2	0.1
	10+ times	0.1	0.1	0.2	0.2	0.3	0.2	0.3
Carried weapon to school	No Answer	2.8	3.0	2.6	3.0	3.2	3.3	3.1
	Never	96.1	94.9	94.1	93.6	93.4	93.6	94.4
	1-2 times	0.8	1.5	2.2	2.2	1.9	1.7	1.1
	3-9 times	0.2	0.3	0.5	0.5	0.6	0.7	0.4
	10+ times	0.1	0.3	0.6	0.7	0.9	0.8	0.9

Female



**Annual Prevalence of Weapon Use and Violent Behavior
by Indiana Female 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Threatened by weapon on school property	No Answer	3.0	3.0	2.7	3.2	3.3	3.3	3.2
	Never	92.0	91.2	91.4	91.3	92.1	93.7	94.7
	1-2 times	4.1	4.6	4.6	4.4	3.6	2.5	1.6
	3-9 times	0.6	0.8	0.7	0.7	0.7	0.4	0.2
Property stolen while at school	10+ times	0.4	0.3	0.4	0.4	0.3	0.2	0.4
	No Answer	3.3	3.1	2.8	3.3	3.4	3.3	3.1
	Never	69.8	63.2	63.3	63.0	65.9	71.2	75.6
	1-2 times	20.8	26.1	26.0	27.0	25.0	21.5	18.3
Missed school: Felt unsafe on school property	3-9 times	4.6	5.5	6.2	5.5	4.7	3.5	2.4
	10+ times	1.5	2.0	1.7	1.2	0.9	0.6	0.6
	No Answer	3.1	3.1	2.8	3.2	3.3	3.3	3.1
	Never	92.4	91.8	91.1	91.2	91.0	91.8	93.0
Missed school: Felt unsafe on way to/from school	1-2 times	3.5	3.9	4.7	4.3	4.2	3.7	3.1
	3-9 times	0.7	0.7	0.9	0.8	1.0	0.8	0.4
	10+ times	0.3	0.5	0.5	0.5	0.4	0.4	0.4
	No Answer	3.0	3.0	2.7	3.1	3.3	3.4	3.1
	Never	94.1	94.5	94.2	94.4	94.1	94.9	94.9
	1-2 times	2.3	1.9	2.2	1.7	1.7	1.2	1.4
	3-9 times	0.3	0.3	0.5	0.4	0.5	0.3	0.2
	10+ times	0.3	0.3	0.3	0.4	0.3	0.3	0.4

**Monthly Prevalence of Weapon Use and Violent Behavior
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Physical fight	No Answer	4.9	4.9	5.0	5.4	5.6	5.9	6.2
	Never	74.9	75.6	75.2	79.0	81.2	83.5	85.5
	1-2 times	15.2	15.0	15.2	12.2	10.4	8.4	6.7
	3-9 times	3.4	3.0	3.0	2.2	1.8	1.4	0.8
	10+ times	1.5	1.5	1.6	1.2	1.0	0.8	0.8
Shouting argument	No Answer	5.3	5.4	5.5	5.7	5.9	6.1	6.2
	Never	60.1	57.2	51.8	53.0	53.7	56.9	59.6
	1-2 times	23.1	23.5	26.4	26.3	25.5	24.6	24.1
	3-9 times	7.5	8.8	10.7	10.0	10.1	8.8	7.3
	10+ times	4.0	5.1	5.6	5.0	4.7	3.6	2.9
Carried gun for hunting	No Answer	5.2	5.4	5.3	5.8	6.0	6.2	6.3
	Never	81.6	81.0	80.8	82.2	83.1	84.6	84.2
	1-2 times	7.4	7.2	7.4	6.1	5.6	4.5	4.5
	3-9 times	3.3	3.4	3.7	3.1	2.9	2.5	2.8
	10+ times	2.5	3.0	2.9	2.8	2.4	2.2	2.3
Carried gun for other reasons	No Answer	5.2	5.3	5.4	5.7	5.9	6.2	6.3
	Never	91.7	90.8	89.9	89.7	89.7	90.0	89.0
	1-2 times	1.8	2.2	2.6	2.3	2.3	1.9	2.3
	3-9 times	0.6	0.8	1.0	0.9	0.9	0.7	0.9
	10+ times	0.7	0.9	1.1	1.4	1.2	1.2	1.5
Carried weapon for hunting	No Answer	5.3	5.4	5.3	5.8	5.9	6.3	6.2
	Never	87.3	86.4	86.0	86.6	87.2	87.5	87.5
	1-2 times	4.2	4.2	4.7	3.8	3.3	2.8	2.8
	3-9 times	1.5	1.9	1.8	1.6	1.6	1.4	1.5
	10+ times	1.7	2.1	2.2	2.1	2.0	2.0	1.9
Carried weapon for other reasons	No Answer	5.3	5.5	5.4	5.8	6.0	6.3	6.3
	Never	89.3	87.8	85.7	86.0	85.3	86.1	85.7
	1-2 times	3.1	3.9	4.8	3.7	4.0	3.0	3.2
	3-9 times	1.0	1.2	1.7	1.7	1.8	1.6	1.4
	10+ times	1.3	1.6	2.5	2.8	2.9	3.0	3.4
Carried gun to school	No Answer	5.1	5.4	5.4	5.8	5.9	6.3	6.3
	Never	94.3	93.7	93.5	93.0	92.7	92.3	92.1
	1-2 times	0.2	0.4	0.5	0.5	0.5	0.6	0.6
	3-9 times	0.1	0.2	0.1	0.2	0.3	0.2	0.3
	10+ times	0.2	0.3	0.4	0.5	0.5	0.6	0.7
Carried weapon to school	No Answer	5.3	5.4	5.5	5.9	6.0	6.3	6.3
	Never	93.3	92.4	91.7	90.7	90.1	90.1	89.8
	1-2 times	0.9	1.3	1.7	1.7	2.0	1.5	1.4
	3-9 times	0.2	0.3	0.4	0.6	0.6	0.5	0.6
	10+ times	0.3	0.5	0.7	1.1	1.4	1.7	1.9

General



**Monthly Prevalence of Weapon Use and Violent Behavior
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Threatened by weapon on school property	No Answer	5.4	5.6	5.5	5.9	6.0	6.3	6.3
	Never	91.6	91.2	91.1	91.1	91.2	91.5	91.7
	1-2 times	2.2	2.2	2.5	1.9	1.9	1.4	1.2
	3-9 times	0.4	0.5	0.4	0.5	0.4	0.4	0.2
Property stolen while at school	10+ times	0.4	0.5	0.5	0.6	0.5	0.5	0.6
	No Answer	5.7	5.7	5.7	6.0	6.1	6.4	6.4
	Never	81.2	79.5	80.7	81.1	83.8	86.0	87.3
	1-2 times	10.6	12.0	11.0	10.8	8.4	6.1	5.0
Missed school: Felt unsafe on school property	3-9 times	1.7	1.9	1.7	1.2	1.1	0.9	0.6
	10+ times	0.8	0.9	0.9	0.8	0.7	0.5	0.7
	No Answer	5.3	5.5	5.5	5.9	6.1	6.4	6.4
	Never	92.8	92.1	92.1	91.8	91.5	91.7	91.4
Missed school: Felt unsafe on way to/from school	1-2 times	1.4	1.7	1.7	1.3	1.6	1.1	1.2
	3-9 times	0.3	0.4	0.3	0.4	0.4	0.4	0.3
	10+ times	0.3	0.4	0.4	0.5	0.5	0.5	0.7
	No Answer	5.8	5.8	5.9	6.4	6.4	6.7	6.6
	Never	92.9	92.6	92.6	92.0	91.9	91.9	91.8
	1-2 times	0.9	1.0	0.9	0.8	0.9	0.7	0.7
	3-9 times	0.2	0.2	0.2	0.3	0.4	0.3	0.2
	10+ times	0.2	0.4	0.4	0.5	0.5	0.4	0.7

General

**Monthly Prevalence of Weapon Use and Violent Behavior
by Indiana Male 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Physical fight	No Answer	6.0	6.0	6.5	6.9	7.2	7.7	7.9
	Never	67.1	69.5	69.2	74.0	76.3	78.2	80.9
	1-2 times	20.3	18.9	18.7	14.4	12.6	10.9	8.9
	3-9 times	4.5	3.7	3.6	2.9	2.4	2.1	1.1
	10+ times	2.3	1.9	2.1	1.8	1.6	1.2	1.3
Shouting argument	No Answer	6.3	6.6	7.1	7.3	7.5	7.9	7.8
	Never	57.7	56.7	52.4	54.5	55.2	57.0	58.2
	1-2 times	24.0	23.8	25.7	24.5	23.9	23.2	23.6
	3-9 times	7.7	8.2	9.6	8.7	9.0	8.2	7.0
	10+ times	4.3	4.8	5.1	5.0	4.4	3.7	3.3
Carried gun for hunting	No Answer	6.4	6.8	7.0	7.5	7.5	8.0	8.0
	Never	71.7	70.5	70.0	71.8	73.0	75.8	75.5
	1-2 times	11.6	11.5	11.4	9.8	9.4	7.5	7.3
	3-9 times	5.6	5.8	6.3	5.6	5.4	4.5	5.1
	10+ times	4.7	5.4	5.3	5.3	4.6	4.3	4.2
Carried gun for other reasons	No Answer	6.4	6.7	7.1	7.5	7.5	8.1	7.9
	Never	88.5	86.7	84.9	84.5	84.6	85.0	83.9
	1-2 times	2.8	3.6	4.3	3.9	4.0	3.5	3.8
	3-9 times	1.0	1.3	1.7	1.7	1.6	1.3	1.7
	10+ times	1.2	1.6	2.1	2.5	2.3	2.1	2.7
Carried weapon for hunting	No Answer	6.5	6.8	7.0	7.6	7.5	8.3	7.9
	Never	81.3	79.5	78.5	79.2	80.1	80.5	80.9
	1-2 times	6.6	6.6	7.4	6.2	5.5	5.0	5.0
	3-9 times	2.5	3.2	3.2	2.9	3.0	2.5	2.8
	10+ times	3.2	3.9	4.0	4.2	3.8	3.7	3.5
Carried weapon for other reasons	No Answer	6.5	6.9	7.1	7.5	7.6	8.2	8.0
	Never	84.9	82.9	79.2	79.1	78.1	79.1	78.8
	1-2 times	4.7	5.3	6.8	5.7	6.0	4.8	5.1
	3-9 times	1.7	2.0	2.7	2.9	3.1	2.7	2.3
	10+ times	2.3	2.8	4.2	4.9	5.1	5.3	5.8
Carried gun to school	No Answer	6.3	6.8	7.0	7.7	7.6	8.1	8.0
	Never	93.0	91.8	91.3	90.4	90.2	89.5	89.4
	1-2 times	0.3	0.6	0.8	0.8	0.8	0.9	1.0
	3-9 times	0.1	0.2	0.2	0.3	0.5	0.4	0.5
	10+ times	0.4	0.6	0.7	0.8	1.0	1.0	1.2
Carried weapon to school	No Answer	6.3	6.9	7.1	7.6	7.6	8.2	8.0
	Never	91.6	90.1	89.0	87.4	86.2	85.8	85.6
	1-2 times	1.3	1.8	2.3	2.4	3.0	2.3	2.3
	3-9 times	0.3	0.5	0.5	0.9	1.0	0.8	0.9
	10+ times	0.4	0.8	1.0	1.7	2.2	2.9	3.3

Male



**Monthly Prevalence of Weapon Use and Violent Behavior
by Indiana Male 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Threatened by weapon on school property	No Answer	6.4	6.9	7.2	7.6	7.6	8.2	8.1
	Never	89.7	89.1	88.5	88.2	88.3	88.3	88.9
	1-2 times	2.7	2.6	2.9	2.5	2.6	2.1	1.7
	3-9 times	0.5	0.6	0.6	0.7	0.6	0.6	0.3
Property stolen while at school	10+ times	0.6	0.8	0.8	1.0	0.9	0.8	1.0
	No Answer	6.7	7.1	7.3	7.7	7.6	8.4	8.1
	Never	79.6	78.0	78.9	78.4	81.7	82.8	84.5
	1-2 times	10.6	11.6	10.8	11.1	8.1	6.6	5.4
Missed school: Felt unsafe on school property	3-9 times	2.0	2.0	1.8	1.6	1.4	1.3	0.9
	10+ times	1.0	1.3	1.2	1.3	1.1	0.8	1.0
	No Answer	6.4	6.9	7.2	7.7	7.7	8.3	8.1
	Never	91.9	90.8	90.6	90.0	89.7	89.6	89.2
Missed school: Felt unsafe on way to/from school	1-2 times	1.1	1.4	1.3	1.1	1.3	0.9	1.2
	3-9 times	0.3	0.4	0.3	0.4	0.5	0.5	0.4
	10+ times	0.3	0.5	0.6	0.8	0.8	0.7	1.1
	No Answer	6.9	7.3	7.6	8.3	8.1	8.7	8.5
	Never	91.9	91.1	90.8	89.7	89.7	89.5	89.2
	1-2 times	0.7	0.8	0.8	0.8	0.9	0.8	0.9
	3-9 times	0.2	0.3	0.2	0.3	0.5	0.3	0.3
	10+ times	0.3	0.5	0.6	0.8	0.9	0.6	1.1

Male

**Monthly Prevalence of Weapon Use and Violent Behavior
by Indiana Female 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Physical fight	No Answer	3.9	3.7	3.4	3.8	4.1	4.2	4.2
	Never	82.7	82.0	81.3	83.9	85.8	88.5	90.3
	1-2 times	10.3	11.0	11.7	10.2	8.4	6.2	4.7
	3-9 times	2.3	2.3	2.5	1.5	1.3	0.8	0.6
	10+ times	0.8	1.0	1.1	0.6	0.5	0.4	0.3
Shouting argument	No Answer	4.4	4.2	3.9	4.2	4.3	4.4	4.3
	Never	62.6	57.7	51.3	51.7	52.4	56.8	61.2
	1-2 times	22.2	23.3	27.1	28.0	27.1	26.0	24.5
	3-9 times	7.2	9.4	11.7	11.1	11.2	9.3	7.6
	10+ times	3.7	5.3	6.1	5.0	5.0	3.5	2.4
Carried gun for hunting	No Answer	4.1	3.9	3.6	4.1	4.4	4.5	4.4
	Never	91.5	91.8	91.5	92.0	92.4	92.7	93.0
	1-2 times	3.1	2.9	3.4	2.6	2.2	1.8	1.7
	3-9 times	0.9	0.9	1.0	0.7	0.6	0.6	0.5
	10+ times	0.3	0.5	0.5	0.5	0.3	0.3	0.3
Carried gun for other reasons	No Answer	4.0	3.8	3.7	4.0	4.3	4.5	4.4
	Never	94.9	95.0	94.9	94.7	94.5	94.6	94.4
	1-2 times	0.7	0.7	0.9	0.7	0.8	0.5	0.8
	3-9 times	0.2	0.2	0.3	0.2	0.2	0.1	0.1
	10+ times	0.2	0.2	0.2	0.3	0.2	0.3	0.3
Carried weapon for hunting	No Answer	4.1	3.9	3.6	4.0	4.4	4.5	4.3
	Never	93.3	93.5	93.5	93.6	93.8	93.9	94.4
	1-2 times	1.8	1.8	1.9	1.7	1.2	0.8	0.7
	3-9 times	0.5	0.5	0.5	0.4	0.3	0.3	0.3
	10+ times	0.3	0.3	0.4	0.3	0.3	0.4	0.3
Carried weapon for other reasons	No Answer	4.1	3.9	3.7	4.2	4.4	4.5	4.3
	Never	93.7	92.9	92.2	92.5	91.9	92.6	92.8
	1-2 times	1.5	2.3	2.8	1.9	2.1	1.4	1.4
	3-9 times	0.4	0.4	0.6	0.6	0.7	0.6	0.5
	10+ times	0.3	0.4	0.7	0.8	0.8	0.9	1.0
Carried gun to school	No Answer	4.0	3.9	3.8	4.1	4.4	4.5	4.4
	Never	95.7	95.7	95.8	95.5	95.1	95.0	95.1
	1-2 times	0.1	0.2	0.2	0.2	0.3	0.2	0.2
	3-9 times	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	10+ times	0.1	0.1	0.1	0.2	0.1	0.3	0.2
Carried weapon to school	No Answer	4.2	4.0	3.8	4.1	4.4	4.5	4.3
	Never	95.1	94.8	94.5	94.0	93.7	94.1	94.4
	1-2 times	0.5	0.8	1.1	1.1	1.0	0.7	0.6
	3-9 times	0.1	0.2	0.3	0.2	0.3	0.2	0.3
	10+ times	0.1	0.2	0.4	0.5	0.6	0.5	0.5

Female



**Monthly Prevalence of Weapon Use and Violent Behavior
by Indiana Female 6th - 12th Graders, 2006**
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Threatened by weapon on school property	No Answer	4.3	4.1	3.8	4.2	4.4	4.4	4.4
	Never	93.4	93.4	93.7	93.9	94.0	94.4	94.8
	1-2 times	1.8	1.8	2.0	1.4	1.2	0.8	0.6
	3-9 times	0.3	0.4	0.3	0.3	0.2	0.1	0.1
Property stolen while at school	10+ times	0.2	0.2	0.2	0.2	0.2	0.2	0.2
	No Answer	4.6	4.2	4.1	4.4	4.5	4.6	4.5
	Never	82.7	81.0	82.6	83.7	85.9	89.1	90.4
	1-2 times	10.7	12.4	11.2	10.6	8.6	5.6	4.6
Missed school: Felt unsafe on school property	3-9 times	1.4	1.9	1.6	0.8	0.7	0.4	0.3
	10+ times	0.6	0.5	0.5	0.4	0.3	0.3	0.2
	No Answer	4.2	4.1	3.8	4.2	4.5	4.5	4.5
	Never	93.6	93.4	93.6	93.7	93.2	93.7	94.0
Missed school: Felt unsafe on way to/from school	1-2 times	1.7	1.9	2.1	1.5	1.7	1.2	1.2
	3-9 times	0.2	0.4	0.3	0.4	0.3	0.3	0.1
	10+ times	0.2	0.2	0.2	0.2	0.2	0.2	0.2
	No Answer	4.6	4.3	4.1	4.4	4.7	4.9	4.6
	Never	93.9	94.2	94.5	94.3	94.0	94.1	94.6
	1-2 times	1.1	1.2	1.0	0.8	0.9	0.6	0.5
	3-9 times	0.3	0.2	0.2	0.2	0.3	0.1	0.1
	10+ times	0.1	0.2	0.2	0.2	0.2	0.2	0.2

Female



IV. GAMBLING

- Gambling during the Last Twelve Months,
Grades 6-12



**Gambling during the Last Twelve Months
by Indiana 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Played cards	No Answer	6.3	6.3	5.9	5.7	5.8	5.7	5.7
	Not at all	75.6	70.3	64.2	63.0	61.4	61.9	59.9
	Less than once a month	10.2	13.1	15.3	16.0	16.2	15.9	16.4
	About once a month	4.5	6.3	8.5	8.9	9.8	9.3	10.1
	About once a week	2.4	2.7	4.3	4.6	5.0	5.4	6.0
	Daily	1.0	1.3	1.8	1.8	1.8	1.8	1.9
Bet on games	No Answer	6.5	6.6	6.1	5.9	5.9	5.7	5.8
	Not at all	75.1	71.0	65.6	66.0	66.4	67.6	67.1
	Less than once a month	10.5	12.4	14.4	14.5	13.6	13.2	13.3
	About once a month	4.5	6.0	7.9	7.8	7.9	7.7	8.0
	About once a week	2.0	2.7	3.9	4.2	4.4	3.9	3.8
	Daily	1.3	1.5	2.0	1.8	1.9	1.9	2.0
Bet on sports	No Answer	6.8	6.8	6.3	6.2	6.3	6.0	6.1
	Not at all	76.5	73.0	68.4	69.1	69.6	70.5	71.4
	Less than once a month	9.9	11.6	13.6	13.0	12.8	12.8	11.9
	About once a month	3.7	5.0	6.4	6.6	6.3	6.1	6.0
	About once a week	2.0	2.4	3.5	3.4	3.2	2.9	2.8
	Daily	1.2	1.3	1.8	1.8	1.8	1.7	1.7
Bought lottery tickets	No Answer	6.7	6.6	6.2	6.0	6.0	5.7	5.9
	Not at all	78.2	77.7	77.0	77.8	78.8	77.7	57.8
	Less than once a month	8.9	9.1	9.4	9.3	8.4	8.7	20.1
	About once a month	3.6	3.6	4.2	3.6	3.9	4.1	8.9
	About once a week	1.6	1.9	2.0	2.2	1.8	2.4	5.1
	Daily	1.1	1.1	1.3	1.2	1.1	1.3	2.2
Gambled in a casino	No Answer	6.6	6.6	6.1	6.0	6.1	5.9	5.9
	Not at all	92.1	91.9	91.6	91.5	91.6	91.1	90.0
	Less than once a month	0.5	0.7	1.0	1.1	1.0	1.4	2.3
	About once a month	0.2	0.2	0.3	0.4	0.4	0.4	0.5
	About once a week	0.1	0.1	0.2	0.2	0.2	0.2	0.4
	Daily	0.3	0.5	0.7	0.8	0.7	0.9	0.9
Felt bad about the amount bet	No Answer	7.8	7.3	6.6	6.1	6.2	5.7	6.0
	Yes	5.2	5.0	5.4	4.6	4.6	4.1	4.5
	No	21.1	25.8	32.4	33.5	34.1	34.7	37.9
Would like to stop betting but could not	I do not bet for money	65.9	61.9	55.6	55.7	55.2	55.6	51.6
	No Answer	7.6	7.4	6.6	6.1	6.2	5.9	5.9
	Yes	4.5	3.6	3.2	2.3	1.9	1.6	1.6
	No	21.5	26.7	33.8	35.1	36.1	36.5	40.1
	I do not bet for money	66.4	62.4	56.3	56.5	55.7	56.0	52.4

General

**Gambling during the Last Twelve Months
by Indiana Male 6th - 12th Graders, 2006
(Values expressed as percentages)**


		Grade						
		6th	7th	8th	9th	10th	11th	12th
Played cards	No Answer	7.3	7.5	7.2	7.1	7.2	7.0	7.3
	Not at all	66.9	60.6	51.2	48.3	44.8	43.8	41.8
	Less than once a month	13.8	16.7	19.3	19.4	20.1	20.9	20.6
	About once a month	6.8	9.0	12.6	14.2	15.7	15.4	16.6
	About once a week	3.8	4.1	6.8	7.8	8.9	9.7	10.6
	Daily	1.5	2.1	2.8	3.1	3.4	3.3	3.1
Bet on games	No Answer	7.5	7.7	7.4	7.3	7.2	7.1	7.5
	Not at all	66.4	60.5	52.3	51.6	50.9	50.4	49.3
	Less than once a month	13.7	16.2	18.6	18.5	17.8	18.4	18.9
	About once a month	7.1	8.8	11.9	12.3	12.6	13.4	13.7
	About once a week	3.2	4.4	6.4	7.2	7.9	7.1	7.1
	Daily	2.1	2.3	3.4	3.2	3.5	3.6	3.4
Bet on sports	No Answer	7.8	8.0	7.7	7.7	7.7	7.5	7.8
	Not at all	67.8	62.6	55.4	55.3	55.4	54.4	56.2
	Less than once a month	13.6	15.9	18.0	17.2	17.4	18.9	17.4
	About once a month	5.9	7.5	9.8	10.5	10.4	10.8	10.7
	About once a week	3.1	4.0	5.9	6.0	5.6	5.3	5.2
	Daily	1.9	2.1	3.1	3.3	3.4	3.0	2.8
Bought lottery tickets	No Answer	7.7	7.8	7.5	7.5	7.4	7.3	7.5
	Not at all	76.3	75.2	74.0	74.1	75.3	72.3	51.7
	Less than once a month	8.8	9.0	9.4	9.5	8.6	9.6	19.0
	About once a month	4.0	4.2	4.7	4.2	4.5	5.3	11.3
	About once a week	2.0	2.3	2.6	2.9	2.3	3.4	7.0
	Daily	1.4	1.5	1.9	1.9	1.9	2.1	3.5
Gambled in a casino	No Answer	7.7	7.8	7.5	7.6	7.6	7.3	7.4
	Not at all	90.6	90.1	89.2	88.6	88.9	88.1	86.4
	Less than once a month	0.7	0.9	1.3	1.6	1.4	2.0	3.3
	About once a month	0.3	0.2	0.4	0.5	0.6	0.7	0.9
	About once a week	0.2	0.1	0.3	0.3	0.3	0.3	0.6
	Daily	0.5	0.8	1.2	1.3	1.2	1.5	1.5
Felt bad about the amount bet	No Answer	8.7	8.7	7.8	7.5	7.6	7.1	7.5
	Yes	7.4	6.9	7.8	7.1	7.4	7.1	7.4
	No	29.1	34.8	43.7	45.5	47.6	49.5	51.9
Would like to stop betting but could not	I do not bet for money	54.9	49.6	40.7	39.8	37.4	36.3	33.3
	No Answer	8.4	8.7	8.1	7.5	7.6	7.5	7.5
	Yes	6.1	4.9	4.6	3.3	2.9	2.5	2.5
	No	30.2	36.2	46.0	48.7	51.5	53.4	56.1
	I do not bet for money	55.4	50.2	41.4	40.6	37.9	36.6	33.9



**Gambling during the Last Twelve Months
by Indiana Female 6th - 12th Graders, 2006
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Played cards	No Answer	5.4	5.1	4.5	4.3	4.4	4.4	3.9
	Not at all	84.3	80.1	77.3	76.7	76.7	78.2	77.8
	Less than once a month	6.6	9.5	11.4	12.9	12.7	11.5	12.4
	About once a month	2.2	3.5	4.4	4.0	4.4	3.8	3.9
	About once a week	1.1	1.2	1.7	1.6	1.5	1.5	1.4
	Daily	0.5	0.5	0.7	0.4	0.4	0.5	0.5
Bet on games	No Answer	5.5	5.3	4.6	4.5	4.5	4.3	4.0
	Not at all	83.9	81.6	78.9	79.4	80.5	83.2	84.8
	Less than once a month	7.3	8.5	10.3	10.8	9.7	8.6	7.7
	About once a month	1.9	3.1	4.0	3.6	3.7	2.6	2.4
	About once a week	0.9	0.9	1.5	1.3	1.2	0.9	0.6
	Daily	0.5	0.5	0.7	0.4	0.4	0.4	0.5
Bet on sports	No Answer	5.8	5.5	4.8	4.7	4.8	4.7	4.4
	Not at all	85.1	83.6	81.4	81.9	82.6	85.1	86.6
	Less than once a month	6.2	7.2	9.2	9.1	8.6	7.4	6.6
	About once a month	1.6	2.4	3.0	2.9	2.6	1.8	1.5
	About once a week	0.9	0.8	1.1	1.0	1.0	0.7	0.4
	Daily	0.5	0.4	0.6	0.4	0.4	0.4	0.5
Bought lottery tickets	No Answer	5.6	5.4	4.8	4.5	4.7	4.3	4.2
	Not at all	80.1	80.3	80.0	81.3	82.1	82.6	63.8
	Less than once a month	9.0	9.2	9.3	9.1	8.2	7.8	21.2
	About once a month	3.2	3.0	3.7	3.0	3.2	3.0	6.7
	About once a week	1.3	1.5	1.5	1.6	1.4	1.6	3.3
	Daily	0.8	0.6	0.7	0.5	0.4	0.5	0.8
Gambled in a casino	No Answer	5.5	5.3	4.7	4.5	4.7	4.6	4.2
	Not at all	93.7	93.8	94.2	94.3	94.2	93.9	93.8
	Less than once a month	0.3	0.6	0.6	0.6	0.6	0.8	1.4
	About once a month	0.2	0.1	0.1	0.2	0.2	0.2	0.2
	About once a week	0.1	0.1	0.1	0.1	0.0	0.1	0.1
	Daily	0.2	0.1	0.3	0.3	0.3	0.4	0.3
Felt bad about the amount bet	No Answer	6.8	5.9	5.2	4.7	4.9	4.4	4.3
	Yes	3.0	3.0	3.1	2.3	1.9	1.3	1.8
	No	13.2	16.6	21.1	22.4	21.8	21.2	24.3
	I do not bet for money	76.9	74.4	70.6	70.6	71.4	73.1	69.6
Would like to stop betting but could not	No Answer	6.9	5.9	5.2	4.8	4.8	4.4	4.3
	Yes	2.9	2.2	1.9	1.4	1.0	0.7	0.7
	No	12.9	17.1	21.7	22.4	22.1	21.2	24.4
	I do not bet for money	77.3	74.8	71.3	71.4	72.0	73.7	70.6

Female



APPENDIX: INSTRUMENT

Indiana Prevention Resource Center

ALCOHOL, TOBACCO AND OTHER DRUG USE SURVEY

© COPYRIGHT 2005, 2002, 2001, 1995, 1993, 1991 The Trustees of Indiana University. All rights reserved.

Questions # 24 - 26 used with permission from the Minnesota Department of Education.

This survey is funded, in part, by a contract with the Indiana Family and Social Services Administration, Division of Mental Health and Addiction, HHS / SAMHSA, CSAP, SAPT Block Grant.

PRIVACY STATEMENT

This survey is completely confidential. Complete the survey in private and deposit your completed survey in the privacy envelope, as instructed by your teacher or supervisor. Do not place your name or any identifying marks on the survey form or envelope. There is no way for anyone to identify your individual responses. The survey envelopes will not be opened until they are received at the statewide processing center.

YOUR PARTICIPATION IS VOLUNTARY

Your participation in this survey is voluntary. If you do not wish to answer a question, just leave it blank. If you do not wish to participate in this survey, you may:

- Return the entire survey form blank
- Answer questions with random answers and then answer "not truthfully at all" to the last question. Your form will be discarded and no one will know that you did not participate.
- Inform the teacher or supervisor that you choose not to participate in the survey.

You will not be penalized in any way for deciding not to participate in this survey. We do value your participation, however. Accurate results are dependent upon getting as many students as possible to volunteer to participate in this confidential survey. The results are used by many state and federal agencies to help determine how much money will be given to youth programs and to evaluate the effectiveness of drug programs in your community. Your help is important to this effort.


Thank you for your assistance.

SCHOOL CODE									
A	B	C	D	E	F	G	H	I	J
<input type="text"/>									
0	0	0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9	9	9



Please enter the unique ten-digit school code provided by your teacher or supervisor.

- Use a No. 2 pencil only.
- Do not use ink, ballpoint, or felt tip pens.
- Make solid marks that fill the circle completely.
- Erase cleanly any marks you wish to change.
- Make no stray marks on this form.
- Any mention of "drug" refers to *non-prescription drugs*.



CORRECT: ●

INCORRECT: ○ X ○

INFORMATION ABOUT YOU:

1. SEX: Male Female

2. GRADE: 6th 7th 8th 9th 10th 11th 12th

3. ETHNIC ORIGIN (mark all that apply): White or Caucasian Black or African-American
 Mexican American Puerto Rican or Other Latin American
 Asian or Asian American Other

4. IN THE PAST YEAR, HOW OFTEN HAVE YOU BEEN ABSENT FROM SCHOOL?

Never 1-5 days 6-10 days 11-19 days 20 or more days

5. DURING THE PAST YEAR, DID YOU ATTEND OR PARTICIPATE IN...(mark all that apply)?

An "Afternoons R.O.C.K. in Indiana" after-school prevention program
 Youth Leadership Programs (L.E.A.D. Initiative, Teen Institute, 4-H Junior Leaders, etc.)
 S.A.D.D., S.T.A.N.D., or other prevention group at your school

6. HOW OFTEN IN THE PAST YEAR HAVE YOU PARTICIPATED IN ORGANIZED FAMILY ACTIVITIES (e.g., picnics, watching movies, trips, games, etc.)?

Never Once per week Twice per week 3 or more times per week

TOBACCO CIGARETTE USE:

7. HAVE YOU EVER SMOKED CIGARETTES?

Never Smoked Occasionally but not regularly Regularly now
 Once or Twice Regularly in the past

8. IF YOU HAVE EVER SMOKED CIGARETTES, AT WHAT AGE DID YOU FIRST SMOKE THEM?

Never Smoked 7 years old or less 8 to 9 years old 10 to 11 years old
 12 or 13 years old 14 to 15 years old 16 to 17 years old 18 years or older

9. HOW OFTEN IN THE PAST YEAR HAVE YOU SMOKED CIGARETTES?

None A few times 1 to 5 cigarettes per day
 About one half pack per day About 1 pack per day
 About 1 and a half packs per day Two or more packs per day

10. HOW OFTEN IN THE PAST MONTH HAVE YOU SMOKED CIGARETTES?

None A few times 1 to 5 cigarettes per day
 About one half pack per day About 1 pack per day
 About 1 and a half packs per day Two or more packs per day

Continue on to the next page

PLEASE DO NOT WRITE IN THIS AREA



SERIAL

SERIAL

**DO NOT REPORT USE UNDER DOCTOR'S ORDERS FOR QUESTIONS ON THIS PAGE****USE OF ALCOHOL AND OTHER DRUGS****LIFETIME USE:**

11. HAVE YOU EVER USED...?

	Never	1-5 Times	6-19 Times	20-40 Times	More Than 40 Times
--	-------	-----------	------------	-------------	--------------------

Snuff/Smokeless tobacco	<input type="radio"/>				
Cigars (Tobacco)	<input type="radio"/>				
Pipe (Tobacco)	<input type="radio"/>				
Alcohol (beer, wine, wine coolers, liquor)	<input type="radio"/>				
Marijuana (hashish or hash oil)	<input type="radio"/>				
Cocaine	<input type="radio"/>				
Crack	<input type="radio"/>				
Inhalants (glue, fumes, amyls)	<input type="radio"/>				
Amphetamines (uppers)	<input type="radio"/>				
Methamphetamines (meth, crank, crystal)	<input type="radio"/>				
Ritalin (non-prescribed use only)	<input type="radio"/>				
Tranquilizers or Sleeping Pills (downers) (non-prescribed)	<input type="radio"/>				
Narcotics (opium, morphine, codeine) (non-prescribed)	<input type="radio"/>				
Heroin	<input type="radio"/>				
LSD (acid)	<input type="radio"/>				
MDMA (ecstacy, XTC, X)	<input type="radio"/>				
Other Psychedelics (psilocybin, mescaline, etc.)	<input type="radio"/>				
Rohypnol (Roofies)	<input type="radio"/>				
GHB	<input type="radio"/>				
Steroids (non-prescribed)	<input type="radio"/>				
A needle or syringe to inject a drug	<input type="radio"/>				

AGE OF FIRST USE:12. IF YOU HAVE EVER USED THESE DRUGS,
AT WHAT AGE DID YOU FIRST USE THEM?

	Never Used	7 or Less Years	8-9 Years	10-11 Years	12-13 Years	14-15 Years	16-17 Years	18 or More Years
--	------------	-----------------	-----------	-------------	-------------	-------------	-------------	------------------

Snuff/Smokeless tobacco	<input type="radio"/>							
Cigars (Tobacco)	<input type="radio"/>							
Pipe (Tobacco)	<input type="radio"/>							
Alcohol (beer, wine, wine coolers, liquor)	<input type="radio"/>							
Marijuana (hashish or hash oil)	<input type="radio"/>							
Cocaine	<input type="radio"/>							
Crack	<input type="radio"/>							
Inhalants (glue, fumes, amyls)	<input type="radio"/>							
Amphetamines (uppers)	<input type="radio"/>							
Methamphetamines (meth, crank, crystal)	<input type="radio"/>							
Ritalin (non-prescribed use only)	<input type="radio"/>							
Tranquilizers or Sleeping Pills (downers) (non-prescribed)	<input type="radio"/>							
Narcotics (opium, morphine, codeine) (non-prescribed)	<input type="radio"/>							
Heroin	<input type="radio"/>							
LSD (acid)	<input type="radio"/>							
MDMA (ecstacy, XTC, X)	<input type="radio"/>							
Other Psychedelics (psilocybin, mescaline, etc.)	<input type="radio"/>							
Rohypnol (Roofies)	<input type="radio"/>							
GHB	<input type="radio"/>							
Steroids (non-prescribed)	<input type="radio"/>							
A needle or syringe to inject a drug	<input type="radio"/>							

DO NOT REPORT USE UNDER DOCTOR'S ORDERS FOR QUESTIONS ON THIS PAGE

SPECIAL QUESTION ON ALCOHOL USE:

13. HOW MANY TIMES IN THE LAST TWO WEEKS HAVE YOU HAD FIVE OR MORE ALCOHOLIC DRINKS
 (Beer, Wine, Liquor) AT A SITTING?

- None Once Twice 3 to 5 times 6 to 9 times 10 or more times

ANNUAL USE:

14. HOW MANY TIMES IN THE LAST YEAR
 HAVE YOU USED...

	Never	1-5 Times	6-19 Times	20-40 Times	More Than 40 Times
--	-------	-----------	------------	-------------	--------------------

Snuff/Smokeless tobacco	<input type="radio"/>				
Cigars (Tobacco)	<input type="radio"/>				
Pipe (Tobacco)	<input type="radio"/>				
Alcohol (beer, wine, wine coolers, liquor)	<input type="radio"/>				
Marijuana (hashish or hash oil)	<input type="radio"/>				
Cocaine	<input type="radio"/>				
Crack	<input type="radio"/>				
Inhalants (glue, fumes, amyls)	<input type="radio"/>				
Amphetamines (uppers)	<input type="radio"/>				
Methamphetamines (meth, crank, crystal)	<input type="radio"/>				
Ritalin (non-prescribed use only)	<input type="radio"/>				
Tranquilizers or Sleeping Pills (downers) (non-prescribed)	<input type="radio"/>				
Narcotics (opium, morphine, codeine) (non-prescribed)	<input type="radio"/>				
Heroin	<input type="radio"/>				
LSD (acid)	<input type="radio"/>				
MDMA (ecstacy, XTC, X)	<input type="radio"/>				
Other Psychedelics (psilocybin, mescaline, etc.)	<input type="radio"/>				
Rohypnol (Roofies)	<input type="radio"/>				
GHB	<input type="radio"/>				
Steroids (non-prescribed)	<input type="radio"/>				
A needle or syringe to inject a drug	<input type="radio"/>				

USE IN PAST MONTH:

15. HOW MANY TIMES IN THE LAST
 MONTH (30 DAYS) HAVE YOU USED...

	Never	1-5 Times	6-19 Times	20-40 Times	More Than 40 Times
--	-------	-----------	------------	-------------	--------------------

Snuff/Smokeless tobacco	<input type="radio"/>				
Cigars (Tobacco)	<input type="radio"/>				
Pipe (Tobacco)	<input type="radio"/>				
Alcohol (beer, wine, wine coolers, liquor)	<input type="radio"/>				
Marijuana (hashish or hash oil)	<input type="radio"/>				
Cocaine	<input type="radio"/>				
Crack	<input type="radio"/>				
Inhalants (glue, fumes, amyls)	<input type="radio"/>				
Amphetamines (uppers)	<input type="radio"/>				
Methamphetamines (meth, crank, crystal)	<input type="radio"/>				
Ritalin (non-prescribed use only)	<input type="radio"/>				
Tranquilizers or Sleeping Pills (downers) (non-prescribed)	<input type="radio"/>				
Narcotics (opium, morphine, codeine) (non-prescribed)	<input type="radio"/>				
Heroin	<input type="radio"/>				
LSD (acid)	<input type="radio"/>				
MDMA (ecstacy, XTC, X)	<input type="radio"/>				
Other Psychedelics (psilocybin, mescaline, etc.)	<input type="radio"/>				
Rohypnol (Roofies)	<input type="radio"/>				
GHB	<input type="radio"/>				
Steroids (non-prescribed)	<input type="radio"/>				
A needle or syringe to inject a drug	<input type="radio"/>				

Continue on to the next page

CONSEQUENCES OF USE:

16. HOW MANY TIMES HAVE YOU EXPERIENCED THE FOLLOWING DUE TO YOUR DRINKING OR DRUG USE DURING THE PAST YEAR?

	Never	Once	Twice	3-5 Times	6-9 Times	10 or More Times
--	-------	------	-------	-----------	-----------	------------------

Had a hangover	<input type="radio"/>					
Performed poorly on a test or project	<input type="radio"/>					
Been in trouble with the police	<input type="radio"/>					
Been in trouble with your parents	<input type="radio"/>					
Damaged property, pulled fire alarms, etc.	<input type="radio"/>					
Got into a fight or argument	<input type="radio"/>					
Got nauseated or vomited	<input type="radio"/>					
Driven a car while under the influence	<input type="radio"/>					
Ridden in a car driven by someone who had been drinking alcohol	<input type="radio"/>					
Missed school	<input type="radio"/>					
Had a memory loss	<input type="radio"/>					

BELIEFS AND PERCEPTIONS**BELIEFS ABOUT DRUGS:**

17. HOW MUCH DO YOU THINK PEOPLE RISK HARMING THEMSELVES (physically or in other ways) IF THEY...

	No Risk	Slight Risk	Moderate Risk	Great Risk
--	---------	-------------	---------------	------------

Smoke one or more packs of cigarettes per day	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Smoke marijuana (pot) occasionally	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Smoke marijuana regularly	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Use cocaine occasionally	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Use cocaine regularly	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Take one or two drinks of alcohol (beer, wine, liquor) occasionally	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Have five or more drinks once or twice each weekend	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

PEER PERCEPTIONS:

18. HOW DO YOU THINK YOUR CLOSE FRIENDS FEEL (or would feel) ABOUT YOU DOING EACH OF THE FOLLOWING THINGS?

	Strongly Approve	Approve	Don't Know	Disapprove	Strongly Disapprove
--	------------------	---------	------------	------------	---------------------

Smoke one or more packs of cigarettes per day	<input type="radio"/>				
Smoke marijuana (pot) occasionally	<input type="radio"/>				
Smoke marijuana regularly	<input type="radio"/>				
Use cocaine occasionally	<input type="radio"/>				
Use cocaine regularly	<input type="radio"/>				
Take one or two drinks of alcohol (beer, wine, liquor) occasionally	<input type="radio"/>				
Have five or more drinks once or twice each weekend	<input type="radio"/>				

Continue on to the next page

SERIAL

**SAFETY AND SECURITY****NEIGHBORHOOD ENVIRONMENT:**

19. IN YOUR NEIGHBORHOOD, HOW SAFE DO YOU FEEL...



Being out alone during the day

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Being out alone after dark

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Being inside your home alone during the day

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Being inside your home alone after dark

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

While traveling to or from school

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

SCHOOL ENVIRONMENT:

20. IN YOUR SCHOOL, HOW SAFE DO YOU FEEL...



While in class

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Alone in the hallways

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Staying for after-school activities

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

PERSONAL SAFETY ISSUES:

21. IN THE PAST YEAR, HOW OFTEN HAVE YOU...



Been in a physical fight

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Been in a serious argument involving shouting

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried a gun for hunting or target practice

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried a gun for other reasons

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried some other weapon for hunting

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried some other weapon for other reasons

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried a gun to school

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried some other weapon to school

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Been threatened by a weapon on school property

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Had your property stolen or damaged while on school property

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Missed school because you felt unsafe on school property

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Missed school because you felt unsafe on your way to or from school

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

22. IN THE PAST MONTH, HOW OFTEN HAVE YOU...



Been in a physical fight

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Been in a serious argument involving shouting

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried a gun for hunting or target practice

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried a gun for other reasons

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried some other weapon for hunting

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried some other weapon for other reasons

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried a gun to school

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Carried some other weapon to school

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Been threatened by a weapon on school property

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Had your property stolen or damaged while on school property

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Missed school because you felt unsafe on school property

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Missed school because you felt unsafe on your way to or from school

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
-----------------------	-----------------------	-----------------------	-----------------------

Continue on to the next page

AFTER SCHOOL HOURS (approximately 3 p.m. to 6 p.m.)

23. THERE ARE ABOUT 180 SCHOOL DAYS PER YEAR. ON HOW MANY OF THOSE DAYS THIS YEAR WILL YOU SPEND MOST OF THE AFTER SCHOOL HOURS...

(mark all that apply)

None	1 to 9	10 to 29	30 to 59	60 to 89	90 to 129	130 to 180
------	--------	----------	----------	----------	-----------	------------

Participating in...

- or playing on official school sports teams
- official school arts, music, or drama programs
- other official school activities such as clubs and intramural
- supervised activities at youth centers (Boys & Girls Clubs, Parks Department, etc.)
- extended day programs at school
- other supervised youth activities (4-H, scouts, little league, etc.)
- classes such as music, art, computer, dance, sport lessons, etc.

Socializing...

- with friends without adults supervision (e.g., going to the mall, hanging out in private homes, on streets, in parks, etc.)
- at home without adult supervision
- at home with adult supervision

Working....

- for pay outside your home
- for pay inside a home (housekeeping, child care, etc.)
- as a volunteer outside your home
- as a volunteer inside a home (housekeeping, child care, etc.)

GAMBLING

24. DURING THE LAST 12 MONTHS, HOW OFTEN HAVE YOU DONE THESE ACTIVITIES?

Not at all	Less than once a month	About once a month	About once a week	Daily
------------	------------------------	--------------------	-------------------	-------

- Played cards for money
- Bet money on games of personal skill like pool, golf, or bowling
- Bet money on sports teams
- Bought lottery tickets or scratch offs
- Gambled in a casino

25. DURING THE LAST 12 MONTHS, HAVE YOU EVER FELT BAD ABOUT THE AMOUNT YOU BET, OR ABOUT WHAT HAPPENS WHEN YOU BET MONEY?

Yes No I don't bet for money

26. DURING THE LAST 12 MONTHS, HAVE YOU EVER FELT THAT YOU WOULD LIKE TO STOP BETTING MONEY BUT DIDN'T THINK YOU COULD?

Yes No I don't bet for money

27. HOW TRUTHFULLY HAVE YOU ANSWERED THESE QUESTIONS?

Not truthfully at all Somewhat truthfully Completely truthfully

PLEASE DO NOT WRITE IN THIS AREA



SERIAL

FORM SS-02

Indiana Prevention Resource Center

ALCOHOL, TOBACCO AND OTHER DRUG USE SURVEY

© COPYRIGHT 2005, 2002, 2001, 1995, 1993, 1991 The Trustees of Indiana University. All rights reserved.

Questions # 24 - 26 used with permission from the Minnesota Department of Education.



THANK YOU
FOR YOUR PARTICIPATION



THE INDIANA PREVENTION RESOURCE CENTER
2006 PREVALENCE STATISTICS
MAIN FINDINGS



ALCOHOL, TOBACCO AND OTHER DRUG USE BY INDIANA CHILDREN AND ADOLESCENTS

Survey Conducted March & April 2006
Report Dated August 2006

Conducted by:
Indiana Prevention Resource Center

Published by:
Institute for Drug Abuse Prevention

Department of Applied Health Science
School of Health, Physical Education, and Recreation
Indiana University

Funded, in part by a contract with the
Indiana Family and Social Services Administration
Division of Mental Health and Addiction, and the
School of Health, Physical Education, and Recreation

Contact:
Indiana Prevention Resource Center
2735 East 10th street
Creative Arts Building, Room 110
Bloomington, IN 47408
<http://www.drugs.indiana.edu>
drugprc@indiana.edu
Toll Free: 1-800-346-3077
Fax: 812-855-4940

