

THE INDIANA PREVENTION RESOURCE CENTER  
2007 PREVALENCE STATISTICS  
MAIN FINDINGS

**ALCOHOL, TOBACCO AND OTHER DRUG USE  
BY INDIANA CHILDREN AND ADOLESCENTS**





THE INDIANA PREVENTION RESOURCE CENTER  
2007 PREVALENCE STATISTICS  
MAIN FINDINGS

# ALCOHOL, TOBACCO AND OTHER DRUG USE BY INDIANA CHILDREN AND ADOLESCENTS

Survey Conducted March & April 2007  
Report Dated August 2007

*Conducted by:*  
Indiana Prevention Resource Center  
*Published by:*  
Institute for Drug Abuse Prevention  
  
Department of Applied Health Science  
School of Health, Physical Education, and Recreation  
Indiana University

*Contact:*  
**Indiana Prevention Resource Center**  
501 North Morton Street Suite 110  
Bloomington, IN 47404  
  
<http://www.drugs.indiana.edu>  
drugprc@indiana.edu  
Toll Free: 1-800-346-3077  
Fax: 812-855-4940

*Funded, in part by a contract with the  
Indiana Family and Social Services Administration  
Division of Mental Health and Addiction, and the  
School of Health, Physical Education, and Recreation*

**Source Citation:**

Gassman, R., Jun, M. K., Samuel, S., Martin, E. V., Lee, J., Agley, J. D., Athavale, A. M., Choi, R. U., DeWitt, N. K., Ickes, N. L., Lim, H. J., Mawanda, F., Mueller, E. A., Park, S. J., Pelto-Wheeler, S. J., Roditis, M. L., Shin, H., Smith, D. C., Smith, N. R., and Wells, T. J. (2007). Alcohol, Tobacco, and Other Drug Use by Indiana Children and Adolescents: The Indiana Prevention Resource Center Survey – 2007 (IDAP Monograph No. 07-01). Bloomington, IN: Indiana Prevention Resource Center.

# Table of Contents

<b>I. Report .....</b>	<b>1</b>																																									
Introduction .....	2																																									
Results .....	7																																									
Methodology .....	29																																									
How to Interpret the Results .....	41																																									
Future of the Indiana Survey .....	45																																									
Note on the Reporting of Detailed Data .....	46																																									
References .....	48																																									
<b>II. Statewide Data .....</b>	<b>51</b>																																									
<table><thead><tr><th style="background-color: #d3d3d3;"></th><th style="background-color: #d3d3d3;">Grade</th><th style="background-color: #d3d3d3;">6<sup>th</sup></th><th style="background-color: #d3d3d3;">7<sup>th</sup></th><th style="background-color: #d3d3d3;">8<sup>th</sup></th><th style="background-color: #d3d3d3;">9<sup>th</sup></th><th style="background-color: #d3d3d3;">10<sup>th</sup></th><th style="background-color: #d3d3d3;">11<sup>th</sup></th><th style="background-color: #d3d3d3;">12<sup>th</sup></th></tr></thead><tbody><tr><td>Percentage Reporting Lifetime Use of ATOD .....</td><td>52</td><td>56</td><td>60</td><td>64</td><td>68</td><td>72</td><td>76</td></tr><tr><td>Percentage Reporting Annual Use of ATOD .....</td><td>53</td><td>57</td><td>61</td><td>65</td><td>69</td><td>73</td><td>77</td></tr><tr><td>Percentage Reporting Monthly Use of ATOD .....</td><td>54</td><td>58</td><td>62</td><td>66</td><td>70</td><td>74</td><td>78</td></tr><tr><td>Percentage Reporting Daily and Special Use of ATOD .....</td><td>55</td><td>59</td><td>63</td><td>67</td><td>71</td><td>75</td><td>79</td></tr></tbody></table>		Grade	6 <sup>th</sup>	7 <sup>th</sup>	8 <sup>th</sup>	9 <sup>th</sup>	10 <sup>th</sup>	11 <sup>th</sup>	12 <sup>th</sup>	Percentage Reporting Lifetime Use of ATOD .....	52	56	60	64	68	72	76	Percentage Reporting Annual Use of ATOD .....	53	57	61	65	69	73	77	Percentage Reporting Monthly Use of ATOD .....	54	58	62	66	70	74	78	Percentage Reporting Daily and Special Use of ATOD .....	55	59	63	67	71	75	79	
	Grade	6 <sup>th</sup>	7 <sup>th</sup>	8 <sup>th</sup>	9 <sup>th</sup>	10 <sup>th</sup>	11 <sup>th</sup>	12 <sup>th</sup>																																		
Percentage Reporting Lifetime Use of ATOD .....	52	56	60	64	68	72	76																																			
Percentage Reporting Annual Use of ATOD .....	53	57	61	65	69	73	77																																			
Percentage Reporting Monthly Use of ATOD .....	54	58	62	66	70	74	78																																			
Percentage Reporting Daily and Special Use of ATOD .....	55	59	63	67	71	75	79																																			
Mean Age of First Time Use of Alcohol, Tobacco, and Other Drugs .....	80																																									
<b>III. Regional Data .....</b>	<b>81</b>																																									
<table><thead><tr><th style="background-color: #d3d3d3;"></th><th style="background-color: #d3d3d3;">Grade</th><th style="background-color: #d3d3d3;">6<sup>th</sup></th><th style="background-color: #d3d3d3;">7<sup>th</sup></th><th style="background-color: #d3d3d3;">8<sup>th</sup></th><th style="background-color: #d3d3d3;">9<sup>th</sup></th><th style="background-color: #d3d3d3;">10<sup>th</sup></th><th style="background-color: #d3d3d3;">11<sup>th</sup></th><th style="background-color: #d3d3d3;">12<sup>th</sup></th></tr></thead><tbody><tr><td>Percentage Reporting Lifetime Use of ATOD .....</td><td>82</td><td>86</td><td>90</td><td>94</td><td>98</td><td>102</td><td>106</td></tr><tr><td>Percentage Reporting Annual Use of ATOD .....</td><td>83</td><td>87</td><td>91</td><td>95</td><td>99</td><td>103</td><td>107</td></tr><tr><td>Percentage Reporting Monthly Use of ATOD .....</td><td>84</td><td>88</td><td>92</td><td>96</td><td>100</td><td>104</td><td>108</td></tr><tr><td>Percentage Reporting Daily and Special Use of ATOD .....</td><td>85</td><td>89</td><td>93</td><td>97</td><td>101</td><td>105</td><td>109</td></tr></tbody></table>		Grade	6 <sup>th</sup>	7 <sup>th</sup>	8 <sup>th</sup>	9 <sup>th</sup>	10 <sup>th</sup>	11 <sup>th</sup>	12 <sup>th</sup>	Percentage Reporting Lifetime Use of ATOD .....	82	86	90	94	98	102	106	Percentage Reporting Annual Use of ATOD .....	83	87	91	95	99	103	107	Percentage Reporting Monthly Use of ATOD .....	84	88	92	96	100	104	108	Percentage Reporting Daily and Special Use of ATOD .....	85	89	93	97	101	105	109	
	Grade	6 <sup>th</sup>	7 <sup>th</sup>	8 <sup>th</sup>	9 <sup>th</sup>	10 <sup>th</sup>	11 <sup>th</sup>	12 <sup>th</sup>																																		
Percentage Reporting Lifetime Use of ATOD .....	82	86	90	94	98	102	106																																			
Percentage Reporting Annual Use of ATOD .....	83	87	91	95	99	103	107																																			
Percentage Reporting Monthly Use of ATOD .....	84	88	92	96	100	104	108																																			
Percentage Reporting Daily and Special Use of ATOD .....	85	89	93	97	101	105	109																																			
<b>VI. Trends and Prevalence Graphs .....</b>	<b>111</b>																																									
Trends Graphs .....	112																																									
Prevalence Graphs .....	126																																									

## V. Frequency Tables ..... 137

	Gender	Both	Male	Female
Age Distribution of First Time ATOD Use .....	138	175	212	
Lifetime Use of ATOD .....	146	183	220	
Annual Use of ATOD .....	150	187	224	
Monthly Use of ATOD .....	154	191	228	
Binge Drinking in the Past Two Weeks .....	158	195	232	
Main Sources of Alcoholic Beverages .....	158	195	232	
Most Important Reasons for Drinking .....	159	196	233	
Consequences of ATOD Use .....	160	197	234	

### Risk and Protective Factors

	Gender	Both	Male	Female
Perceived Risk of Selected Substance Use Behavior .....	163	200	237	
Perceived Peer Approval of Using Selected Substances .....	164	201	238	
Perceived Parental Approval of Using Selected Substances .....	165	202	239	
Days Being Absent from School in the Past Year .....	166	203	240	
Participation in Some Form of Camp .....	166	203	240	
Participation in Organized Family Events .....	166	203	240	
Number of Days per School Year Spent in Various Afterschool Activities .....	167	204	241	
Perceived Safety in Neighborhood and in School .....	170	207	244	
Annual Prevalence of Weapon Use and Violent Behavior .....	171	208	245	

### Gambling

	Gender	Both	Male	Female
Gambling during the Last Twelve Months .....	173	210	247	

## Appendix: Instrument ..... 249

Data tables and graphs can be downloaded from the IPRC website:  
[http://www.drugs.indiana.edu/data-survey\\_monograph.html](http://www.drugs.indiana.edu/data-survey_monograph.html)



# I. REPORT

• Introduction .....	2
• Results .....	7
• Methodology .....	29
• How to Interpret the Results .....	41
• Future of the Indiana Survey .....	45
• Note on the Reporting of Detailed Data .....	46
• References .....	48

# INTRODUCTION

## Overview

---

In Spring, 2007, the Indiana Prevention Resource Center (IPRC) conducted the Seventeenth Annual Survey of Alcohol, Tobacco, and Other Drug Use by Indiana Children and Adolescents, referred to hereafter as the Indiana Survey. The IPCR has conducted the Indiana Survey, managed and reported the resulting data since 1991. This project is administered through a contract with the Division of Mental Health and Addiction (DMHA) of the Indiana Family and Social Services Administration (FSSA). The purpose of this project is to provide data for state and local planning with respect to the use of alcohol, tobacco, and other drugs (ATOD), gambling behaviors, and risk and protective factors.

In March and April, 2007, local school officials administered surveys to students in grades 6 through 12 in 478 schools throughout Indiana. A total of 175,712 youth from both public and nonpublic schools completed a written survey that asked about their use of various drugs, the age of onset of first use of various drugs, and risk and protective factors. This resulted in 158,632 usable surveys, the primary source of the data reported here. The Methodology section describes this process in more detail. The instrument is in the appendix.

This report summarizes the findings of the 2007 Indiana Survey. These findings include statistics and other information about the prevalence of ATOD use, gambling behaviors, and risk and protective factors. Statewide prevalence-of-use data are shown for past years to help distinguish trends and facilitate comparisons among years. National prevalence estimates, reported by Monitoring the Future (MTF) (Johnston, O'Malley, Bachman, & Schulenberg, 2005), are provided to facilitate comparison with the United States as a whole. For a summary of trends and statistically significant findings, see the Results section.

Other surveys, such as the Youth Risk Behavior Surveillance System (YRBSS) and MTF, collect representative state-level data for Indiana. However, the results of these data are not reported for sub-state regions, nor are local results available to local participants. The strength of the Indiana Survey is its ability to describe reported ATOD use at the local level. This reporting facilitates local needs assessment, planning, and evaluation of drug abuse prevention activities. The IPCR provides a report of local results to each participating school corporation. Data from an individual school corporation is the property of that corporation. Local data may neither be released nor obtained without the permission of the corporation where the survey was conducted.

## National Outcome Measures (NOMs)

The Government Performance and Results Act of 1993 requires government agencies to report the results of their activities. In response, the United States Department of Health and Human Services (HHS), Substance Abuse and Mental Health Services Administration (SAMHSA) designated a set of measures, called the National Outcome Measures (NOMs), for substance abuse prevention. SAMHSA collects the NOMs through its National Survey on Drug Use and Health (NSDH) for all states, and it reports these data for the states and sub-state regions. The Indiana Survey collects data on the majority of the NOMs, and the IPRC also reports that data for Indiana as a whole and for the sub-state regions.

Increasingly, federal and state agencies require the NOMs for needs assessments and evaluation of local prevention programs that they fund. To compete for grants and to measure outcomes without developing new surveillance capacity, communities will need to be able to capture, track, and analyze the NOMs at the local level.

A unique benefit of participating in the Indiana Survey is that participants receive NOMs data on their own schools and communities. The Indiana Survey collects NOMs data on alcohol, tobacco, marijuana and a variety of other illicit drugs for the following measures:

- thirty day prevalence of use,
- perceived risk of harm,
- age of first use, and
- perception of peer disapproval.

## Changes to the Survey Instrument

### *New Items*

We added the following items to the survey in 2007. Response choices, if any, follow the item in a bulleted list.

Age of respondent

During the past month (30 days), how did you usually get your alcohol?

- I did not drink alcohol during the past month (30 days)
- I bought it at a store such as a liquor store, convenience store, supermarket,

discount store, or gas station

- I bought it at a restaurant, bar, or club
- I bought it at a public event such as a concert or sporting event
- I gave someone else money to buy it for me
- A person 21 years or older gave it to me
- I took it from a store
- I got it from a family member
- I got it some other way

What have been the most important reasons for your drinking alcoholic beverages?  
(mark all that apply)

- I did not drink alcoholic beverages
- To experiment (to see what it's like)
- To relax or relieve tension
- To feel good or get high
- To seek deeper insights and understanding
- To have a good time with my friends
- To fit in with a group I like
- To get away from my problems or troubles
- Because of boredom, nothing else to do
- Because of anger or frustration
- To get through the day
- To increase the effects of some other drug(s)
- To decrease (offset) the effects of some other drug(s)
- To get to sleep
- Because it tastes good
- Because I am hooked ( I feel I have to drink)

How many times have you experienced the following due to your drinking or drug use during the past year? The following response options were added as components of the CRAFFT adolescent screening test (Knight, Sherritt, Harris, Gates, & Chang, 2003):

- Forgot things you did while using alcohol or drugs

- Your family or friends ever told you that you should cut down on your drinking or drug use
- Used alcohol or drugs to relax, feel better about yourself, or fit in
- Used alcohol or drugs while you were by yourself or alone

How do you think your parents/guardians feel (or would feel) about you doing each of the following things?

- Smoke one or more pack of cigarettes per day
- Smoke marijuana (pot) occasionally
- Smoke marijuana regularly
- Use cocaine occasionally
- Use cocaine regularly
- Take one or two drinks of alcohol (beer, wine, liquor) occasionally
- Have five or more drinks once or twice each weekend

During the last 12 months, how often have you done these activities? The following response option was added:

- Played online for money

## ***Changes to Existing Items***

The following existing items were modified in the 2007 survey to increase their usefulness, to update their content, or to align them with prevailing data collection standards. Response choices, if any, follow the item in a bulleted list. Words added this year appear in *italics*.

Ethnicity (*Hispanic or non-Hispanic*)

Race (*White or Caucasian, Asian, Black or African-American, Native Hawaiian or Other Pacific Islander, American Indian or Alaskan Native, Other*)

[To items about lifetime use, age of first use, annual use and past-month use]

- Pipe (*tobacco, hookah, water pipe*)
- Ritalin, Adderall (*non-prescribed use only*)
- Narcotics (*opium, morphine, codeine, OxyContin*)
- Over-the-counter drugs (*non-medical use*)

If you have ever used these drugs, at what age did you first use them? [In past years,

the response options to this item were age ranges such as 8-9 yrs, 10-11 yrs, etc.]

- 8 yrs, 9 yrs, 10 yrs, 11 yrs, 12 yrs, 13 yrs, 14 yrs, 15 yrs, 16 yrs, 17 yrs

## **Regional Reporting**

---

This year, for the second time, this report presents prevalence data broken out at the level of FSSA's sub-state planning regions. These eight regions are aggregates of the State's 92 counties, as indicated in Table 1. Prevalence data for sub-state planning regions provides more targeted information than statewide data. Sub-state regional results are detailed in the Results section.

Table 1. *Family and Social Service Administrations Planning Regions*

Sub-state Regions	Counties
Central	Boone, Hamilton, Hancock, Hendricks, Johnson, Marion, Morgan, Shelby
East	Blackford, Delaware, Fayette, Grant, Henry, Jay, Madison, Randolph, Rush, Union, Wayne
North Central	Cass, Elkhart, Fulton, Howard, Kosciusko, La Porte, Marshall, Miami, St. Joseph, Tipton, Wabash
Northeast	Adams, Allen, De Kalb, Huntington, LaGrange, Noble, Steuben, Wells, Whitley
Northwest	Jasper, Lake, Newton, Porter, Pulaski, Starke
West	Benton, Carroll, Clay, Clinton, Fountain, Monroe, Montgomery, Owen, Parke, Putnam, Sullivan, Tippecanoe, Vermillion, Vigo, Warren, White
Southeast	Brown, Bartholomew, Decatur, Franklin, Lawrence, Jackson, Jennings, Ripley, Dearborn, Orange, Washington, Scott, Jefferson, Ohio, Switzerland, Crawford, Clark, Harrison, Floyd
Southwest	Greene, Knox, Daviess, Martin, Gibson, Pike, Dubois, Posey, Vanderburgh, Warrick, Spencer, Perry

# RESULTS

This section of the monograph summarizes:

- statistically significant changes in statewide prevalence measures from 2006 to 2007.
- regional findings.
- important patterns relating risk and protective factors to substance use.

Data tables included in this monograph include statewide prevalence rates by grade for lifetime, annual, and monthly use of all drugs, as well as daily and special use of selected drugs.

Other data tables and graphs that this monograph has included in the past will now be available for download in a supplemental report via our website at <http://www.drugs.indiana.edu>.

For details of the data included in the supplemental report, see the note on reporting of detailed data, at the end of this monograph.

## **Alcohol, Tobacco, and Other Drugs**

### ***Statewide Prevalence Rates***

For many measures and grades, the Indiana Survey results yielded no statistically significant difference between the two years. Because this section focuses on changes, prevalence rates for certain grades and drugs may not be mentioned in this section. Complete results from the Indiana Survey appear in subsequent sections of this report.

### **Gateway Drugs**

Research has shown that the majority of individuals' illicit drug use occurs only after they use cigarettes, alcohol, or marijuana. These three are known as the "gateway drugs." (Bailey, 1992; Donovan & Jessor, 1983; Fleming, Leventhal, Glynn, & Ershler, 1989; Golub & Johnson, 2001; Kandel & Yamaguchi, 1993).

## *Gateway Drug Use in Grades 6-8*

### **Tobacco use in Grades 6-8**

Statewide, the reported prevalence of cigarette, smokeless tobacco and alcohol use held steady or declined in Grades 6, 7, and 8. However, cigar and pipe use increased in Grades 7 and 8 compared to last year. The observed increase in pipe smoking may be a result of adding hookah and water pipe to the survey this year.

### **Alcohol use in Grades 6-8**

The reported lifetime and annual use of alcohol among youth in Grades 6 and 8 decreased, and monthly prevalence decreased for students in Grades 6, 7 and 8. Daily prevalence of drinking alcohol did not change for these grade levels from last year.

The reported prevalence of binge drinking increased for students in Grades 6 through 8 compared to last year. This result should be interpreted with caution, however. Formatting changes in the instrument gave the binge drinking item a more visible placement on the survey this year. This changed accompanied a quadrupling of the response rate for this item compared to last year. As a result, more binge drinkers may have responded this year.

Higher rates of response typically improve the validity of the results. Thus what appears to be an increase between this year and last may instead reflect the survey's underestimation of binge drinking in the past.

### **Marijuana use in Grade 6-8**

The reported lifetime and annual prevalence of marijuana use increased for students in Grade 7 compared to last year. It is too, early, however to say whether this indicates a reverse in the decreasing trend seen in previous years.

## *Gateway Drug Use in Grades 9-12*

### **Tobacco use in grades 9-12.**

Reported lifetime, annual and monthly use of cigarettes in Grades 9-12 decreased again, consistent with the trend observed in previous year. Reported use of smokeless tobacco use held steady or decreased in these grades.

In contrast, reported daily use of cigars increased among youth in Grades 9-12. In conjunction, lifetime, annual and monthly use of pipe smoking increased for youth in these grades. As previously mentioned, the increased prevalence in pipe smoking may be a result of adding hookah and water pipe to the survey this year.

## **Alcohol use in Grades 9-12**

The reported lifetime, annual and monthly prevalence of alcohol use among students in Grades 9 through 12 continued to decline.

The reported prevalence of binge drinking increased for youth in Grades 9 through 12 compared to last year. This increase may partly reflect this year's higher rate of response to this item (as explained under the heading "Alcohol use in Grades 6-8," above).

## **Marijuana use in Grades 9-12**

The reported lifetime, annual and monthly prevalence of marijuana use for older students held steady or decreased as compared with 2006. The reported daily prevalence of marijuana use among youth in Grades 9 and 12 decreased, and monthly prevalence decreased for students in Grades 9, 11 and 12. In addition, lifetime prevalence of marijuana use decreased for students in Grades 9, 11 and 12. These results are consistent with the continuing trend in decreased marijuana use seen in previous years. There was no change in the prevalence of marijuana use for youth in Grade 10.

## **Methamphetamine**

This is the third year that the Indiana Survey included questions about methamphetamine.

Compared to last year, the reported prevalence of methamphetamine either held steady or declined across grade levels (see Table 2). Reported lifetime, annual and monthly use of methamphetamine decreased among students in Grades 8 through 12, with the exception that monthly use by 10th-grade students did not change. Monthly use decreased for students in Grade 6. These results are consistent with a trend in decreased methamphetamine use seen since the item was introduced to the survey in 2005. Prevalence of lifetime methamphetamine use among Indiana's eighth, tenth, and twelfth grade youth are lower than available 2006 national prevalence rates reported in Monitoring the Future (Johnston, O'Malley, Bachman, & Schulenberg, 2005).

Table 2. *Methamphetamine use by youth in grades 6-12*

Grade	2005	2006	2007	Change 2006-07	Nation 2006**
6					
Lifetime	0.7	0.5	0.4	-0.1	
Annual	0.5	0.3	0.2	-0.1	
Monthly	0.3	0.2	0.1	-0.1*	
7					
Lifetime	1.2	1.2	1.0	-0.2	
Annual	0.9	0.8	0.7	-0.1	
Monthly	0.6	0.6	0.5	-0.1	
8					
Lifetime	2.4	2.0	1.6	-0.4*	2.7
Annual	1.7	1.4	1.2	-0.2*	1.8
Monthly	1.1	0.9	0.7	-0.2*	0.6
9					
Lifetime	3.5	3.0	2.2	-0.8*	
Annual	2.5	2.2	1.4	-0.8*	
Monthly	1.5	1.2	0.8	-0.4*	
10					
Lifetime	4.5	3.5	3.0	-0.5*	3.2
Annual	3.1	2.4	1.9	-0.5*	1.8
Monthly	1.6	1.2	1.0	-0.2	0.7
11					
Lifetime	4.6	4.2	3.3	-0.9*	
Annual	3.0	2.7	2.0	-0.7*	
Monthly	1.5	1.5	1.1	-0.4*	
12					
Lifetime	5.5	5.0	3.4	-1.6*	4.4
Annual	3.3	3.1	1.9	-1.2*	2.5
Monthly	1.7	1.5	1.0	-0.5*	0.9

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p<0.05$ ).

\*\* Johnston, O'Malley, Bachman, & Schulenberg, 2006.

## Other Drugs

In the 2007 Survey, the word "Adderall" was added to the item that had previously contained only "Ritalin," so that the item would measure the use of either Ritalin or Adderall. The prevalence rates associated with this item are higher this year than last, and this difference is statistically significant. However, because of the change in the Survey instrument, the rates for the item this year and last are not comparable.

Lifetime and annual use of non-prescribed narcotics increased for Grades 8 and 10 compared to last year. Non-prescribed narcotics include opium, morphine, codeine and OxyContin. The addition of OxyContin to the survey this year may have contributed to the observed increase this year. Although rates specific to OxyContin

use cannot be extracted from the Indiana Survey, the Monitoring the Future Survey indicated that 1.8% of eighth graders, 3.2% of tenth graders and 5.5% of twelfth graders reported past year use of OxyContin (Adlaf et al., 2006). The National Survey on Drug Use and Health has found that use of gateway drugs typically precede that of OxyContin, as well as the non-medical use of other prescription analgesics (Sees et al., 2005). Individuals who report using OxyContin typically have a history of multiple drug use including non-medical use of other prescription medications such as tranquilizers, sedatives and stimulants (Sees et al., 2005).

## Needle Injection of Drugs

Among students in Grade 12, the lifetime prevalence of injecting drugs with a needle (see Table 3) dropped to the level seen before 2006. Annual and monthly injection drug use remained the same. Injection drug use (IDU) is strongly correlated with the transmission of secondary infections such as human immunodeficiency virus (HIV), the virus that causes acquired immunodeficiency syndrome (AIDS). Injection drug use is also strongly associated with the spread of both hepatitis B and C, viral infections that lead to chronic disease and death. According to the Indiana State Department of Health, 11% of all HIV infections in 2005 were directly associated with IDU.

Table 3. *Injected drug use by Indiana 12th grade students: 2001-2007 (percentages)*

	2001	2002	2003	2004	2005	2006	2007	Change 2006-2007
Lifetime	1.8	1.4	1.7	1.8	1.7	2.2	1.7	-0.5 *
Annual	1.1	0.8	0.9	1.2	1.0	1.2	1.3	0.1
Monthly	0.6	0.6	0.6	0.7	0.6	0.9	0.9	0.0

\*Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

The Indiana Survey does not directly ask which drugs respondents injected. However, a sub-analysis of responses indicating any use of a needle to inject a drug suggests that the drugs most commonly injected by 12th-grade respondents were heroin, methamphetamine, and steroids (see Table 4). Among lifetime use of these drugs, heroin was more prevalent than methamphetamine and steroids. Also for lifetime use, methamphetamine use was more prevalent than steroid use. For annual and monthly use, heroin was more prevalent than steroids. These differences were statistically significant. The data suggest that the drug most often injected on a lifetime basis was heroin.

Table 4. *Injected drug use by Indiana 12th-grade students*

	N	%
Lifetime	300	
Heroin	186	62.0
Steroids	128	42.7
Methamphetamines	158	52.7
Annual	222	
Heroin	140	63.1
Steroids	107	48.2
Methamphetamines	125	56.3
Monthly	155	
Heroin	109	70.3
Steroids	88	56.8
Methamphetamines	95	61.3


## ***Regional Prevalence Rates***

This is the second year that this report presents the Indiana Survey results for Indiana's Family and Social Service Administration's sub-state planning regions (see Figure 1). Regional data provide more targeted information for local analyses. For the drugs studied, use prevalence rates varied considerably among regions and between any given region and the state. This summary does not attempt to capture the extent of this variation; however, a few examples are provided.

Compared to all respondents statewide, in the...

- Southeast Region, seventh through twelfth graders were more likely to report daily use of smokeless tobacco.
- East Region, seventh through eleventh graders were more likely to report daily use of cigarettes.
- Central Region, seventh through eleventh graders were less likely to report monthly use of alcohol.
- Northeast Region, sixth through tenth graders were less likely to report monthly use of cigarettes and alcohol.

The reasons for the regional variation are not addressed in this report. Communities may gain a better understanding of their own youth drug use by comparing their prevalence rates (in their local report) to the regional rates presented in this report.



**Figure 1.** Indiana's Family and Social Service Administration's substate planning regions.

## ***Risk and Protective Factors***

### **Age of First Use**

Research has shown that the younger a person is when she or he begins using alcohol, the more likely the person is to experience alcohol dependence and abuse (Grant, Stinson, & Harford, 2001; Warner & White, 2003). A recent study found that compared to persons who began drinking at age 21 or older, those who began drinking before age 14 were more likely to experience alcohol dependence later in life (R. W. Hingson, Heeren, & Winter, 2006).

In the Indiana Survey data, age of first use was measured by asking, "If you have ever used these drugs, at what age did you first use them?" From 1993 to 2006, responses to this question consisted of two-year intervals for ages 8 through 17 (i.e., 8-9 years, 10-11 years, etc.) and open-ended ranges for extreme responses ("7 or less years" and "18 or more years"). From 1993-2006, average age of first use was calculated using the midpoints of the intervals and the upper and lower endpoints for the extreme responses. In 2007, responses included open-ended ranges at the extremes ("7 or younger" and 18 or older") and one-year intervals representing discrete ages (i.e., "8 yrs," "9 yrs," etc.). Average age of first use for 2007 was calculated, as before, with the endpoints of the extreme responses, and with the

actual age of first use given for ages 8-17 years.

Over the years of Indiana Survey data, the average age of first use of gateway drugs has ranged consistently from 12 to 14 (see Figure 2). Reported initiation of the gateway drugs exhibits a nearly consistent pattern: cigarettes first, then alcohol, followed by marijuana. For marijuana and alcohol, changes from 2006 to 2007 in the average reported age of first use were statistically significant. Both increased by .2 percent.




Figure 2. Trends in average reported age of first use of cigarettes, alcohol, and marijuana.

## Perceived Risk of Harm

Prior research has demonstrated that perceived risk of harm in using drugs is negatively related to prevalence of use (Millstein & Halpern-Felsher, 2002; Wild, Hinson, Cunningham, & Bacchichiochi, 2001).

Table 5. *Glossary of prevalence of gateway drug use for all correlation analyses*

<b>Annual Use</b>	
Cigarettes	How often in the past year have you smoked cigarettes?
Alcohol	How many times in the last year have you used alcohol (beer, wine, wine coolers, liquor)?
Marijuana	How many times in the last year have you used marijuana (hashish or hash oil)?
<b>Monthly Use</b>	
Cigarettes	How often in the past month have you smoked cigarettes?
Alcohol	How many times in the last month (30 days) have you used alcohol (beer, wine, wine coolers, liquor)?
Marijuana	How many times in the last month (30 days) have you used marijuana (hashish or hash oil)?

To examine this relationship in the Indiana Survey data, respondents' perceived harm in using gateway drugs was correlated with reported monthly and annual use of gateway drugs. See Table 5 for definitions of annual and monthly use of gateway drugs. Perceived risk of harm was measured for the three gateway substances using this question: "How much do you think people risk harming themselves (physically or in other ways) if they... (a) smoke one or more packs of cigarettes per day, (b) smoke marijuana regularly, (c) have five or more drinks once or twice each weekend."

The results show that the higher the perceived risk of harm, the lower the reported annual and monthly levels of smoking cigarettes, smoking marijuana, or drinking alcohol (see Figures 3 and 4).




Figure 3. Perceived risk of harm and annual prevalence of gateway drugs.




Figure 4. Perceived risk of harm and monthly prevalence of gateway drugs.

## Perceptions of Peer Disapproval

Research has demonstrated that peer perceptions of disapproval exert an influence that lowers drug use (Butters, 2004; Chassin, Presson, & Sherman, 1984). To examine this relationship in the Indiana Survey data, respondents' perceptions of peer approval were correlated with reported use of gateway drugs. See Table 5 for definitions of annual and monthly use of gateway drugs.

Perceived peer approval was measured for three drugs of first use with the survey items: "How do you think your close friends feel (or would feel) about you doing each of the following things... (a) smoke one or more packs of cigarettes per day; (b) smoke marijuana regularly, (c) have five or more drinks once or twice each weekend."

The results illustrate that when reported perception of peer disapproval for a particular behavior is stronger, the reported level of the behavior is lower (see Figures 5 and 6).




Figure 5. Perceived peer approval and annual prevalence of gateway drugs.




Figure 6. Perceived peer approval and monthly prevalence of gateway drugs.

## Perceptions of Parents' and Guardians' Approval

The Indiana Survey data measured perceived parental approval for the use of three gateway drugs with the following items: "How do you think your parents/guardians feel (or would feel) about you doing each of the following things... (a) smoke one or more packs of cigarettes per day; (b) smoke marijuana regularly, (c) have five or more drinks once or twice each weekend ." The results appear in the prevalence tables.

Adolescents' perceptions of parental approval of substance use are positively related to the frequency of adolescents' use of alcohol, tobacco and marijuana (McMaster & Wintre, 1996; Lee et al., 2000; Jessor & Donovan, 1978; Barnes & Welte, 1986). To examine this relationship in the Indiana Survey data, respondents' perceptions of parents' and guardians' approval were correlated with reported use of gateway drugs in the previous month and previous year. Respondents tended to report engaging in these behaviors more often when they also reported stronger perceived parental approval for the behavior (see Figures 7 and 8). These findings are consistent with the research cited.




Figure 7. Perceived parental approval and annual prevalence of gateway drugs.




Figure 8. Perceived parental approval and monthly prevalence of gateway drugs.

## After-School Activities

Research has demonstrated that supervised after-school activities protect adolescents from using alcohol and other drugs (Grossman et al., 2002; Riggs & Greenberg, 2004). To determine if this relationship exists in the Indiana Survey data, prevalence of monthly and annual use of the gateway drugs was correlated with an index score measuring student participation in after-school activities. These activities included playing on a school sports team; participating in arts, music, or drama programs; participating in supervised activities at youth centers; and classes such as music, art, computer, dance, or sports lessons. The results illustrate that the more frequent the participation in after-school activities, the lower the prevalence of reported gateway drug use (see Figures 9 and 10). These findings support the value of after-school programming as a strategy to prevent or reduce substance use among adolescent populations.




Figure 9. After-school activities and annual prevalence of gateway drugs.




Figure 10. After-school activities and monthly prevalence of gateway drugs.

## Personal Safety

Research suggests that violent interactions and weapon carrying are positively related to prevalence of drug use (Ash & Kellermann, 1996; Dukarm, Byrd, Auinger, & Weitzman, 1996). To investigate whether these relationships exist in the Indiana Survey data, prevalence of annual use of gateway drugs was correlated with an index score measuring violent interactions. Violent interactions were measured by asking youth "In the past year, how often have you...? (a) been in a physical fight, (b) been in a serious argument involving shouting."

The findings indicate that the higher the frequency of involvement in violent interactions, the higher the reported prevalence of annual alcohol, cigarettes and marijuana use (see Figure 11). It is important to mention that correlation does not mean causation, but rather a relationship or association.

In addition, the Indiana prevalence data on annual use of gateway drugs (see Table 5) was correlated with an index score measuring weapon carrying to school. Weapon carrying was measured by asking youth: "In the past year, how often have you...? (a) carried a gun to school, (b) carried some other weapon to school."

The results illustrate that the higher the frequency of weapon carrying, the higher the reported prevalence of annual gateway drug use (see Figure 12). It is important to note that all Indiana Survey data was collected at a single point in time. It therefore cannot distinguish the order in which the reported behaviors occurred. Therefore, the data cannot suggest a causal relationship between behaviors such as weapon carrying and drug use. However, research indicates that other deviant behaviors, such as truancy, vandalism, poor grades, and fighting, occur before drug use

(Ensminger, Juon, & Fothergill, 2002; Poikolainen, 2002).




Figure 11. Violence and annual prevalence of gateway drugs.




Figure 12. Carrying weapons and annual prevalence of gateway drugs.

## Access to Alcohol

A public health strategy to reduce alcohol abuse by youth is to identify the primary sources of alcohol access and to target interventions to eliminate access at those points (Dent et al., 2005). Adults over the age of 21 are common sources of alcohol access for youth (Wagenaar et al, 1996; Jones-Webb et al., 1997). However, the sources tend to vary by age. Younger adolescents tend to get alcohol from their

homes and families, while older adolescents tend to get alcohol from friends and commercial sources (Harrison et al., 2000; Hearst et al., 2007).

The percentage of respondents who reported accessing alcohol from specific sources is shown for each grade in Table 6. The results are consistent with the research cited above. Family members were the primary source of alcohol for grades 6, 7, 8, and 9. The higher a respondent's grade, the more likely the respondent was to report either (1) having had someone buy it for him or her or (2) having received it from a person aged 21 or older. Respondents in higher grades reported obtaining alcohol from commercial outlets at higher rates than did those in lower grades.

These results suggest that youth drinking could be reduced if parents and family members better understood the risk of harm to youth that alcohol presents, as well as ways to prevent youth from accessing alcohol kept in the home. In addition, the results suggest that persons age 21 and older may need to be informed of the laws and penalties that apply to purchasing and/or otherwise supplying alcohol to minors.

*Table 6. Main sources of alcoholic beverages in the past month (percentages)*

	Grade						
	6	7	8	9	10	11	12
No Answer	11.1	10.1	9.7	11.3	11.6	12.3	12.7
No drink	81.9	77.9	71.3	66.1	60.2	57.7	51.9
Liquor stores/convenience stores/ supermarkets	0.1	0.1	0.2	0.2	0.7	0.9	1.6
Restaurants/bars/clubs	0.0	0.0	0.1	0.1	0.1	0.2	0.5
Public events	0.0	0.1	0.1	0.1	0.1	0.1	0.1
Had someone else buy it	0.3	0.9	2.6	4.3	8.0	10.1	13.0
Received from person 21 or older	0.9	1.5	3.0	4.0	5.5	6.5	8.8
Took it from a store	0.1	0.2	0.3	0.4	0.5	0.4	0.3
Family members	3.0	4.1	5.3	4.9	4.3	3.6	3.5
Other ways	2.5	5.0	7.6	8.7	9.0	8.1	7.6

## Reasons for Drinking Alcoholic Beverages

The percentage of respondents who reported specific reasons for drinking alcoholic beverages is shown in Table 7. Note that the majority of adolescents reported no consumption of alcoholic beverages. Of those who did drink alcoholic beverages, the leading reasons reported were "to experiment," "to have a good time with friends" and "because it tastes good."

Table 7. Most important reasons for drinking (percentages; multiple responses allowed)

	Grade						
	6	7	8	9	10	11	12
Did not drink alcoholic beverages	80.6	70.8	58.5	51.8	43.1	40.1	34.7
Because it tastes good	4.1	7.6	12.0	14.4	16.4	18.3	20.1
To have a good time with friends	1.0	2.4	4.8	8.2	12.1	14.9	18.2
To experiment	9.4	10.9	11.4	9.6	8.3	6.7	5.7
Because of boredom	1.0	1.8	3.2	3.9	5.2	5.1	5.4
To get away from my problems	0.7	1.0	1.8	2.2	2.8	2.8	3.2
Because of anger	0.9	1.6	2.5	30.	3.5	3.4	3.1
To relax or relieve tension	0.3	0.6	1.0	1.2	1.6	1.7	2.3
Because I am hooked	0.6	1.1	1.4	1.5	1.8	1.7	1.6
To get to sleep	0.3	0.5	0.7	0.8	1.1	1.2	1.4
To feel good or get high	0.3	0.5	0.9	1.0	1.3	1.3	1.3
To increase the effects of other drugs	0.1	0.2	0.4	0.7	1.0	1.0	1.1
To fit in with a group I like	0.4	0.5	0.7	0.7	0.9	0.9	0.9
To get through the day	0.2	0.4	0.6	0.6	0.7	0.5	0.6
To seek deeper insights or understanding	0.1	0.1	0.2	0.3	0.3	0.3	0.4
To decrease the effects of other drugs	0.1	-	0.1	0.1	0.1	0.1	0.1

In the Indiana Survey data, respondents were instructed to mark as many reasons for drinking alcoholic beverages as apply. Figure 13 shows that across grade levels, the more reasons a respondent gave for drinking, the more often they reported consuming alcoholic beverage in the previous year.




Figure 13. Number of reasons for drinking versus frequency of drinking in the past year.

To analyze the relationship between frequency of drinking and reasons given for drinking, respondents who reported any drinking in the previous year were divided into three groups: least-frequent drinkers, intermediate-frequency drinkers, and highest-frequency drinkers. Figure 14 below shows the relative frequency with which these three groups cited various reasons for their drinking.

Compared to the more frequent drinkers, the least frequent drinkers were markedly more likely to report drinking for experimentation, to fit into a group, or to gain deeper insights or understanding. The most frequent drinkers were the most likely to report drinking because they were hooked, to increase or reduce the effects of other drugs, and to get through the day. Intermediate-frequency drinkers selected most reasons at about as often as did the highest frequency drinkers. However, these two groups were distinguished from one another chiefly by their likelihood of reporting drinking to increase effects of other drugs and because of being hooked. The most frequent drinking group chose these options, respectively, about two and three times as often as the intermediate frequency drinkers.

Overall, the data suggest that infrequent drinkers tend to drink for social, recreational or coping reasons (e.g. anger, to relax, to get away from problems); whereas the most frequent drinkers do so to manage the physiological effects of alcohol or other drugs.




Figure 14. Annual alcohol use and reasons for drinking.

## ***Consequences of Use***


Research has shown that prevention and/or reduction of alcohol and drug use among student populations leads to a decrease in related problems (Hingson et al., 1996; Wodarski, 1988). Communities are better positioned to address substance use if they understand the relationship of that use to its consequences; these include, for example, poor performance on tests, missing school, and being hung over.

The percentage of respondents who reported specific consequences of alcohol or other drug use is shown for each grade in Table 8. The most commonly reported consequences across grades were having a hangover, nausea and vomiting, and riding in a car driven by someone who was "high" or had been using alcohol or drugs.

Table 8. Consequences of *alcohol, tobacco, and other drug use*

Consequences of Use	Grade						
	6	7	8	9	10	11	12
Had a hangover	4.0	9.6	18.1	23.3	30.3	33.9	38.2
Got nauseated or vomited	4.1	8.0	14.0	18.7	26.1	30.9	35.7
Had a memory loss	2.4	5.2	10.0	13.9	19.8	21.6	23.9
Performed poorly on a test or project	3.2	5.1	7.3	7.9	8.7	7.6	7.0
Missed school	3.2	4.4	6.1	6.8	7.9	7.3	7.9
Got into a fight or argument	5.4	7.9	11.8	12.9	16.0	15.9	16.2
Damaged property, pulled fire alarms, etc.	1.3	2.6	4.7	4.9	5.8	4.9	4.9
Been in trouble with your parents	4.7	7.2	10.5	12.0	14.8	14.9	14.4
Been in trouble with the police	2.0	3.3	5.3	5.9	6.8	6.7	6.6
Driven a car while under the influence	1.1	1.9	3.6	4.4	8.6	13.8	17.9
Ridden in a car driven by someone who was "high" or had been using alcohol or drugs	5.5	8.3	14.4	17.7	23.3	25.3	26.7
Forgot things you did while using alcohol or drugs	2.4	6.3	12.1	16.6	22.7	24.7	26.3
Your family or friends ever told you that you should cut down on your drinking or drug use	1.4	2.9	5.4	7.1	9.3	9.4	9.9
Used alcohol or drugs to relax, feel better about yourself, or fit in	3.2	6.3	11.0	14.2	17.9	19.2	20.3
Used alcohol or drugs while you were by yourself or alone	4.0	7.7	14.0	16.5	20.1	19.6	19.7

Figure 15 below illustrates the prevalence of number of consequences (0, 1 and 2+) among respondents who report various levels of drinking in the last year. Of those who reported drinking 1-5 times in the last year, roughly 40 percent reported zero consequences, and roughly 40 percent reported two or more consequences. In contrast, at least 80 percent of those who reported drinking 6-19 times and 20 plus times in the past year reported two or more consequence due to substance use.



*Figure 15. Frequency of alcohol use in the past year and number of reported consequences of ATOD use.*

## Gambling

As gambling options and venues in the state expand, monitoring the prevalence of gambling activities among children and adolescents is increasingly important for planning prevention efforts. This is the third year that the Indiana Survey has included gambling prevalence items. The Indiana Survey included questions about playing cards, betting on games, betting on sports, buying lottery tickets and gambling in a casino. Compared to last year, the proportion of respondents who reported any gambling behavior declined significantly in all grades except seventh (see Table 9). Seventh graders were more likely to have reported gambling than sixth graders, and eighth graders were more likely to have reported gambling than seventh graders. Twelfth graders were the most likely to report having gambled. Males

reported significantly more gambling than females (not shown).

Table 9. Any Gambling Behavior by Indiana Adolescents (percentages)

Grade	2005	2006	2007**	Change	
6 <sup>th</sup>	36.3	35.4	33.6	-1.8	*
7 <sup>th</sup>	40.7	39.1	38.6	-0.5	
8 <sup>th</sup>	47.4	45.3	43.3	-2.0	*
9 <sup>th</sup>	47.5	45.3	42.2	-3.1	*
10 <sup>th</sup>	48.0	45.2	42.5	-2.7	*
11 <sup>th</sup>	47.1	45.1	41.1	-4.0	*
12 <sup>th</sup>	55.1	55.0	50.8	-4.2	*

\* Statistically significant changes between 2006 and 2007 prevalence rates (p<.05).

\*\* 2007 data includes online gambling.

Problem gambling items were "During the last 12 months, have you ever felt...? (a) bad about the amount you bet, or about what happens when you bet money, (b) that you would like to stop betting money but didn't think you could." These items, used with the permission of the Minnesota Department of Education, help identify gambling problems (Johnson, Hamer, & Nora, 1998, and Johnson, et al., 1997). The findings indicate that compared to 2006, a smaller proportion of students reported feeling bad about the amount of money they bet (see Table 10) or about what happened when they bet money (see Table 11). The exceptions were Grade 6, where a larger proportions of students reported feeling bad, and Grade 7, which stayed the same. In addition, a greater proportion of students in Grades 6 and 7 reported they would like to stop betting money but did not think that they could.

Table 10. Trend in felt bad about the amount bet

Grade	2005	2006	2007**	Change 2006-2007	
6 <sup>th</sup>	5.6	5.2	5.6	0.4	*
7 <sup>th</sup>	5.5	5.0	5.1	0.1	
8 <sup>th</sup>	5.4	5.4	5.0	-0.4	*
9 <sup>th</sup>	5.4	4.6	4.1	-0.5	*
10 <sup>th</sup>	5.2	4.6	3.9	-0.7	*
11 <sup>th</sup>	4.8	4.1	3.4	-0.7	*
12 <sup>th</sup>	5.4	4.5	3.7	-0.8	*

\* Statistically significant changes between 2006 and 2007 prevalence rates (p<.05).

Table 11. *Trend in would like to stop betting but could not*

Grade	2005	2006	2007**	Change 2006-2007
6 <sup>th</sup>	4.9	4.5	4.9	0.4 *
7 <sup>th</sup>	3.7	3.6	4.0	0.4 *
8 <sup>th</sup>	3.4	3.2	3.5	0.3
9 <sup>th</sup>	2.6	2.3	2.6	0.3
10 <sup>th</sup>	2.2	1.9	2.1	0.2
11 <sup>th</sup>	1.8	1.6	1.7	0.1
12 <sup>th</sup>	1.7	1.6	1.5	-0.1

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

For the first time this year, the survey included the gambling prevalence item "played online for money." The results appear in Table 12.

 Table 12. *Reported any gambling online for money*

Grade	6	7	8	9	10	11	12
2007	3.3	4.3	4.6	4.2	4.3	3.8	3.8

## METHODOLOGY

The methodological history of the Indiana Survey can be read about in previous reports. The 2006 version can be found at [http://www.drugs.indiana.edu/data-survey\\_monograph.html](http://www.drugs.indiana.edu/data-survey_monograph.html).

### Sample

The sampling frame of the Indiana Survey is the universe of all Indiana school corporations and dioceses. The IPRC obtained from the Indiana Department of Education a list of names and addresses of all 2,774 public and nonpublic schools. In November 2006, recruitment materials were sent to all school superintendents, principals, and Safe and Drug-Free School Coordinators. The recruitment packet consisted of:

- an invitation letter.
- a memorandum of understanding (MOU).
- a statement on parental consent.
- an application form.
- a list of frequently asked questions.
- a stamped addressed return envelope.
- a blank survey form.

The corporations and dioceses had a 7-8 week window to respond by sending back an application and a signed MOU to the IPRC. In order to participate, a school or corporation was required to:

- identify a designated coordinator at each public school corporation or multiple coordinators if schools are scattered in several locations.
- describe its policy on parental consent.
- identify its spring break period.
- identify desired survey administration date.
- tell the number of students scheduled to participate in each grade level.

Schools were responsible for obtaining parental consent for students to participate in the Indiana Survey. Information about whether a particular school or corporation

participated in any year is strictly confidential. The MOU designated the results of school and corporation-level surveys as the property of the participating school or corporation. These results are released to third parties only with the explicit written permission of the school or corporation's survey coordinator or other appropriate authority.

## Participation

---

Since 1993, participation in the Indiana Survey has been open to all Indiana school corporations, public and nonpublic. The number of usable surveys, shown in Table 13, has fluctuated from year to year. While it is not possible to explain all of the fluctuation, the causes are likely to include:

- The exclusion from participation of 5th grade students since 1994.
- The participation of some schools on biennial and triennial bases.
- The arbitrary selection by some schools of the grade levels surveyed.
- Variation in the interpretation of parental consent requirements.
- The growing length of the *Indiana Survey* instrument.

Table 13. Number of Usable Surveys (1993-2007)

Year	Usable Surveys	Year	Usable Surveys	Year	Usable Surveys
1993	90,586	1998	44,232	2003	141,342
1994	81,732	1999	81,685	2004	91,577
1995	63,631	2000	72,523	2005	136,782
1996	36,586	2001	89,861	2006	120,914
1997	72,571	2002	77,068	2007	158,632

The rate of participation for school corporations, schools and students statewide and for each region is shown in Table 14. Rate of participation is the proportion of corporations, schools and students scheduled to participate in the survey out of the entire student population. There is wide variation in rates of participation across regions. The West Region had the highest rate of participation among corporations (64.1 percent), schools (21.5 percent) and students (38.8 percent) the North West Region the lowest rates, 31.7, 10.9 and 21.2 percent, respectively.

Table 14. Participation Rates for Students, Schools and Corporations by Region

	State	North West	North Central	North East	West	Central	East	South West	South East
<b>Students</b>									
Participants*	175,460	14,745	21,156	13,348	19,159	51,150	10,340	18,861	26,701
Total Population	558,429	69,527	88,922	60,824	56,695	170,069	34,715	41,396	60,846
Participation Rate <sup>1</sup>	31.4	21.2	23.8	22.0	33.8	30.1	29.8	45.6	43.9
<b>Schools</b>									
Participants	478	35	62	28	65	95	35	81	77
Total Population	3,203	321	614	479	303	732	204	248	319
Participation Rate <sup>2</sup>	14.9	10.9	10.1	5.8	21.5	13.0	17.2	32.7	24.1
<b>Corporations</b>									
Participants	152	13	20	10	25	28	13	18	25
Total Population	326	41	48	16	39	73	34	33	42
Participation Rate <sup>3</sup>	46.6	31.7	41.7	62.5	64.1	38.4	38.2	54.5	59.5

\* 252 cases were not included because the regional information was not available.

<sup>1</sup> The percentage of student participants out of the total population of 6-12th grade students.

<sup>2</sup> The percentage of schools that participated out of the total population of schools.

<sup>3</sup> The percentage of school corporations that participated out of the total population of corporations.

## Administration of the Survey

---

Two weeks before the designated survey date, the IPRC mailed to participant schools a packet containing:

- the survey forms.
- machine-readable answer sheets.
- written instructions for administering the survey.
- a twelve-minute survey administration training video produced by the IPRC.

The training video describes and demonstrates for school personnel each step of administering the survey to a group of students. The video and the written instructions were provided to improve the consistency of cross-site survey administration.

The written and video documentation included the following instructions for school personnel who administered the survey:

1. Inform students that:

- their participation in the survey is completely voluntary.
- the results are confidential.
- there are no penalties for deciding not to participate or for not responding to specific items.

2. Students will need about 30 minutes to complete the survey.

3. Remain seated while students fill out the survey.

4. Instruct the students that, when they complete the survey, they should place their survey forms into the single envelope, which the IPRC has provided and which is labeled with a school and classroom code.

No individually identifying information is collected, except gender, grade level, ethnic background and school code. Indiana Survey staff are available to provide support and answer questions through a toll-free 800 telephone number from Monday through Friday. Schools have a six-week window to administer the surveys between early March and mid-April and are requested to return them immediately to the IPRC.

## **Survey Instrument**

---

### ***Origin and Relevance of the Questions***

Year-to-year changes in the Indiana Survey instrument are documented in the reports corresponding to the years in which they occur. In general, survey items have been derived from a variety of sources for purposes of comparability with national and other state survey data. All items use multiple choice response options.

Alcohol and drug use prevalence items were developed based on those used in the MTF (Johnston, Bachman, & O'Malley, 1989), conducted by the University of Michigan. The Indiana Survey asked respondents about their use of twenty different types of drugs or drug classifications (See Indiana Survey instrument in appendix). In some instances, examples and/or slang descriptions appeared in parentheses next to the name of the drug or drug classification. The purpose of this was to elicit responses on the use of a single substance that might be a common ingredient in multiple products (i.e., alcohol in beer, wine, and liquor) or that might have multiple common names, some of them distributed geographically within the state. Other examples include "ecstasy," "XTC" and "X" for MDMA, and "meth," "crank," and "crystal" for methamphetamine.

The Indiana Survey also asks about the use of a needle or syringe to inject a drug. This question is meaningful because drugs can have multiple routes of administration, and drug injection can introduce secondary infection and disease (Center for Disease Control and Prevention, 2002; Miller et al., 2004).

Prevalence of alcohol, tobacco, and a variety of other drug use is a primary focus of the Indiana Survey. Prevalence refers to the rate of total cases of a condition in a population. For the Indiana Survey, prevalence is the percentage of reported drug use among child and adolescent populations over specified time frames. Definitions for the various prevalence-of-use measures used in this report are listed below.

*Lifetime prevalence:* the percentage of respondents who report using a particular drug at least once in their lifetime.

*Annual prevalence:* the percentage of respondents who report using a particular drug at least once during the year prior to the administration of the survey.

*Current use or monthly prevalence:* the percentage of respondents who report using a particular drug at least once in the 30 days prior to the administration of the survey.

Binge drinking prevalence: the percentage of respondents who report drinking at least five alcoholic drinks at a sitting in the two weeks prior to administration of the survey.

*Daily prevalence:* the percentage of respondents that report using a particular drug an average of once per day during the 30 day period prior to the administration of the survey. This rate is formulated based on responses of "20-40 times" or "more than 40 times" during the past month. This definition is also used by the MTF Study.

The perceived-risk of harm items have origins in both MTF and the National Household Survey on Drug Abuse (NHSDA). The perceived parental approval items have recently become performance measurement requirements for SAMHSA Drug-Free Communities Support Program grantees. The item set for age of first use is similar to that in the NHSDA. However, in the Indiana Survey these items are closed ended. The higher the perceived risk of harm the lower the reported level of drinking and other drug use.

The access to alcohol and the reasons for drinking items both have origins in the Youth Risk Behavior Survey. Reducing or eliminating identified points of access to alcohol is a primary public health approach to substance use prevention. Adolescents' reasons or expectancies for drinking alcoholic beverages can help explain different patterns of drinking (Kline, 1996; Bradizza, Reifman & Barnes, 1999).

The item sets about safety and security in the environment (for example, "How safe do you feel in school?", "How safe do you feel in the neighborhood?") are similar to items found in the National Longitudinal Study for Adolescent Health (Kingery, Coggesshall, & Alford, 1998). Items pertaining to personal safety such as having been in a physical fight and having been threatened by a weapon on school property are similar to those found in the YRBSS (Kingery, Coggesshall, & Alford, 1998). The questions that address carrying a gun to school or some other weapon have similarities to items used in the Teen Assessment Project (Atav & Spencer, 2002). To distinguish various purposes or motivations for carrying weapons, the Indiana Survey asks whether the respondent carried a gun or weapon for self defense, for hunting, or for some other reason.

The set of gambling items, added in 2005, originated from the Minnesota Student Survey (Stinchfield, Kushner, & Winters, 2005; Winters & Anderson, 2000).

## ***Reliability***

Reliability is an estimate of how consistently survey items measure the same concept each time they are administered under the same conditions and with the same

subjects. Internal reliability coefficients describe the consistency of results across sets of items. Internal reliability coefficients (i.e. Cronbach's alpha) based on the average correlation among item sets are shown in Table 15. Alpha coefficients range in value from 0 to 1. The higher the score is, the more reliable the item sets. Nunnaly (1978) has indicated 0.7 to be an acceptable reliability coefficient, and many researchers require at least 0.8 for a "good scale". Based on these criteria, the internal reliability of the Indiana Survey's item sets is good to excellent. This indicates that the item sets are consistent measures of the concepts in the Indiana Survey.

Table 15. *Internal Reliability of Item Sets*

Item Set	Question Wording	Number of Items	Cronbach's Alpha
Lifetime Use	Have you ever used...?	23	.902
Annual Use	How many times in the last year have you used...	23	.889
Use in the Past Month	How many times in the last month (30 days) have you used...	23	.844
Age of First Use	If you have ever used these drugs, at what age did you first use them?	22	.875
Consequences of Use	How many times have you experienced the following due to your drinking or drug use during the past year?	15	.924
Beliefs about Drugs	How much do you think people risk harming themselves (physically or in other ways) if they...	7	.895
Peer Perceptions	How do you think close friends feel (or would feel) about you doing each of the following things?	7	.935
Parental/guardian's Perceptions	How do you think your parents/guardians feel (or would feel) about you doing each of the following things?	7	.957
Neighborhood Environment	In your neighborhood, how safe do you feel?	5	.842
School Environment	In your school, how safe do you feel?	3	.884
Personal Safety Issues	In the past year, how often have you?	12	.798
After School Hours	There are about 180 school days per year. On how many of those days this year will you spend most of the after school hours...	14	.792
Gambling	During the last 12 months, how often have you done these activities?	5	.826

## Validity

The validity of a survey is the extent to which it measures what its designers intend it to measure. The Indiana Survey is intended to measure the reported prevalence of substance use, the consequences of use, perceptions of peer approval of substance use, perceptions of personal safety, prevalence of gambling, and other concepts. The Indiana Survey measures these concepts among children and adolescents in grades 6 through 12 located in schools across Indiana at a given point in time. The Indiana Survey uses a nonrandom sampling procedure, which presents a threat to the survey's external validity. A nonrandom procedure was used because a longstanding priority of the Indiana Survey has been to provide local level data to any and all schools or school corporations that request it. This nonrandom procedure may introduce bias, so the findings of the Indiana Survey should be generalized with caution.

To the extent that school districts have obtained information from every student or everyone in a particular grade level, the local results represent those community populations. In this respect, participating school districts can determine their own student response rate and decide how representative their survey data are for their local youth population. Formula 1.0 shows how the student response rate may be calculated from data in the local report provided to each participating school or school district.

**Formula 1.0**   Response Rate = Number of Respondents / Total Number of Students × 100

Where number of respondents is the students who completed and turned the survey in, and where total number of students is the number of students intended to receive the survey.

If the response rate is 90% or greater, then the district level results (not shown in this report) are likely to be representative of the local student population.

At the regional level, the ability to generalize the data is far less because many school districts and therefore many children and adolescents in the regional population did not participate in the Indiana Survey. As shown in Table 14, only 21.2 to 45.6 percent of students participated in the Indiana Survey across regions. The best approach to address this limitation is to assess the extent to which the samples of student respondents compare to the entire population of students in each region. This comparison of regional respondents and the population is shown by gender, ethnicity, race, and grade level in Table 16. Although this comparison does not provide a basis for generalizing results at the regional level, it provides guidance on where the data do and do not coincide proportionally with demographic subgroups in

the regional populations.

As shown in Table 16, the majority of samples either under or over-represent the proportions of students by gender, ethnicity, race, or grade level in a region. For example, in the Northwest region the proportion of males are underrepresented in the Indiana Survey sample and the proportion of Hispanics are over represented. This means that the sample has proportionally fewer males than the Northwest region as a whole, and this bias should be taken into consideration when interpreting the results of the Indiana Survey data. Likewise, the sample has proportionally more Hispanic students. Thus, this is a known limitation to applying the Indiana Survey results to non-Hispanic groups, which are under represented in the sample.

Table 16. Regional Comparison of Samples and Population by Demographics

	Indiana		Northwest		Northcentral		Northeast		West	
	Pop.	Sample	Pop.	Sample	Pop.	Sample	Pop.	Sample	Pop.	Sample
<b>Gender</b>										
Male	54.4	49.0 *	51.0	48.7 *	51.1	49.4 *	51.0	50.1 *	51.4	48.7 *
Female	51.8	51.0 *	49.0	51.3 *	48.9	50.6 *	49.0	49.9 *	48.6	51.3 *
<b>Ethnicity</b>										
Hispanics	5.2	7.0 *	9.6	14.9 *	7.6	13.6	5.0	7.0 *	3.7	5.9 *
Non-Hispanics	94.8	93.0 *	90.4	85.1 *	92.4	86.4 *	95.0	93.0 *	96.3	94.1 *
<b>Race</b>										
White	80.4	83.1 *	63.2	70.6 *	78.8	78.6	81.4	84.1 *	89.5	91.0
Black	11.8	7.7 *	23.1	14.2 *	9.2	7.9 *	9.8	6.6 *	2.7	1.8 *
Other#	7.9	9.2 *	13.7	15.2 *	12.0	13.5 *	8.9	9.3	7.8	7.2 *
<b>Grade</b>										
6	14.5	15.2	15.1	15.2	13.8	14.3	14.5	12.9 *	13.9	12.5 *
7	14.6	13.6 *	15.2	13.9 *	14.4	13.2 *	14.3	12.2 *	14.3	12.9 *
8	14.7	19.3 *	15.4	18.0 *	14.6	19.2 *	14.6	14.9	14.8	20.2 *
9	15.4	13.4 *	16.1	14.3 *	15.3	13.1 *	15.6	17.6 *	15.3	12.7 *
10	14.7	17.3 *	15.5	16.0	14.7	19.4 *	14.7	17.4 *	14.6	19.0 *
11	13.7	10.3 *	9.4	11.6 *	14.1	10.0 *	13.5	13.8	14.5	11.0 *
12	12.4	11.0 *	13.3	11.0 *	13.0	10.8 *	12.7	11.0 *	12.7	11.6 *
	Indiana		Central		East		Southwest		Southeast	
	Pop.	Sample	Pop.	Sample	Pop.	Sample	Pop.	Sample	Pop.	Sample
<b>Gender</b>										
Male	54.4	49.0 *	51.2	48.5 *	51.2	49.2 *	51.4	49.0 *	51.4	49.4 *
Female	51.8	51.0 *	48.8	51.5 *	48.8	50.8 *	48.6	51.0 *	48.6	50.6 *
<b>Ethnicity</b>										
Hispanics	5.2	7.0 *	4.4	6.1 *	1.5	3.7 *	1.5	3.2 *	1.6	4.2 *
Non-Hispanics	94.8	93.0 *	95.6	93.9 *	98.5	96.3 *	98.5	96.8 *	98.4	95.8 *
<b>Race</b>										
White	80.4	83.1 *	72.3	77.7 *	89.1	87.3 *	91.1	89.7 *	92.9	91.3 *
Black	11.8	7.7 *	18.8	12.5 *	5.5	6.2 *	4.4	4.6	2.5	2.5
Other#	7.9	9.2 *	8.9	9.8 *	5.4	6.5 *	4.4	5.7 *	4.5	6.2 *
<b>Grade</b>										
6	14.5	15.2	15.1	15.6 *	13.9	15.0 *	13.9	2.6 *	14.3	14.2
7	14.6	13.6 *	14.9	17.0 *	14.3	12.8 *	14.5	7.6 *	14.5	12.9 *
8	14.7	19.3 *	14.8	19.6 *	14.7	17.5 *	14.6	21.4 *	14.8	20.0 *
9	15.4	13.4 *	15.8	14.9 *	15.0	16.6 *	15.0	6.7 *	15.3	12.1 *
10	14.7	17.3 *	14.5	13.8 *	15.0	16.3 *	14.8	22.0 *	14.8	18.9 *
11	13.7	10.3 *	13.3	10.5 *	14.1	12.0 *	14.4	6.6 *	13.9	9.3 *
12	12.4	11.0 *	11.7	8.7 *	13.0	9.9 *	12.8	15.4 *	12.5	12.6

\* The sample is significantly different from the population ( $p < .05$ )

# Asian, American Indian, and multicultural students

The strength of the Indiana Survey is the ability to collect and report valid and reliable data for local school corporations. At the state level, the data has the same degree of reliability. However, according to sampling theory, the state-level data is less valid (Levy & Lemeshow, 2003). This tradeoff reflects the state's current priorities for serving the substance use prevention needs of local communities. At the same time, statewide results from the Indiana Survey have a longstanding and high degree of concurrence with state-level surveys that use probability sampling, such as the Youth Risk Behavior Survey and the Monitoring the Future survey. This high degree of concurrence is due to the large percentage of school corporations (46.6 percent in 2007) and students who participate in the Indiana Survey. Due to this level of agreement, it is reasonable to compare the statewide Indiana Survey data with national survey results. Furthermore, the results are consistent with findings from other prevention studies and this reinforces our confidence in the validity of Indiana Survey data at the state level.

## Data

The IPRC entered the Indiana Survey data in house using Pearson optical scanner equipment. The data were cleaned using multiple criteria. A survey was eliminated from the analysis for three reasons: failure of the respondent to report a grade level, failure of the respondent to answer a majority of the questions, or failure of the error check protocol. Table 17 shows the frequency and proportions of surveys eliminated for these reasons.

*Table 17: Reasons for Excluding Respondents from the Analysis*

	Frequency	Percent	Cumulative Percent
<b>Rejected Surveys</b>			
No grade level reported	3,894	2.2	2.2
Refused to answer majority of questions	6,909	3.9	6.1
Rejected due to failing error check protocol	6,277	3.6	9.7
Total Number of Usable Surveys	158,632	90.3	100.0
<b>Total</b>	<b>175,712</b>	<b>100.0</b>	<b>100.0</b>

The error check protocol eliminated surveys that met any one of four conditions:

- Responded provided a substantially inconsistent pattern of responses to prevalence questions (i.e., reported "never used" a particular drug in their lifetime and also reported using that drug in the past year or month)/

- Pattern of responses that was pharmacologically implausible (i.e., a combination of drugs and frequencies of use whose cumulative effect would be lethal).
- Reported a combination of age and grade that was implausible (i.e., a seven-year-old in the eleventh grade.)
- Respondent indicated that he or she did not respond truthfully at all. Measurement of this criterion is explained below.

The final item of the Indiana Survey asked students, "How truthfully have you answered these questions," with response options, (a) "not truthfully at all", (b) "somewhat truthfully", and (c) "completely truthfully". Those who responded "not truthfully at all" were eliminated from the analysis. The distribution of responses to this item is shown in Table 18. A significant degree of variation, from 1.1 to 1.9 percent (95% confidence interval), was found across regions for students who reported they did not answer truthfully at all. Further study is required to understand factors that contributed to this inter-regional variation.

Table 18. Response to Item on Truthful Completion of the Indiana Survey by Region

Regions	No Answer		Not Truthfully at All		Somewhat Truthfully		Completely Truthfully		Total
	Number	%	Number	%	Number	%	Number	%	
Northwest	2,262	15.3	247	1.7	1,011	6.9	11,225	76.1	14,745
North Central	3,141	14.8	317	1.5	1,307	6.2	16,391	77.5	21,156
Northeast	1,882	14.1	147	1.1	704	5.3	10,615	79.5	13,348
West	2,559	13.4	259	1.4	1,062	5.5	15,279	79.7	19,159
Central	7,884	15.4	767	1.5	3,028	5.9	39,471	77.2	51,150
East	1,709	16.5	150	1.5	610	5.9	7,871	76.1	10,340
Southwest	2,171	11.5	252	1.3	1,107	5.9	15,331	81.3	18,861
Southeast	3,656	13.7	323	1.2	1,555	5.8	21,167	79.3	26,701
Total	25,264	14.4	2,462	1.4	10,384	5.9	137,350	78.3	175,460

Note. 252 cases were not included because the regional information was not available.

## HOW TO INTERPRET THE RESULTS

### Margin of Error

The Indiana Survey has a margin of error plus or minus less than 1 percent with a 95 percent confidence interval. This means that if we selected a group of students 100 times and asked a given question from this survey each time, then in 95 of those instances, the percentage of students giving a particular answer would be within one percentage point of the percentage who gave that same answer in this year's results of the Indiana Survey. Margin of error and confidence intervals, however, have limited applicability to the Indiana Survey because respondents were chosen nonrandomly.

### 2006-2007 Changes

For the statewide data tables, values shown in the 'change' column refer to statistically significant observed differences in prevalence of use between years 2006 and 2007. According to Daniel (1991), statistical significance describes a mathematical measure of difference between groups. The difference is said to be statistically significant if it is greater than what might be expected to happen by chance alone. "P" is an estimate of the probability that the result has occurred by statistical accident. In this report, the p-value is set at 0.05 which means the differences or changes indicated has a 95 percent chance of being true. The plus and minus sign next to the change value indicates whether the 2007 percentage is an increase or decrease, respectively, from the previous year.

For the regional data tables, the symbols ▲ or ▼ are used to indicate statistically significant regional differences in prevalence of use compared to the entire state. The ▲ symbol indicates that the regional rate is higher than the state rate, and the ▼ symbol indicates that the regional rate is lower than the state rate.

### Generalizing Local Indiana Survey Data to Local Populations

The IPRC provides each participant school corporation with a local report of the Indiana Survey results from its district. This section describes the method for estimating how well local data applies to local populations. The IPRC strongly encourages school corporations to share the results of this analysis with citizens and organizations engaged in local surveillance and planning.

## The Response Rate

---

The response rate estimates how well the local results of the Indiana Survey describe all the youth in grade or grades, in a school or schools, or in an entire school corporation.

If the response rate is 90% or greater, then the results are likely to be representative of the population defined. If the response rate is lower than 90%, results should be interpreted cautiously. In this case, the results might not be representative of students who were absent the day the survey was administered or of those who did not turn in a survey.

In any case, results do not generalize to those whose responses the IPRC systematically excluded from tabulation (see the "Data" section, above, for details).

The IPRC is unable to calculate the response rate for individual localities because one of the components of the response rate depends on information not collected along with the survey. This information, the Total Number of Students, is explained below.

Two pieces of information are required to calculate the response rate: "the total number of students" and "the total number of respondents."

### ***The Total Number of Students***

The total number of students can only be determined by each school district.

The total number of students is the number of students that the school district intended to receive the survey. For instance, if the intention was to survey all 9th through 10th grade students in all schools then the total enrollment number in grades 9 through 10 would constitute the total number of students.

### ***Total Number of Respondents***

Local reports of the Indiana Survey issued to participating school corporations include a table similar to Table 19. The asterisk (\*) in Table 19 illustrates where to locate the total number of respondents.

Table 19. Sample Table from Local Report

	Frequency	Percent
No grade level reported		
Refused to answer majority of questions		
Rejected due to failing error check protocol		
Total Number of Usable Surveys		
Total	#,### *	100.0

## ***Calculating the Response Rate***

To determine the response rate for a locality, divide the Total Number of Respondents by the Total Number of Students and multiply the result by 100.

$$\text{Response Rate} = \left( \frac{\text{Number of Respondents}}{\text{Total Number of Students}} \right) \times 100$$

## **Identifying Significant Differences between Local and State Prevalence Rates**

In the local report, the first set of tables, titled "Significant Difference between Local and State Drug Use Rates, 2007," show where a school corporation had statistically significant higher or lower rates compared to the state. Positive numbers indicate that a school corporation had rates higher than those of the state. Negative numbers indicate that a school corporation had rates lower than those of the state. Dash marks appear where there were no statistically significant differences between the school corporation and the state.

If a school corporation had prevalence rates higher than those of the state on particular drugs, the community might want to consider targeting prevention efforts at those drugs.

## **Practical Significance**

The next set of tables in a local report shows the percentage of student participants who reported that they used specific drugs daily, during the past month, during the

past year and during their lifetime. These rates are shown for the school district over a five year period if such data are available and compared to the state and most recently available national rates.

Targeting prevention efforts at particular drugs might be appropriate even though local prevalence rates are not higher than those of the state. For instance, if 6th grade students in a corporation had a prevalence rate similar to that of the state for a specific drug, but the rate itself was high, prevention or treatment programs might be needed for 6th graders.

Key points for interpreting prevalence rate results of the Indiana Survey:

- Monthly prevalence is a general indicator of frequent or regular use of a drug.
- Annual prevalence is a general indicator of occasional or experimental use of a drug.
- "Prevalence" indicates any use of a drug within the specified time frame, regardless of its severity or frequency within that time frame (for example, any use of marijuana in the last year could be one time or two hundred times). Use monthly and annual prevalence rates in conjunction with detailed frequencies of use (for example, 1-5 times, 6-19 times, 20-40 times, more than 40 times). These are provided in tables in the back of a local report.

## **Interpreting Local Prevalence Rates of Gateway Drug Use**

---

- Cigarette smoking is the best predictor of future drug use.
- Binge drinking is a key indicator of future substance abuse.
- Alcohol use indicates a need for student assistance programs.
- Marijuana use is a key indicator of future drug use.
- Grades with the most rapid increases indicate appropriate timing of secondary prevention strategies.

If a school corporation has participated more than three times in the Indiana Survey, it has sufficient data to examine changes over time. If enrollment is relatively stable over time, trends and patterns may appear. A commonly observed pattern is the "cohort effect." For certain groups of youth, prevalence rates are inconsistent with the average use rates. These differences often persist over time. For instance, 8th grade students this year may have used particular drugs 10% more than 8th grade students two years ago. This pattern may continue from year to year until the students graduate.

## FUTURE OF THE INDIANA SURVEY

### **Collaboration with the Youth Risk Behavior Survey**

Beginning in 2009, and in odd-numbered years thereafter, the IPRC will collaborate with the Indiana Department of Education (IDOE) and Indiana State Department of Health (ISDH) on the administration of the Youth Risk Behavior Survey (YRBS). In 2009 and in subsequent years the Indiana Survey will continue to be administered to all schools that wish to participate. This collaboration will affect only the relatively few school classrooms that will be randomly selected to participate in the YRBS.

If a school is selected to participate in the YRBS, and that school also wishes to participate in the Indiana Survey, the IPRC will require the school to participate in the YRBS as a condition of receiving the Indiana Survey. Otherwise the Indiana Survey will be administered with no preconditions.

The YRBS already contains substance use prevalence items, and the IPRC collaboration will contribute additional substance use-related items beginning in 2009.

### **Addition of a random sample methodology**

Beginning in 2010, and in subsequent even-numbered years, the IPRC will implement a random sample method, in conjunction with administering the survey to all schools that wish to participate. The random sample selection approach will benefit local communities and the state by providing information that is scientifically representative at the regional and/or county level.

## NOTE ON THE REPORTING OF DETAILED DATA

Unless otherwise specified, detailed data is reported for the following substances or delivery mechanisms:

Cigarettes	Inhalants	Heroin
Cigars	Amphetamines	Steroids
Pipe	Methamphetamines	Injected drugs
Smokeless Tobacco	Ritalin or Adderall	MDMA
Alcohol	Tranquilizers	Rohypnol
Marijuana	Narcotics	GBH
Cocaine	LSD	Over the counter drugs
Crack	Psychedelics other than LSD	

The supplemental report available at <http://www.drugs.indiana.edu> includes:

### I. Statewide prevalence of use

- A. Frequency of use by gender (all, male, female), Grades 6-12 (tables)
- B. Distribution of age of first-time use of ATOD by gender (all/male/female) (tables)
- C. Monthly prevalence, Grade 12, Indiana and USA (graph)
- D. Prevalence of gateway drugs (graphs)
  - 1. Monthly, annual, and lifetime by grade (Grades 6-12)
  - 2. Monthly by grade and gender
- E. Trends in the prevalence of gateway drug use, Grades 6, 8, 10, and 12 (graphs)
  - 1. Annual and monthly
  - 2. Monthly, Grade 12, Indiana and USA Comparison
  - 3. Trend

### II. Regional prevalence of use

Regional prevalence rates by grade for lifetime, annual, and monthly use of all drugs, as well as daily and special use of selected drugs.

III. Risk and Protective Factors

- A. Perceptions, Grades 6-12
- B. After-school and extra-curricular activities, Grades 6-12
- C. Access to alcohol, Grades 6-12
- D. Reasons for drinking alcoholic beverages, Grades 6-12
- E. Consequences of use, Grades 6-12
- F. Safety, Grades 6-12

IV. Gambling, Grades 6-12

Annual prevalence rates

V. Appendix: Survey instrument

## References

- Ash, P., & Kellermann, A. (1996). Gun acquisition and use by juvenile offenders. *JAMA*, 275(22), 1754-1758.
- Atav, S., & Spencer, G. A. (2002). Health risk behaviors among adolescents attending rural, suburban, and urban schools: A comparative study. *Family & Community Health* 25(2), 53-64.
- Bailey, S. L. (1992). Adolescents' multisubstance use patterns: The role of heavy alcohol and cigarette use. *American Journal of Public Health*, 82(9), 1220-1224.
- Barnes, G.M., & Welte, J.W. (1986). Patterns and predictors of alcohol use among 7-12th grade students in New York state. *Journal of Studies on Alcohol*, 47(1), 53-62.
- Butters, J. E. (2004). The impact of peers and social disapproval on high-risk cannabis use: Gender differences and implications for drug education. *Drugs Education Prevention Policy*, 11(5), 381-390.
- Center for Disease Control and Prevention. (2002). Viral hepatitis and injection drug users. Retrieved July 14, 2006, from [http://www.cdc.gov/idiu/hepatitis/viral\\_hep\\_drug\\_use.htm](http://www.cdc.gov/idiu/hepatitis/viral_hep_drug_use.htm)
- Chassin, L., Presson, C. C., & Sherman, S. J. (1984). Cognitive and social influence factors in adolescent smoking cessation. *Addictive Behaviors*, 9(4), 383-390.
- Daniel, W. W. (1991). *Biostatistics: A foundation for analysis in the health sciences* (5th ed.). New York, NY: Wiley.
- Dent, C.W., Grube, J.W., & Biglan, A. (2005). Community level alcohol availability and enforcement of possession laws as predictors of youth drinking. *Preventive Medicine*, 40(3), 355-362.
- Donovan, J. E., & Jessor, R. (1983). Problem drinking and the dimension of involvement with drugs: A Guttman scalogram analysis of adolescent drug use. *American Journal of Public Health*, 73(5), 543-552.
- Dukarm, C. P., Byrd, R. S., Auinger, P., & Weitzman, M. (1996). Illicit substance use, gender, and the risk of violent behavior among adolescents. *Archives of Pediatrics & Adolescent Medicine*, 150(8), 797-801.
- Ensminger, M. E., Juon, H. S., & Fothergill, K. E. (2002). Childhood and adolescent antecedents of substance use in adulthood. *Addiction*, 97(7), 833-844.
- Fleming, R., Leventhal, H., Glynn, K., & Ershler, J. (1989). The role of cigarettes in the initiation and progression of early substance use. *Addictive Behaviors*, 14(3), 261-272.
- Golub, A., & Johnson, B. D. (2001). Variation in youthful risks of progression from alcohol and tobacco to marijuana and to hard drugs across generations. *American Journal of Public Health*, 91(2), 225-232.
- Grant, B. F., Stinson, F. S., & Harford, T. C. (2001). Age at onset of alcohol use and DSM-IV alcohol abuse and dependence: A 12-year follow-up. *Journal of Substance Abuse*, 13(4), 493-504.
- Grossman, J. B., Price, M. L., Fellerath, V., Jucovy, L. Z., Kotloff, L. J., Raley, R., et

- al. (2002). *Multiple choices after school: Findings from the extended-service schools initiative*. Philadelphia, PA. : Public/Private Ventures.
- Harrison, P.A., Fulkerson, J.A., & Park, E. (2000). The relative importance of social versus commercial sources in youth access to tobacco, alcohol, and other drugs. *Preventive Medicine*, 31(1), 39-48.
- Hearst, M.O., Fulkerson, J.A., Maldonado-Moline, M.M., Perry, C.L., & Komro, K.A. (2007). Who needs liquor stores when parents will do? The importance of social sources of alcohol among young urban teens. *Preventive Medicine*, 44, 471-476.
- Hingson, R., McGovern, T., Howland, J., Heeren, T., Winter, M., & Zakocs, R. (1996). Reducing alcohol-impaired driving in Massachusetts: The saving lives program. *American Journal of Public Health*, 86(6), p791-791.
- Hingson, R. W., Heeren, T., & Winter, M. R. (2006). Age at drinking onset and alcohol dependence: Age at onset, duration, and severity. *Archives of Pediatrics & Adolescent Medicine*, 160(7), 739-746.
- Johnson, E.E., Hamer, R., & Nora, R.M. (1998). The Lie/Bet questionnaire for screening pathological gamblers: A follow-up study. *Psychological Reports* 83, 1219-1224.
- Johnson, E.E., Hamer, R., Nora, R.M., Tan, B., Eisenstein, N., and Engelhart, C. (1997). The Lie/Bet Questionnaire for screening pathological gamblers. *Psychological Reports* 80, 83-88.
- Johnston, L. D., Bachman, J. G., & O'Malley, P. M. (1989). *Drug use, drinking, and smoking: National survey results from high school, college, and young adult populations, 1975-1989*. Rockville, MD: National Institute on Drug Abuse.
- Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2005). Teen drug use down but progress halts among youngest teens. Retrieved December 19, 2005, from [www.monitoringthefuture.org](http://www.monitoringthefuture.org)
- Johnston, L. D., O'Malley, P. M., Bachman, J. G. & Schulenberg, J. E. (2006). Teen drug use continues down in 2006, particularly among older teens; but use of prescription-type drugs remains high. Retrieved December 31, 2006, from <http://www.monitoringthefuture.org>.
- Jones-Webb, R., Toomey, T., Miner, K., Wagenaar, AC., Wolfson, M., & Poon, R. (1997). Why and in what context adolescents obtain alcohol from adults: a pilot study. *Substance Use and Misuse*, 32(2), 219-228.
- Jessor, R., & Donovan, J.E. (1978). Drinking, problem drinking, and illicit drug use among students: A psychosocial study of a nationwide sample. Report.
- Kandel, D., & Yamaguchi, K. (1993). From beer to crack: Developmental patterns of drug involvement. *American Journal of Public Health*, 83(6), 851-855.
- Kingery, P. M., Coggeshall, M. B., & Alford, A. A. (1998). Violence at school: Recent evidence from four national surveys. *Psychology in the Schools*, 35(3), 247-258.
- Knight, J.R., Sherritt, L., Harris, S.K., Gates, E.C., & Chang, G. (2003). Validity of brief alcohol screening tests among adolescents: A comparison of the AUDIT, POSIT, CAGE, and CRAFFT. *Alcoholism Clinical and*

- Experimental Research*, 27(1), 67-73.
- Lee, C.Y., Eyre, S.L., & Millstein, S.G. (2000). Gender and ethnic differences in adolescent alcohol use; Problem behavior theory approach. *Journal of Adolescent Health*, 26(2), 143.
- Levy, P. S., & Lemeshow, S. (2003). *Sampling of populations: Methods and applications* (3rd ed.). New York, NY: John Wiley & Sons.
- McMaster, L.E., & Wintre, M.G. (1996). Relations between perceived parental reciprocity, perceived parental approval, and adolescent substance use. *Journal of Adolescent Research*, 11(4), 440-460.
- Miller, C. L., Wood, E., Spittal, P. M., Li, K., Frankish, J. C., Braitstein, P., et al. (2004). The future face of coinfection: Prevalence and incidence of HIV and hepatitis C virus coinfection among young injection drug users. *Journal of Acquired Immune Deficiency Syndromes*, 36(2), 743-749.
- Millstein, S. G., & Halpern-Felsher, B. L. (2002). Perceptions of risk and vulnerability. *The Journal of Adolescent Health*, 31(S), 10-27.
- Nunnally, J. C. (1978). *Psychometric theory* (2nd ed.). New York, NY: McGraw Hill.
- Poikolainen, K. (2002). Antecedents of substance use in adolescence. *Current Opinion in Psychiatry*, 15(3), 241-245.
- Riggs, N. R., & Greenberg, M. T. (2004). After-school youth development programs: A developmental-ecological model of current research. *Clinical Child & Family Psychology Review*, 7(3), 177-190.
- Stinchfield, R., Kushner, M. G., & Winters, K. C. (2005). Alcohol use and prior substance abuse treatment in relation to gambling problem severity and gambling treatment outcome. *Journal of Gambling Studies*, 21(3), 273-297.
- Wagenaar, A.C., & Toomey, T.L. (1996). Sources of alcohol for underage drinkers. *Journal of Studies on Alcohol*, 57(3), 325-333.
- Warner, L. A., & White, H. R. (2003). Longitudinal effects of age at onset and first drinking situations on problem drinking. *Substance Use & Misuse*, 38(14), 1983-2016.
- Wild, T. C., Hinson, R., Cunningham, J., & Bacchiori, J. (2001). Perceived vulnerability to alcohol-related harm in young adults : Independent effects of risky alcohol use and drinking motives. *Experimental and Clinical Psychopharmacology*, 9(1), 117-125.
- Winters, K. C., & Anderson, N. (2000). Gambling involvement and drug use among adolescents. *Journal Of Gambling Studies* 16(2-3), 175-198.
- Wodarski, J. S. (1988). Teams-Games-Tournaments: Teaching adolescents about alcohol and driving. *Journal of Alcohol and Drug Education*, 33( 3), 46-57.

## II. STATEWIDE DATA

<b>Statewide Prevalence Rates by Grade: Lifetime, Annual, Monthly, Daily and Special Use</b>	52
.....	52
• Grade 6 .....	52
• Grade 7 .....	56
• Grade 8 .....	60
• Grade 9 .....	64
• Grade 10 .....	68
• Grade 11 .....	72
• Grade 12 .....	76
<b>Mean Age of First Time Use of Alcohol, Tobacco, and Other Drugs .....</b>	80

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 6th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
Cigarettes	29.2	28.8	27.6	27.7	27.3	26.3	23.3	21.8	20.6	17.7	17.7	14.6	11.9	11.1	-0.8	
Cigars	--	--	--	--	--	9.7	8.4	7.7	10.5	6.9	6.8	7.0	6.0	5.4	4.8	-0.6
Pipe	--	--	--	--	--	3.2	3.1	2.9	--	2.1	2.2	2.4	2.0	1.8	1.3	-0.5
Smokeless Tobacco	8.5	9.0	7.7	7.1	6.7	4.6	3.7	4.0	3.5	2.9	3.2	3.0	2.9	2.7	2.4	
Alcohol	37.8	36.1	32.4	29.3	32.3	25.7	26.9	25.8	29.6	29.4	29.4	29.7	26.1	25.2	22.5	-2.7
Marijuana	2.2	2.9	3.6	5.1	4.7	5.0	4.2	3.8	4.5	4.8	4.6	4.5	3.9	2.9	3.0	
Cocaine	0.1	0.5	0.6	0.8	0.8	0.9	0.7	0.7	0.8	0.7	0.8	0.7	0.8	0.6	0.5	
Crack	0.1	0.6	0.6	1.0	0.9	1.0	0.9	0.8	1.1	1.0	1.0	1.0	1.1	0.8	0.7	
Inhalants	7.8	8.4	8.1	8.1	10.1	7.9	7.3	6.4	6.9	6.1	7.2	8.1	7.3	6.5	6.2	
Amphetamines	1.5	1.6	1.3	1.5	1.5	1.3	0.9	0.9	1.0	0.8	0.7	0.7	0.6	0.5	0.4	-0.1
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.7	0.5	0.4
Ritalin	--	--	--	--	--	1.4	1.0	1.2	1.3	1.1	1.1	1.1	1.1	0.9	1.0	
Tranquilizers	4.3	4.4	4.0	4.2	5.1	4.7	4.1	3.7	4.0	4.3	4.4	4.4	4.1	3.9	4.0	
Narcotics	1.7	1.4	1.2	1.3	1.5	1.4	1.0	0.9	1.1	1.0	1.0	1.2	0.9	0.8	0.8	
Psychedelics#	0.1	1.0	0.8	1.4	1.5	1.5	1.1	0.9	--	1.0	0.9	1.0	0.8	0.6	0.5	
LSD	--	--	--	--	--	--	--	--	--	0.8	0.7	0.7	0.5	0.5	0.3	
Others	--	--	--	--	--	--	--	--	--	0.5	0.5	0.6	0.5	0.4	0.3	
Heroin	0.1	0.5	0.4	0.8	0.7	0.9	0.6	0.6	0.8	0.7	0.7	0.7	0.6	0.5	0.4	
Steroids	1.2	1.5	1.0	1.4	1.4	1.3	1.4	1.2	1.1	0.9	1.0	1.1	0.9	0.8	0.7	
Injected drugs	--	--	--	--	--	--	--	--	0.9	0.8	0.8	1.0	1.0	0.9	0.8	
MDMA	--	--	--	--	--	--	--	--	--	0.7	0.6	0.7	0.5	0.4	0.3	-0.1
Rohypnol	--	--	--	--	--	--	--	--	--	0.4	0.4	0.4	0.4	0.4	0.2	-0.2
GHB	--	--	--	--	--	--	--	--	--	0.3	0.3	0.3	0.3	0.2	-0.1	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	4.7	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs by Indiana 6th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
Cigarettes	19.0	19.5	18.3	18.4	18.3	16.4	14.2	12.5	11.2	9.4	9.5	9.5	7.9	6.2	5.3	-0.9
Cigars	--	--	--	--	--	7.4	6.3	5.5	7.0	4.5	4.3	4.6	4.0	3.4	3.1	
Pipe	--	--	--	--	--	2.3	1.7	1.7	--	1.4	1.5	1.6	1.3	1.1	1.0	
Smokeless Tobacco	6.4	6.8	5.8	5.4	4.9	3.2	2.6	2.6	2.5	2.1	2.2	2.2	2.0	1.9	1.7	
Alcohol	27.7	27.3	23.2	21.7	23.7	19.3	21.0	20.3	21.4	20.4	20.7	20.7	17.6	17.3	15.3	-2.0
Marijuana	1.7	2.5	3.1	4.4	4.0	4.2	3.6	3.2	3.6	3.6	3.5	3.7	3.0	2.3	2.3	
Cocaine	0.1	0.4	0.4	0.8	0.7	0.7	0.5	0.6	0.7	0.5	0.6	0.6	0.5	0.4	0.4	
Crack	0.1	0.4	0.3	0.9	0.7	0.8	0.7	0.6	0.8	0.6	0.8	0.8	0.7	0.5	0.5	
Inhalants	6.1	6.2	5.8	6.4	7.9	6.0	5.4	4.5	5.2	4.1	5.2	5.9	5.1	4.5	4.1	-0.4
Amphetamines	1.2	1.3	1.0	1.4	1.3	1.1	0.7	0.7	0.7	0.6	0.6	0.6	0.4	0.4	0.3	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	0.5	0.3	0.2
Ritalin	--	--	--	--	--	1.2	0.8	0.9	0.9	0.7	0.7	0.8	0.7	0.5	0.7	0.2
Tranquilizers	3.5	3.5	3.0	3.2	4.1	3.8	3.2	2.8	3.0	3.0	3.1	3.2	3.0	2.7	2.8	
Narcotics	1.3	1.0	0.9	1.0	1.3	1.0	0.6	0.7	0.9	0.6	0.7	0.8	0.6	0.5	0.5	
Psychedelics#	0.1	0.7	0.7	1.2	1.2	1.3	0.8	0.7	--	0.6	0.6	0.7	0.5	0.4	0.4	
LSD	--	--	--	--	--	--	--	--	--	0.5	0.5	0.5	0.3	0.3	0.2	
Others	--	--	--	--	--	--	--	--	--	0.3	0.3	0.4	0.3	0.2	0.3	
Heroin	0.1	0.4	0.2	0.6	0.5	0.7	0.4	0.4	0.6	0.4	0.5	0.5	0.4	0.3	0.2	
Steroids	0.1	1.0	0.6	1.0	1.0	1.1	0.8	0.8	0.8	0.6	0.7	0.8	0.6	0.5	0.5	
Injected drugs	--	--	--	--	--	--	--	--	0.6	0.5	0.7	0.7	0.7	0.5	0.6	
MDMA	--	--	--	--	--	--	--	--	--	0.5	0.5	0.5	0.4	0.3	0.2	-0.1
Rohypnol	--	--	--	--	--	--	--	--	--	0.2	0.3	0.3	0.2	0.1	0.1	-0.1
GHB	--	--	--	--	--	--	--	--	--	0.2	0.2	0.2	0.1	0.1		
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	3.8	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 6th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
Cigarettes	8.5	10.0	9.3	9.7	9.5	8.2	7.0	5.7	6.0	4.8	5.1	5.4	4.2	3.3	2.8	-0.5
Cigars	--	--	--	--	--	3.9	3.3	2.9	3.8	2.4	2.5	2.5	2.2	1.8	1.6	
Pipe	--	--	--	--	--	1.5	1.0	0.8	--	0.8	0.9	0.8	0.7	0.7	0.5	
Smokeless Tobacco	3.5	4.0	3.3	3.1	3.1	1.9	1.4	1.4	1.4	1.2	1.4	1.3	1.1	1.1	0.9	
Alcohol	13.6	13.7	11.4	10.7	11.5	9.5	9.9	9.2	9.5	9.5	9.9	9.8	8.1	7.6	6.9	-0.7
Marijuana	1.2	1.7	1.9	3.0	2.7	2.2	1.8	2.4	2.2	2.3	2.5	1.8	1.4	1.4	1.4	
Cocaine	0.1	0.3	0.6	0.5	0.5	0.3	0.4	0.4	0.3	0.4	0.3	0.4	0.4	0.4	0.2	0.2
Crack	0.1	0.3	0.3	0.6	0.5	0.6	0.3	0.4	0.4	0.4	0.5	0.5	0.5	0.3	0.3	
Inhalants	3.8	4.1	3.6	3.8	4.9	3.6	3.4	2.7	3.1	2.5	3.2	3.6	3.1	2.9	2.3	-0.6
Amphetamines	0.1	0.9	0.6	1.0	0.9	0.6	0.5	0.3	0.4	0.4	0.4	0.3	0.2	0.2	0.2	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	
Ritalin	--	--	--	--	--	0.7	0.5	0.5	0.6	0.4	0.4	0.5	0.4	0.3	0.4	0.1
Tranquilizers	2.0	2.2	1.6	1.9	2.4	2.0	1.8	1.6	1.7	1.7	1.8	1.9	1.7	1.5	1.5	
Narcotics	0.1	0.5	0.6	0.6	0.9	0.7	0.4	0.5	0.5	0.4	0.4	0.5	0.3	0.3	0.3	
Psychedelics#	0.1	0.6	0.5	0.8	0.9	0.8	0.5	0.4	--	0.4	0.4	0.4	0.3	0.2	0.2	
LSD	--	--	--	--	--	--	--	--	--	0.4	0.3	0.3	0.3	0.2	0.1	
Others	--	--	--	--	--	--	--	--	--	0.2	0.2	0.2	0.2	0.1	0.1	
Heroin	0.1	0.3	0.2	0.5	0.4	0.4	0.3	0.3	0.4	0.3	0.4	0.3	0.3	0.2	0.2	
Steroids	0.1	0.6	0.4	0.7	0.7	0.6	0.5	0.4	0.5	0.4	0.4	0.5	0.4	0.3	0.3	
Injected drugs	--	--	--	--	--	--	--	--	0.4	0.3	0.4	0.4	0.5	0.3	0.3	
MDMA	--	--	--	--	--	--	--	--	--	0.4	0.3	0.3	0.3	0.1	0.1	
Rohypnol	--	--	--	--	--	--	--	--	--	0.2	0.2	0.1	0.2	0.1	0.1	
GHB	--	--	--	--	--	--	--	--	--	0.1	0.1	0.1	0.1	0.1	0.1	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	2.4	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th Grade Students: 1993-2007**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
<b>CIGARETTES</b>																
Daily Use	3.4	4.3	4.0	4.2	4.3	3.7	3.1	2.3	2.6	1.9	2.2	2.1	1.9	1.4	1.0	-0.4
1/2+ Pack per Day	1.5	2.0	1.9	2.1	2.1	1.7	1.4	1.0	1.2	0.9	0.8	0.9	1.0	0.7	0.4	-0.3
CIGARS	--	--	--	--	--	0.6	0.5	0.3	0.6	0.2	0.3	0.4	0.3	0.2	0.2	
SMOKELESS TOBACCO	0.1	0.6	0.4	0.5	0.5	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	
<b>ALCOHOL</b>																
Daily Use	0.1	0.4	0.5	0.4	0.6	0.5	0.4	0.4	0.5	0.5	0.6	0.4	0.5	0.3	0.3	
Binge Drinking	8.0	8.1	7.2	7.6	7.6	8.7	8.2	7.9	6.7	6.6	5.3	5.9	4.5	4.2	4.9	0.7
MARIJUANA	0.1	0.1	0.3	0.4	0.3	0.4	0.4	0.2	0.4	0.4	0.5	0.4	0.4	0.3	0.2	-0.1

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates (p&lt;.05).

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 7th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
Cigarettes	38.6	40.1	39.8	40.2	38.5	35.1	34.0	31.2	29.1	26.3	23.4	24.6	22.0	19.7	17.9	-1.8
Cigars	--	--	--	--	--	15.1	14.3	12.3	14.9	11.5	9.8	10.6	9.6	8.3	8.6	
Pipe	--	--	--	--	--	5.1	5.0	5.0	--	3.6	3.6	3.8	3.2	2.9	3.0	
Smokeless Tobacco	13.9	14.9	14.6	12.7	11.5	8.1	7.3	5.9	5.2	5.3	4.8	6.1	4.8	5.0	4.3	-0.7
Alcohol	48.7	47.4	44.4	44.0	42.4	36.2	37.5	36.5	39.5	39.4	37.7	38.3	34.0	33.3	32.7	
Marijuana	5.3	7.7	9.5	12.8	10.2	9.9	9.4	8.8	9.8	9.3	8.7	8.7	8.0	6.8	7.7	0.9
Cocaine	1.0	1.2	1.3	2.1	1.6	1.6	1.2	1.2	1.5	1.5	1.6	1.6	1.3	1.5	1.2	-0.3
Crack	0.1	1.3	1.3	1.9	1.4	1.7	1.6	1.4	1.5	1.8	1.6	1.9	1.6	1.5	1.3	-0.2
Inhalants	9.8	12.0	12.3	13.5	12.9	10.6	8.8	7.9	7.8	7.6	8.8	10.3	8.9	9.0	7.9	-1.1
Amphetamines	3.9	4.5	4.6	5.7	3.5	3.0	2.5	2.4	2.1	2.4	2.1	2.0	1.5	1.4	1.2	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	1.2	1.2	1.0
Ritalin	--	--	--	--	--	2.5	2.1	2.4	1.9	2.0	1.9	1.9	1.9	1.8	2.2	0.4
Tranquilizers	6.5	6.8	6.7	8.5	7.4	7.1	6.2	6.4	6.0	6.8	6.5	7.2	6.0	6.0	5.6	
Narcotics	3.0	2.9	2.8	3.6	3.4	2.9	2.4	2.3	2.1	2.4	2.2	2.4	2.0	2.0	2.2	
Psychedelics#	1.8	2.2	2.4	3.7	3.0	3.1	2.5	2.4	--	2.1	2.0	1.8	1.6	1.5	1.2	-0.3
LSD	--	--	--	--	--	--	--	--	--	1.7	1.6	1.5	1.2	1.1	1.0	
Others	--	--	--	--	--	--	--	--	--	0.9	0.9	0.8	0.8	0.8	0.6	-0.2
Heroin	1.0	1.2	1.1	1.2	1.3	1.1	1.0	0.9	1.0	1.1	1.2	1.0	1.1	0.9	-0.2	
Steroids	2.2	2.4	1.7	2.0	1.8	1.6	2.1	1.5	1.3	1.6	1.5	1.6	1.3	1.2	1.2	
Injected drugs	--	--	--	--	--	--	--	--	0.9	1.3	1.3	1.5	1.3	1.3	1.4	
MDMA	--	--	--	--	--	--	--	--	--	1.8	1.8	1.5	1.2	1.2	1.0	
Rohypnol	--	--	--	--	--	--	--	--	--	0.9	0.7	0.7	0.6	0.8	0.6	
GHB	--	--	--	--	--	--	--	--	--	0.5	0.5	0.4	0.4	0.5	0.4	-0.1
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	6.5	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs by Indiana 7th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
Cigarettes	28.4	29.7	29.8	31.4	29.0	25.2	22.7	20.0	18.1	15.7	13.9	15.2	13.0	11.7	10.8	-0.9
Cigars	--	--	--	--	--	12.6	11.0	9.5	10.5	7.9	6.4	7.4	6.6	5.8	6.2	0.4
Pipe	--	--	--	--	--	3.7	3.2	2.9	--	2.6	2.7	2.8	2.1	2.1	2.6	0.5
Smokeless Tobacco	10.9	11.7	11.9	10.2	8.8	6.4	5.1	3.9	3.8	3.9	3.6	4.5	3.5	3.9	3.3	-0.6
Alcohol	39.8	38.7	36.1	36.9	34.5	30.3	31.7	30.8	31.6	30.1	29.2	29.8	25.7	25.5	24.8	
Marijuana	4.5	6.7	8.6	11.5	9.1	8.6	8.1	7.9	8.3	7.8	7.1	7.2	6.3	5.6	6.3	0.7
Cocaine	0.1	1.1	1.1	1.8	1.4	1.3	1.3	1.1	1.0	1.2	1.3	1.3	1.0	1.2	1.0	
Crack	0.1	0.9	1.0	1.6	1.2	1.3	1.2	1.3	1.2	1.1	1.3	1.4	1.1	1.2	0.9	-0.3
Inhalants	7.6	9.6	9.3	11.4	10.3	7.8	6.4	5.7	5.6	4.9	6.2	7.6	6.4	6.3	5.4	-0.9
Amphetamines	3.3	3.9	4.0	5.1	3.2	2.4	2.0	2.0	1.8	1.8	1.5	1.5	1.1	1.1	1.0	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	0.9	0.8	0.7
Ritalin	--	--	--	--	--	2.0	1.6	2.0	1.5	1.4	1.4	1.3	1.3	1.3	1.6	0.3
Tranquilizers	5.2	5.6	5.4	7.2	6.2	5.9	5.0	5.2	5.0	5.1	4.9	5.5	4.5	4.6	4.2	
Narcotics	2.4	2.2	2.2	3.1	2.7	2.2	1.9	1.8	1.7	1.7	1.5	1.7	1.4	1.4	1.6	
Psychedelics#	1.5	1.7	2.2	3.2	2.6	2.6	2.1	2.1	--	1.5	1.5	1.4	1.2	1.1	1.0	
LSD	--	--	--	--	--	--	--	--	--	1.3	1.2	1.1	0.9	0.8	0.7	
Others	--	--	--	--	--	--	--	--	--	0.6	0.6	0.6	0.5	0.6	0.6	
Heroin	0.1	0.9	0.9	1.3	1.0	0.9	0.8	0.7	0.8	0.9	0.9	1.0	0.7	0.8	0.7	
Steroids	1.6	1.8	1.2	1.5	1.4	1.1	1.5	1.1	1.0	1.2	1.1	1.3	0.9	0.9	0.9	
Injected drugs	--	--	--	--	--	--	--	--	0.7	1.0	0.9	1.0	0.9	1.0	1.0	
MDMA	--	--	--	--	--	--	--	--	--	1.4	1.4	1.1	0.9	0.9	0.8	
Rohypnol	--	--	--	--	--	--	--	--	--	0.7	0.5	0.5	0.4	0.6	0.5	
GHB	--	--	--	--	--	--	--	--	--	0.4	0.3	0.3	0.4	0.3		
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	5.1	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 7th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
Cigarettes	14.5	16.9	17.7	19.0	16.3	14.0	12.3	11.0	9.9	9.1	8.1	8.9	7.5	6.7	6.2	
Cigars	--	--	--	--	--	6.9	5.8	5.3	6.0	4.6	3.7	4.5	3.7	3.2	3.6	0.4
Pipe	--	--	--	--	--	2.4	1.8	2.1	--	1.5	1.5	1.6	1.2	1.2	1.5	0.3
Smokeless Tobacco	6.7	7.5	7.1	6.7	5.2	3.9	2.7	2.2	2.1	2.3	2.1	2.8	2.2	2.3	2.0	-0.3
Alcohol	21.8	21.5	19.6	21.3	18.4	16.8	16.5	16.1	15.2	14.7	15.3	12.7	12.9	12.2	-0.7	
Marijuana	2.7	4.8	5.4	7.9	5.7	5.5	5.1	5.2	5.2	4.7	4.6	4.5	4.1	3.6	3.9	
Cocaine	0.1	0.8	0.7	1.3	1.0	0.9	0.7	0.7	0.6	0.8	0.9	0.8	0.7	0.8	0.7	
Crack	0.1	0.7	0.7	1.2	0.9	0.9	0.8	0.7	0.7	0.7	0.9	0.9	0.8	0.8	0.6	-0.2
Inhalants	4.7	6.1	5.6	7.0	6.6	4.8	3.7	3.3	3.1	3.0	3.7	4.8	3.7	3.6	3.1	-0.5
Amphetamines	2.2	2.7	2.4	3.3	2.3	1.7	1.2	1.2	1.1	1.1	1.0	1.1	0.7	0.8	0.6	-0.2
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.6	0.6	0.5
Ritalin	--	--	--	--	--	1.4	1.0	1.2	0.8	0.8	0.9	0.8	0.8	0.8	0.8	1.0
Tranquilizers	3.1	3.7	3.3	5.0	4.1	4.0	3.2	3.3	3.0	3.2	2.9	3.3	2.7	2.8	2.5	-0.3
Narcotics	1.4	1.4	1.3	2.2	1.7	1.6	1.2	1.2	1.0	1.1	1.0	1.1	0.9	0.9	1.0	
Psychedelics#	1.1	1.3	1.4	2.2	1.8	1.9	1.5	1.3	--	1.0	0.9	0.9	0.8	0.7	0.6	
LSD	--	--	--	--	--	--	--	--	--	0.8	0.8	0.8	0.7	0.5	0.5	
Others	--	--	--	--	--	--	--	--	--	0.4	0.4	0.4	0.3	0.4	0.3	
Heroin	0.1	0.8	0.6	0.9	0.8	0.5	0.5	0.5	0.5	0.6	0.6	0.6	0.5	0.5	0.4	
Steroids	0.1	1.3	0.9	1.0	0.8	0.6	1.0	0.7	0.7	0.8	0.7	0.8	0.6	0.6	0.6	
Injected drugs	--	--	--	--	--	--	--	--	0.5	0.7	0.6	0.6	0.6	0.6	0.6	
MDMA	--	--	--	--	--	--	--	--	--	1.0	0.9	0.8	0.6	0.6	0.6	
Rohypnol	--	--	--	--	--	--	--	--	--	0.4	0.4	0.4	0.3	0.3	0.3	
GHB	--	--	--	--	--	--	--	--	--	0.2	0.2	0.2	0.3	0.2	-0.1	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	3.4	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 7th Grade Students: 1993-2007**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
<b>CIGARETTES</b>																
Daily Use	7.3	8.6	8.8	10.4	8.3	6.9	6.0	5.6	4.4	4.2	3.6	4.2	3.3	3.0	2.7	-0.3
1/2+ Pack per Day	3.9	4.4	4.5	5.7	4.3	3.7	2.9	2.8	1.9	2.2	1.7	2.1	1.5	1.4	1.3	
CIGARS	--	--	--	--	--	1.2	1.0	1.0	1.3	0.9	0.7	1.0	0.7	0.6	0.7	
SMOKELESS TOBACCO	1.2	1.7	1.1	1.3	0.9	0.5	0.4	0.4	0.3	0.6	0.3	0.6	0.4	0.4	0.4	
<b>ALCOHOL</b>																
Daily Use	1.2	1.4	1.0	1.4	1.1	1.2	1.0	1.1	0.9	1.0	1.0	1.2	0.9	0.8	0.9	
Binge Drinking	12.7	13.0	12.3	13.5	12.0	12.3	12.4	12.4	9.7	10.0	8.0	8.9	6.8	7.0	8.3	1.3
MARIJUANA	0.1	0.7	0.9	1.4	1.0	1.1	1.0	1.0	0.9	0.9	1.0	0.9	0.7	0.8		

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates (p&lt;.05).

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National## 2006
Cigarettes	50.0	51.3	52.4	53.7	52.6	49.5	45.7	43.7	39.9	37.4	35.1	34.7	32.1	30.2	28.1	-2.1	24.6
Cigars	--	--	--	--	--	24.4	22.5	21.3	22.1	17.4	16.4	16.1	15.8	14.6	14.7	--	--
Pipe	--	--	--	--	--	8.0	8.6	7.8	--	5.5	5.5	5.6	5.2	4.9	5.8	0.9	--
Smokeless Tobacco	18.9	20.1	19.2	17.9	16.7	12.4	10.9	10.9	8.2	7.7	7.3	8.3	7.3	8.3	7.2	-1.1	10.2
Alcohol	60.9	61.7	60.2	59.1	59.4	51.5	53.0	51.0	53.3	52.6	51.7	52.0	49.0	46.5	45.4	-1.1	40.5
Marijuana	12.0	16.0	20.2	24.3	23.0	22.1	19.7	19.0	19.5	20.0	19.1	18.6	17.6	15.6	16.1	15.7	
Cocaine	2.0	2.5	2.9	3.6	3.5	3.2	2.6	2.6	2.7	2.6	2.9	2.7	2.9	2.5	2.4	3.4	
Crack	1.5	2.0	2.4	3.0	2.9	2.9	2.4	2.4	2.5	2.3	2.7	2.4	2.6	2.2	2.0	2.3	
Inhalants	14.5	15.3	18.1	17.4	17.6	13.9	12.0	11.2	10.3	9.3	10.7	11.0	11.3	10.4	10.5	16.1	
Amphetamines	9.0	9.1	10.6	10.0	9.1	6.6	5.8	5.6	5.5	5.0	4.7	4.3	3.7	3.2	3.0	7.3	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	-0.4	2.7
Ritalin	--	--	--	--	--	4.2	3.9	4.2	4.0	3.5	3.6	3.1	3.5	3.1	4.5	1.4	--
Tranquilizers	10.3	10.7	11.4	11.9	12.5	11.5	10.1	10.3	10.2	10.2	10.7	10.2	9.8	8.9	9.1	4.3	
Narcotics	5.3	5.4	5.8	5.5	6.5	5.8	5.2	5.0	4.9	4.8	4.8	4.5	4.5	4.1	5.0	0.9	--
Psychedelics#	3.7	3.7	5.1	6.7	6.3	5.6	5.7	5.4	--	4.2	3.7	3.1	3.3	2.8	2.6	3.4	
LSD	--	--	--	--	--	--	--	--	--	3.8	3.2	2.6	2.7	2.3	2.2	1.6	
Others	--	--	--	--	--	--	--	--	--	1.4	1.4	1.5	1.3	1.3	1.3	2.8	
Heroin	1.4	1.7	1.8	2.2	1.8	1.9	1.3	1.6	1.7	1.4	1.8	1.5	1.6	1.5	1.3	1.4	
Steroids	2.6	2.7	2.6	2.1	2.3	2.5	2.1	2.5	2.2	2.1	2.2	2.2	1.8	1.5	1.5	1.6	--
Injected drugs	--	--	--	--	--	--	--	--	--	1.5	1.4	1.7	1.5	1.8	1.7		
MDMA	--	--	--	--	--	--	--	--	--	3.4	3.3	2.7	2.5	2.3	2.3	2.5	
Rohypnol	--	--	--	--	--	--	--	--	--	1.1	1.3	1.1	1.2	1.2	1.1	1.0	
GHB	--	--	--	--	--	--	--	--	--	0.7	0.7	0.7	0.7	0.8	0.7	--	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	10.2	--	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Monitoring the Future Study, the University of Michigan, 2006

## Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National## 2006
Cigarettes	37.9	39.6	41.4	42.6	41.5	37.3	33.1	30.6	26.3	24.1	22.6	22.2	20.6	19.1	18.0	-1.1	--
Cigars	--	--	--	--	--	20.0	18.1	16.9	16.6	11.9	11.3	11.6	10.3	11.0	0.7	--	
Pipe	--	--	--	--	--	5.8	5.9	5.1	--	3.8	4.0	4.2	3.8	3.5	5.0	1.5	--
Smokeless Tobacco	14.9	16.5	15.4	14.2	12.8	9.5	8.3	8.0	6.2	5.8	5.5	6.3	5.6	6.4	5.8	-0.6	--
Alcohol	52.3	52.9	51.6	51.4	51.2	45.1	47.0	45.1	45.4	43.8	42.4	42.1	39.0	38.3	36.6	-1.7	33.6
Marijuana	10.4	14.3	18.4	22.4	21.0	19.8	17.7	17.1	16.8	16.0	15.4	14.5	12.7	13.1	11.7		
Cocaine	1.6	2.1	2.5	2.9	3.0	2.7	2.1	2.2	2.2	2.0	2.3	2.1	2.3	1.9	1.8	2.0	
Crack	1.2	1.6	2.0	2.3	2.4	2.4	1.9	1.9	2.0	1.7	2.0	1.7	1.8	1.7	1.5	1.3	
Inhalants	11.2	12.4	14.5	13.6	14.0	10.5	8.9	8.3	7.4	6.2	7.5	7.9	7.8	7.2	7.1	9.1	
Amphetamines	7.8	7.8	9.3	8.8	7.9	5.6	4.7	4.7	4.5	3.8	3.7	3.4	2.9	2.5	2.3	4.7	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	1.7	-0.2	1.8
Ritalin	--	--	--	--	--	3.6	3.2	3.5	3.1	2.5	2.6	2.2	2.6	2.3	3.3	1.0	2.6
Tranquilizers	8.6	9.1	9.5	10.3	10.8	9.7	8.5	8.5	8.5	7.9	8.3	8.1	7.5	6.8	6.7	2.6	
Narcotics	4.3	4.0	4.8	4.6	5.5	4.9	4.2	4.1	4.1	3.6	3.5	3.3	3.2	3.0	3.6	0.6	--
Psychedelics#	3.1	3.1	4.5	6.0	5.6	5.0	4.8	4.5	--	2.9	2.7	2.4	2.4	2.1	2.0	2.1	
LSD	--	--	--	--	--	--	--	--	--	2.7	2.3	1.9	2.0	1.7	1.6	0.9	
Others	--	--	--	--	--	--	--	--	--	1.0	1.0	1.1	1.0	1.1	1.0	1.8	
Heroin	1.1	1.5	1.5	1.8	1.6	1.5	1.1	1.2	1.3	1.0	1.4	1.2	1.2	1.0	0.9	0.8	
Steroids	2.0	2.0	1.8	1.6	1.7	2.0	1.5	1.8	1.7	1.6	1.6	1.4	1.1	1.2	0.9		
Injected drugs	--	--	--	--	--	--	--	--	1.1	1.0	1.2	1.2	1.3	1.2	1.3	--	
MDMA	--	--	--	--	--	--	--	--	--	2.7	2.6	2.0	1.8	1.7	1.7	1.4	
Rohypnol	--	--	--	--	--	--	--	--	--	0.8	1.0	0.9	0.9	0.8	0.8	0.5	
GHB	--	--	--	--	--	--	--	--	--	0.4	0.5	0.6	0.5	0.5	0.5	0.8	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	8.0	--	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Monitoring the Future Study, the University of Michigan, 2006

## Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National## 2006	
Cigarettes	21.1	24.2	26.3	27.1	25.8	24.1	19.7	18.2	16.1	14.7	14.0	13.4	12.5	11.6	10.8	-0.8	8.7	
Cigars	--	--	--	--	--	12.0	9.6	9.7	10.0	6.6	6.8	6.7	6.5	5.9	6.4	0.5	--	
Pipe	--	--	--	--	--	3.5	3.5	3.1	--	2.2	2.2	2.3	2.1	2.0	3.1	1.1	--	
Smokeless Tobacco	9.0	10.6	9.3	8.6	7.2	5.6	4.6	4.5	3.5	3.4	3.2	3.8	3.3	3.8	3.3	-0.5	3.7	
Alcohol	30.9	32.0	31.3	31.3	30.3	27.2	27.7	26.4	24.7	24.9	24.3	23.5	21.1	20.9	19.9	-1.0	17.2	
Marijuana	6.5	9.9	13.1	15.2	13.8	12.8	11.3	11.3	11.3	11.1	10.6	9.8	9.3	8.2	8.3	6.5	6.5	
Cocaine	0.1	1.3	1.4	1.7	1.9	1.5	1.1	1.1	1.3	1.2	1.3	1.2	1.4	1.1	1.1	1.0	1.0	
Crack	0.1	1.0	1.1	1.4	1.4	1.3	1.1	1.0	1.2	0.9	1.2	1.0	1.2	1.0	0.9	0.6	0.6	
Inhalants	6.5	7.5	8.4	8.0	8.2	5.7	4.8	4.4	4.0	3.4	4.2	4.7	4.3	4.1	3.7	-0.4	4.1	
Amphetamines	5.1	5.2	5.9	5.6	5.3	3.3	2.8	2.8	2.8	2.4	2.4	2.1	1.9	1.4	1.4	2.1	2.1	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	0.6	
Ritalin	--	--	--	--	--	2.0	1.7	2.0	1.7	1.3	1.4	1.3	1.5	1.3	1.9	0.6	--	
Tranquilizers	5.6	5.7	5.8	6.0	6.5	5.7	5.2	5.1	5.2	4.8	5.1	4.9	4.7	4.1	3.9	1.3	1.3	
Narcotics	2.6	2.5	2.6	2.7	3.2	2.8	2.4	2.4	2.4	2.0	2.1	1.9	1.9	1.8	2.1	0.3	--	
Psychedelics#	2.0	2.1	3.1	3.6	3.6	3.0	2.6	2.7	--	1.6	1.5	1.4	1.5	1.2	1.1	0.9	0.9	
LSD	--	--	--	--	--	--	--	--	--	1.5	1.3	1.2	1.2	0.9	0.9	0.4	0.4	
Others	--	--	--	--	--	--	--	--	--	0.6	0.7	0.7	0.6	0.6	0.6	0.7	0.7	
Heroin	0.1	1.0	0.9	1.1	1.1	1.0	0.6	0.7	0.8	0.7	0.9	0.8	0.8	0.7	0.6	0.3	0.3	
Steroids	1.2	1.3	1.0	1.0	1.0	1.1	0.9	0.9	1.0	0.9	1.0	0.9	0.8	0.7	0.7	0.5	0.5	
Injected drugs	--	--	--	--	--	--	--	--	--	0.7	0.6	0.8	0.8	0.8	0.8	--	--	
MDMA	--	--	--	--	--	--	--	--	--	--	1.6	1.5	1.3	1.1	1.0	0.9	0.7	0.7
Rohypnol	--	--	--	--	--	--	--	--	--	0.5	0.7	0.6	0.6	0.5	0.5	0.4	0.4	
GHB	--	--	--	--	--	--	--	--	--	0.3	0.4	0.4	0.4	0.4	0.3	--	--	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	5.2	--	--	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Monitoring the Future Study, the University of Michigan, 2006

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 8th Grade Students: 1993-2007**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National##	2006
<b>CIGARETTES</b>																		
Daily Use	11.8	13.8	14.9	15.4	14.2	13.7	10.6	10.0	8.8	7.5	7.2	6.9	6.7	5.8	5.5	4.0		
1/2+ Pack per Day	6.7	7.2	8.4	8.8	8.4	7.5	5.6	5.3	4.9	4.3	3.7	3.5	3.3	2.9	2.7	1.5		
<b>CIGARS</b>																		
SMOKELESS TOBACCO	2.3	2.6	2.1	2.2	1.6	1.4	0.9	0.9	0.9	0.8	0.7	0.9	0.8	1.0	0.8	0.7		
<b>ALCOHOL</b>																		
Daily Use	1.8	2.3	2.1	2.2	2.4	2.1	2.4	2.2	1.9	2.1	1.9	1.8	1.6	1.8	0.5			
Binge Drinking	18.5	18.6	18.4	19.0	18.9	19.7	19.5	18.6	14.8	15.5	13.4	13.4	11.6	11.0	13.2	2.2	10.9	
MARIJUANA	0.9	1.9	2.1	3.1	2.7	2.7	2.3	2.7	3.1	2.7	2.6	2.3	2.4	2.0	2.0	1.0		

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p<.05$ ).

## Monitoring the Future Study, the University of Michigan, 2006

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	
Cigarettes	56.5	57.4	58.7	61.4	57.3	55.8	53.7	50.6	47.2	43.6	39.8	40.7	36.8	33.5	-2.8		
Cigars	--	--	--	--	--	31.0	28.7	26.8	29.8	23.1	20.7	21.3	20.6	20.4	19.4	-1.0	
Pipe	--	--	--	--	--	10.9	10.1	9.7	--	7.6	7.1	7.3	6.5	6.9	9.1	2.2	
Smokeless Tobacco	24.3	25.8	26.2	24.2	21.3	18.7	15.0	14.2	11.9	12.4	10.5	12.2	10.7	11.6	10.8	-0.8	
Alcohol	70.2	69.9	68.7	69.7	65.9	61.8	61.4	59.8	62.6	59.9	58.4	58.4	55.2	55.5	51.7	-3.8	
Marijuana	18.4	23.9	26.1	35.0	28.9	30.7	27.8	26.7	28.4	28.1	25.2	24.0	22.9	22.5	21.2	-1.3	
Cocaine	2.5	3.7	4.9	4.8	5.3	4.2	3.9	3.8	4.3	4.3	4.7	4.2	4.4	3.7	-0.7		
Crack	1.8	2.5	2.7	4.0	3.5	4.2	3.3	3.0	3.0	3.1	3.2	3.3	3.0	3.0	2.4	-0.6	
Inhalants	13.3	17.4	17.3	18.3	16.7	15.7	12.3	11.6	11.1	9.9	9.8	11.3	10.9	11.4	9.8	-1.6	
Amphetamines	13.5	14.8	14.3	16.7	13.3	12.0	9.9	10.1	10.0	9.7	8.6	8.6	7.0	6.8	5.3	-1.5	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	3.5	3.0	2.2	-0.8	
Ritalin	--	--	--	--	--	6.8	5.0	6.2	6.3	5.3	5.1	4.9	5.4	5.4	7.3	1.9	
Tranquilizers	11.1	13.8	13.4	15.3	15.0	14.9	13.7	14.0	14.2	14.4	13.2	13.6	12.3	12.4	10.9	-1.5	
Narcotics	6.9	8.5	7.9	9.9	9.1	9.2	8.1	8.1	8.5	9.1	7.9	8.0	7.0	7.4	7.6		
Psychedelics#	5.6	6.4	7.8	12.1	8.9	9.9	9.2	8.0	--	6.9	5.2	5.2	4.7	4.2	3.6	-0.6	
LSD	--	--	--	--	--	--	--	--	--	6.3	4.5	4.3	3.9	3.4	2.9	-0.5	
Others	--	--	--	--	--	--	--	--	--	2.4	2.3	2.4	2.3	2.2	1.8	-0.4	
Heroin	1.5	2.0	1.9	2.4	2.1	2.1	1.7	1.6	1.8	1.8	2.0	2.0	2.1	1.9	1.5	-0.4	
Steroids	2.8	3.0	2.6	2.5	2.3	2.5	2.4	2.8	2.5	2.4	2.6	2.7	2.3	1.8	1.7		
Injected drugs	--	--	--	--	--	--	--	--	--	1.3	1.6	2.0	2.0	2.0	1.8	1.7	
MDMA	--	--	--	--	--	--	--	--	--	5.4	4.2	4.1	3.6	3.7	3.2	-0.5	
Rohypnol	--	--	--	--	--	--	--	--	--	1.6	1.6	1.8	1.8	1.8	1.4	-0.4	
GHB	--	--	--	--	--	--	--	--	--	1.1	1.1	1.1	1.1	1.0	0.9		
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	12.3		

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	
Cigarettes	43.6	45.5	46.2	50.2	46.4	43.4	39.7	37.1	33.3	29.1	27.0	27.6	25.4	25.0	22.6	-2.4	
Cigars	--	--	--	--	--	26.3	23.8	21.8	23.2	16.2	14.6	15.4	15.1	15.3	14.7		
Pipe	--	--	--	--	--	7.9	6.9	6.6	--	5.1	5.1	5.5	4.9	5.1	7.6	2.5	
Smokeless Tobacco	19.2	20.8	20.8	19.6	16.7	14.1	11.5	11.2	9.0	8.8	8.1	9.3	8.3	9.6	8.7	-0.9	
Alcohol	62.5	61.6	60.5	62.2	58.1	56.0	54.6	54.7	50.4	50.1	49.9	46.3	47.3	43.4	-3.9		
Marijuana	16.1	21.7	23.7	31.8	25.8	27.3	24.6	24.0	23.4	20.9	19.6	18.0	18.4	17.0	-1.4		
Cocaine	2.1	2.9	3.1	4.1	4.0	4.5	3.2	3.3	3.1	3.2	3.2	3.5	3.1	3.3	2.6	-0.7	
Crack	1.4	2.0	2.2	3.2	2.9	3.3	2.4	2.4	2.3	2.1	2.2	2.4	2.2	2.2	1.6	-0.6	
Inhalants	9.8	13.5	13.2	13.8	12.1	11.5	8.8	8.1	7.2	6.1	6.3	7.5	6.9	7.5	6.1	-1.4	
Amphetamines	11.6	12.9	12.2	14.4	11.4	9.8	8.2	8.0	8.5	7.2	6.7	6.6	5.2	5.1	3.7	-1.4	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	2.5	2.2	1.4	-0.8
Ritalin	--	--	--	--	--	5.7	3.9	4.8	4.9	3.7	3.6	3.4	3.6	3.8	5.3	1.5	
Tranquilizers	9.2	11.9	10.8	13.0	12.7	12.5	11.7	12.2	12.2	10.8	10.2	10.7	8.9	9.6	7.9	-1.7	
Narcotics	5.5	6.9	6.3	7.9	7.6	7.8	6.9	6.8	7.1	6.6	5.8	5.9	4.9	5.3	5.3		
Psychedelics#	4.7	5.6	6.9	10.9	7.7	8.5	7.9	6.7	--	4.7	3.6	3.7	3.1	3.1	2.5	-0.6	
LSD	--	--	--	--	--	--	--	--	--	4.4	3.1	3.1	2.5	2.4	2.0	-0.4	
Others	--	--	--	--	--	--	--	--	--	1.6	1.6	1.8	1.6	1.6	1.3	-0.3	
Heroin	1.1	1.5	1.7	2.0	1.7	1.6	1.3	1.4	1.3	1.1	1.4	1.4	1.4	1.1	1.1	-0.3	
Steroids	2.0	2.2	1.8	1.8	1.7	1.8	1.7	2.1	1.9	1.7	2.0	1.9	1.6	1.4	1.2	-0.2	
Injected drugs	--	--	--	--	--	--	--	--	0.9	1.1	1.4	1.4	1.4	1.3	1.4		
MDMA	--	--	--	--	--	--	--	--	--	3.9	3.1	3.0	2.5	2.7	2.2	-0.5	
Rohypnol	--	--	--	--	--	--	--	--	--	1.1	1.2	1.3	1.2	1.3	1.0	-0.3	
GHB	--	--	--	--	--	--	--	--	--	0.7	0.8	0.8	0.8	0.8	0.6		
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	9.2		

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
Cigarettes	26.2	29.4	30.8	34.4	30.5	28.9	25.4	24.9	21.8	19.1	17.9	18.2	16.4	16.5	14.7	-1.8
Cigars	--	--	--	--	--	15.5	13.2	12.6	14.1	9.2	8.3	9.1	8.6	8.8	8.7	
Pipe	--	--	--	--	--	4.5	4.1	4.2	--	2.9	2.7	3.2	2.7	3.0	4.4	1.4
Smokeless Tobacco	12.2	13.3	13.2	12.5	9.8	8.3	6.5	6.5	5.4	5.2	4.9	5.6	5.2	5.7	5.3	-0.4
Alcohol	38.2	39.6	38.5	40.8	36.0	36.7	36.2	35.3	33.2	30.5	30.8	29.4	26.6	28.5	24.8	-3.7
Marijuana	10.6	15.2	15.9	22.8	17.2	18.4	16.2	15.6	16.1	14.6	13.9	12.1	11.5	11.4	10.3	-1.1
Cocaine	0.1	1.6	1.7	2.4	2.2	2.4	1.4	1.6	1.6	1.6	1.8	2.0	1.7	1.7	1.4	-0.3
Crack	0.1	1.4	1.3	2.1	1.5	1.9	1.2	1.3	1.2	1.1	1.2	1.4	1.3	1.1	0.9	-0.2
Inhalants	5.2	7.3	6.9	7.2	6.1	5.8	4.4	4.2	3.7	3.0	3.0	3.9	3.6	3.6	2.9	-0.7
Amphetamines	6.4	7.8	7.2	8.9	6.8	5.9	4.8	5.1	5.2	4.3	3.6	3.8	3.0	3.0	2.0	-1.0
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	1.5	1.2	0.8	-0.4
Ritalin	--	--	--	--	--	3.0	2.0	2.5	2.4	1.9	1.8	1.9	2.0	2.0	2.8	0.8
Tranquilizers	5.2	7.1	6.2	7.7	7.9	7.5	7.2	7.4	7.3	6.7	5.9	6.3	5.1	5.7	4.4	-1.3
Narcotics	2.9	3.7	3.8	4.9	4.2	4.2	3.8	3.9	4.2	3.9	3.3	3.1	2.8	2.9	2.9	
Psychedelics#	2.7	3.7	4.9	6.5	4.6	5.0	4.3	3.6	--	2.4	1.8	1.9	1.7	1.6	1.3	-0.3
LSD	--	--	--	--	--	--	--	--	--	2.0	1.5	1.6	1.4	1.3	1.1	
Others	--	--	--	--	--	--	--	--	--	1.0	0.8	1.0	0.9	0.8	0.7	-0.1
Heroin	0.1	1.1	0.9	1.3	1.0	0.9	0.6	0.7	0.8	0.7	0.9	0.9	0.9	0.8	0.7	
Steroids	1.3	1.4	1.2	1.2	1.1	1.1	0.9	1.0	1.1	1.0	1.2	1.1	0.9	0.9	0.8	
Injected drugs	--	--	--	--	--	--	--	--	0.6	0.6	0.8	0.8	0.8	0.8	0.9	
MDMA	--	--	--	--	--	--	--	--	--	1.9	1.5	1.6	1.3	1.5	1.2	-0.3
Rohypnol	--	--	--	--	--	--	--	--	--	0.7	0.6	0.7	0.7	0.8	0.6	-0.2
GHB	--	--	--	--	--	--	--	--	--	0.5	0.5	0.5	0.5	0.5	0.4	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	5.6	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 9th Grade Students: 1993-2007**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
<b>CIGARETTES</b>																
Daily Use	16.0	18.3	18.4	22.0	19.0	18.5	15.5	14.7	12.4	11.6	10.3	9.4	9.6	8.1	-1.5	
1/2+ Pack per Day	9.9	11.7	11.2	14.0	11.8	11.6	9.2	8.4	7.0	7.0	6.1	5.7	5.1	4.4	4.4	-1.0
CIGARS	--	--	--	--	--	3.3	2.8	2.8	4.3	2.1	1.7	2.2	1.8	1.9	2.3	0.4
SMOKELESS TOBACCO	3.8	4.2	4.1	3.9	2.5	2.4	1.8	2.0	1.5	1.6	1.4	1.8	1.5	1.8	1.6	
<b>ALCOHOL</b>																
Daily Use	2.8	3.2	3.1	3.6	3.1	3.6	3.6	3.4	3.0	3.1	2.8	2.9	2.4	2.7	2.3	-0.4
Binge Drinking	23.3	23.6	23.3	25.5	23.1	25.7	24.3	24.3	19.7	19.8	17.9	17.3	14.8	15.6	16.9	1.3
MARIJUANA	1.8	3.3	3.7	6.4	4.1	4.9	4.6	4.5	4.2	4.5	3.8	3.2	3.3	3.2	2.7	-0.5

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates (p&lt;.05).

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 10th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National## 2006
Cigarettes	60.8	61.5	61.8	63.6	64.1	61.4	60.5	57.4	55.5	51.5	47.2	46.7	43.9	43.0	40.6	-2.4	36.1
Cigars	--	--	--	--	--	37.5	36.4	32.4	36.4	30.6	27.2	27.8	26.9	26.8	26.7	--	--
Pipe	--	--	--	--	--	12.8	12.1	10.6	--	9.0	8.3	7.7	7.8	8.6	13.3	4.7	--
Smokeless Tobacco	27.9	28.0	28.7	30.0	26.2	22.1	19.6	17.3	16.0	16.2	13.8	14.1	14.3	15.4	15.0	15.0	15.0
Alcohol	76.5	76.2	73.2	74.3	75.2	68.4	69.2	66.6	71.3	68.4	67.2	65.5	63.5	62.3	61.0	-1.3	61.5
Marijuana	25.6	30.5	34.2	40.2	40.1	38.2	37.4	34.1	37.8	36.9	34.8	33.5	31.6	30.1	29.9	31.8	31.8
Cocaine	3.4	4.3	4.9	6.9	6.7	6.6	6.0	5.1	6.1	6.5	6.0	6.0	6.1	5.9	5.8	4.8	4.8
Crack	1.9	2.9	3.6	4.8	4.7	5.1	4.1	3.4	3.9	3.5	3.7	3.8	3.8	3.4	3.1	2.2	2.2
Inhalants	15.0	16.1	17.5	17.8	17.5	15.9	12.8	11.4	11.9	10.7	10.9	11.3	11.2	11.1	10.6	13.3	13.3
Amphetamines	17.3	17.2	17.7	18.8	17.6	15.1	13.5	12.2	13.7	13.2	12.3	11.3	10.2	9.0	8.2	-0.8	11.2
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Ritalin	--	--	--	--	--	6.8	6.6	6.8	7.5	7.5	6.5	6.1	6.5	6.5	10.6	4.1	--
Tranquilizers	13.0	14.6	13.5	16.3	16.3	16.1	15.5	14.9	16.4	17.4	16.5	16.0	15.0	14.3	13.6	-0.7	7.2
Narcotics	9.0	9.8	9.7	11.7	12.6	11.9	11.1	10.1	11.7	12.2	11.6	11.0	9.9	9.8	10.9	1.1	--
Psychedelics#	7.7	8.3	11.5	14.1	13.2	12.9	12.6	10.9	--	9.4	7.0	6.1	5.7	5.3	4.8	-0.5	6.1
LSD	--	--	--	--	--	--	--	--	--	8.6	6.0	5.1	4.7	4.3	3.9	-0.4	2.7
Others	--	--	--	--	--	--	--	--	--	3.2	3.1	2.9	2.7	2.6	2.6	5.5	5.5
Heroin	1.4	2.0	1.9	2.8	2.2	2.5	2.0	1.9	2.1	2.2	2.1	2.1	2.0	2.0	2.0	1.4	1.4
Steroids	2.4	2.8	2.2	2.9	2.4	3.1	2.3	2.7	2.5	3.0	2.8	2.7	2.2	2.2	2.1	1.8	--
Injected drugs	--	--	--	--	--	--	--	--	--	1.6	1.8	1.9	1.9	2.1	2.0	2.0	--
MDMA	--	--	--	--	--	--	--	--	--	6.9	5.7	4.7	4.6	4.4	4.3	4.5	4.5
Rohypnol	--	--	--	--	--	--	--	--	--	1.8	1.9	1.8	1.8	1.7	1.7	0.8	0.8
GHB	--	--	--	--	--	--	--	--	--	1.3	1.1	1.2	1.2	1.1	1.2	--	--
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	14.4	--	--

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Monitoring the Future Study, the University of Michigan, 2006

**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 10th Grade Students: 1993-2007**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National## 2006	
Cigarettes	46.8	48.0	48.9	50.8	52.4	47.9	45.5	41.4	39.2	35.9	32.2	31.7	30.5	30.0	28.5	-1.5	--	
Cigars	--	--	--	--	--	32.1	30.4	26.0	28.5	22.3	19.6	21.1	19.9	20.2	20.7	--	--	
Pipe	--	--	--	--	--	8.9	8.3	6.6	--	6.2	5.9	5.5	5.8	6.3	11.1	4.8	--	
Smokeless Tobacco	21.6	21.8	22.2	22.9	20.3	16.4	14.6	12.9	11.8	12.0	10.2	10.7	11.4	12.2	12.0	--	--	
Alcohol	68.0	67.7	65.3	65.9	67.3	61.8	63.0	61.1	62.8	59.5	57.6	55.8	54.3	53.7	51.7	-2.0	55.8	
Marijuana	22.2	27.6	30.8	36.1	35.1	33.7	32.8	29.4	32.4	30.3	28.1	26.8	25.0	23.8	23.5	25.2		
Cocaine	2.6	3.5	4.0	5.7	5.5	5.4	5.0	3.8	4.9	4.7	4.5	4.5	4.3	4.0	3.2			
Crack	1.5	2.3	3.0	4.0	3.8	4.0	3.2	2.4	2.7	2.5	2.6	2.7	2.5	2.3	2.0	-0.3	1.3	
Inhalants	10.3	11.3	12.4	11.3	11.7	10.7	8.1	7.1	7.2	6.2	6.3	6.8	6.7	6.5	5.8	-0.7	6.5	
Amphetamines	13.9	14.4	14.4	15.4	14.5	12.1	10.5	9.6	11.1	10.0	9.2	8.5	7.2	6.8	5.7	-1.1	7.9	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	3.1	2.4	1.9	
Ritalin	--	--	--	--	--	5.2	4.9	5.0	5.3	5.0	4.4	4.0	4.3	4.5	7.4	2.9	3.6	
Tranquilizers	10.4	12.1	11.2	13.5	13.5	13.2	12.8	12.1	13.5	13.2	12.3	12.4	11.3	10.7	9.5	-1.2	5.2	
Narcotics	6.9	7.8	7.6	9.3	10.2	9.8	9.0	8.1	9.6	8.8	8.3	8.1	6.9	7.0	7.5	0.5	--	
Psychedelics#	6.4	7.1	10.2	12.3	11.4	10.5	10.5	8.6	--	6.1	4.5	4.0	3.7	3.7	3.2	-0.5	4.1	
LSD	--	--	--	--	--	--	--	--	--	5.5	3.6	3.2	3.0	2.9	2.4	-0.5	1.7	
Others	--	--	--	--	--	--	--	--	--	2.3	2.1	2.0	2.0	2.0	1.8	3.7		
Heroin	1.2	1.6	1.6	2.3	1.8	1.8	1.6	1.2	1.6	1.4	1.5	1.4	1.3	1.3	0.9			
Steroids	1.8	2.1	1.7	2.2	1.7	2.2	1.6	1.9	1.9	2.0	1.9	2.0	1.4	1.6	1.3	-0.3	1.2	
Injected drugs	--	--	--	--	--	--	--	--	1.1	1.2	1.3	1.3	1.3	1.4	1.5	--	--	
MDMA	--	--	--	--	--	--	--	--	--	5.1	3.8	3.2	3.0	3.0	2.9	2.8		
Rohypnol	--	--	--	--	--	--	--	--	--	1.3	1.2	1.2	1.4	1.1	-0.3	0.5		
GHB	--	--	--	--	--	--	--	--	--	0.9	0.8	0.8	0.9	0.7	0.7	0.7		
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	10.3	--		

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Monitoring the Future Study, the University of Michigan, 2006

## Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 10th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National## 2006
Cigarettes	30.4	33.0	34.4	36.7	37.2	33.9	31.5	28.9	27.0	23.8	22.2	22.1	20.8	20.6	19.3	-1.3	14.5
Cigars	--	--	--	--	--	19.0	16.5	14.6	17.8	11.9	11.5	12.4	11.8	11.6	12.3	0.7	--
Pipe	--	--	--	--	--	4.8	4.3	3.6	--	3.2	3.3	2.9	3.2	3.4	6.8	3.4	--
Smokeless Tobacco	13.2	13.6	13.8	14.6	12.3	9.5	8.2	7.4	6.9	7.1	6.0	6.4	7.0	7.8	7.3	-0.5	5.7
Alcohol	43.8	44.9	42.7	43.7	43.9	41.3	41.6	39.7	39.7	38.1	36.9	34.1	33.0	33.0	31.1	-1.9	33.8
Marijuana	14.3	18.9	21.5	24.9	23.7	21.8	21.1	19.0	21.4	19.2	18.2	17.2	16.0	14.6	14.4	14.2	
Cocaine	1.4	1.6	2.2	2.6	2.6	2.9	2.3	1.7	2.4	2.4	2.3	2.4	2.2	2.1	1.9	1.5	
Crack	0.1	1.2	1.6	1.8	1.9	2.1	1.5	1.1	1.5	1.1	1.4	1.4	1.3	1.2	1.0	0.7	
Inhalants	5.0	5.6	5.8	5.1	5.3	4.9	3.4	2.9	3.3	2.9	2.9	3.3	3.0	3.0	2.5	-0.5	2.3
Amphetamines	7.5	8.3	7.9	7.8	7.8	6.5	5.6	5.4	6.5	5.9	5.2	4.8	4.0	3.7	2.9	-0.8	3.5
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	0.7	
Ritalin	--	--	--	--	--	2.5	2.1	2.4	2.6	2.2	2.1	2.0	2.1	2.2	3.7	1.5	--
Tranquilizers	5.7	6.9	6.2	7.0	7.4	7.5	7.1	7.0	8.0	7.6	7.1	7.0	6.4	5.9	5.0	-0.9	2.4
Narcotics	3.4	4.0	4.1	4.4	5.0	5.2	4.4	4.4	5.5	4.6	4.2	4.3	3.6	3.8	3.9		--
Psychedelics#	3.7	4.1	7.0	6.8	6.5	6.0	5.5	4.2	--	2.5	2.1	2.0	1.8	1.9	1.6	-0.3	1.5
LSD	--	--	--	--	--	--	--	--	--	2.2	1.7	1.5	1.4	1.5	1.2	-0.3	0.7
Others	--	--	--	--	--	--	--	--	--	1.1	1.1	1.0	1.0	1.0	0.9	1.3	
Heroin	0.1	0.9	1.0	1.3	1.0	1.3	0.9	0.7	1.0	0.8	0.9	0.9	0.9	0.8	0.8	0.5	
Steroids	1.0	1.3	1.1	1.4	0.9	1.4	0.9	0.9	1.0	1.1	1.0	1.1	0.9	1.1	0.9	0.6	
Injected drugs	--	--	--	--	--	--	--	--	--	0.8	0.7	0.7	0.8	0.9	1.0		--
MDMA	--	--	--	--	--	--	--	--	--	2.4	1.8	1.5	1.5	1.5	1.4	1.2	
Rohypnol	--	--	--	--	--	--	--	--	--	0.6	0.7	0.7	0.7	0.8	0.7	0.2	
GHB	--	--	--	--	--	--	--	--	--	0.5	0.5	0.6	0.5	0.5	0.5	--	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	5.9	--	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Monitoring the Future Study, the University of Michigan, 2006

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 10th Grade Students: 1993-2007**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National##	2006
<b>CIGARETTES</b>																		
Daily Use	19.0	21.9	22.4	24.4	23.9	21.5	20.6	17.9	16.5	14.6	13.7	13.3	12.5	12.2	11.4	-0.8	7.6	
1/2+ Pack per Day	12.3	14.4	14.6	16.2	15.5	13.4	12.9	10.8	9.8	9.4	8.1	7.6	7.1	7.1	6.6	-0.5	3.3	
<b>CIGARS</b>																		
SMOKELESS TOBACCO	4.7	5.2	5.0	5.6	4.1	3.3	2.5	2.3	2.2	2.7	2.2	2.4	2.5	3.2	2.8	-0.4	1.7	
<b>ALCOHOL</b>																		
Daily Use	3.5	3.6	3.5	4.1	3.9	4.2	4.5	4.1	4.2	4.0	4.0	3.5	3.6	3.6	3.4	1.4		
Binge Drinking	27.3	26.8	26.1	27.3	27.8	29.0	28.8	27.1	24.2	24.9	21.8	20.2	19.3	19.9	21.7	1.8	21.9	
MARIJUANA	2.7	4.3	5.1	7.4	6.9	6.3	6.5	5.7	6.3	6.3	5.6	5.4	5.0	4.5	4.6	2.8		

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates (p&lt;.05).

## Monitoring the Future Study, the University of Michigan, 2006

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 11th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
Cigarettes	64.7	64.5	66.5	67.7	66.9	66.9	65.3	62.3	58.7	54.8	51.2	50.1	46.9	46.8	44.6	-2.2
Cigars	--	--	--	--	--	43.9	42.1	39.3	40.7	34.2	31.5	31.8	30.9	30.7	31.0	
Pipe	--	--	--	--	--	14.5	13.1	12.4	--	9.4	8.8	9.3	8.1	9.2	16.2	7.0
Smokeless Tobacco	31.9	31.0	32.2	31.9	29.1	25.6	23.0	22.9	17.4	17.6	16.3	18.8	15.8	17.2	16.9	
Alcohol	82.2	79.7	78.5	78.1	77.5	74.9	73.3	71.9	74.3	72.9	70.2	68.7	66.6	66.7	64.3	-2.4
Marijuana	31.1	33.2	36.6	44.1	42.3	43.7	42.1	39.8	41.0	41.1	38.4	35.9	34.3	34.4	32.4	-2.0
Cocaine	4.2	4.9	5.2	7.5	7.8	8.1	7.6	7.5	7.1	7.8	8.0	7.8	7.1	7.8	6.9	-0.9
Crack	2.0	2.7	3.2	4.1	4.5	5.3	4.2	3.8	3.8	4.0	3.8	4.6	3.7	3.8	3.3	-0.5
Inhalants	14.0	15.7	16.0	17.6	15.4	16.2	12.3	13.0	11.3	10.5	9.9	11.1	9.8	10.1	9.7	
Amphetamines	18.8	18.5	18.2	18.3	18.5	16.8	15.7	15.5	15.1	14.8	14.3	13.6	11.1	10.8	9.1	-1.7
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	-0.9
Ritalin	--	--	--	--	--	7.5	7.3	8.0	7.3	7.6	7.0	5.9	6.6	6.7	10.7	4.0
Tranquilizers	12.9	13.5	13.9	16.6	16.3	17.5	16.0	16.7	17.0	18.6	16.7	15.9	14.6	14.5	13.4	-1.1
Narcotics	9.7	11.3	10.8	13.2	13.9	13.7	13.1	13.3	12.8	14.1	13.1	12.6	11.1	11.3	11.7	
Psychedelics#	10.5	10.1	12.7	16.8	14.9	16.4	15.6	14.0	--	11.4	9.0	7.5	6.3	6.3	5.4	-0.9
LSD	--	--	--	--	--	--	--	--	--	10.7	7.9	6.4	4.9	4.9	4.4	
Others	--	--	--	--	--	--	--	--	--	3.6	3.9	3.7	3.8	3.8	3.0	-0.8
Heroin	1.2	1.5	1.7	1.9	1.9	2.4	2.0	2.5	2.1	2.1	2.4	2.1	2.3	2.0		
Steroids	2.0	2.6	2.5	2.3	2.4	2.7	2.2	2.7	2.3	2.5	2.3	2.1	1.9	1.8		
Injected drugs	--	--	--	--	--	--	--	--	1.5	1.5	1.8	1.9	2.1	2.1	1.9	
MDMA	--	--	--	--	--	--	--	--	--	8.1	6.5	5.7	4.9	5.4	4.8	-0.6
Rohypnol	--	--	--	--	--	--	--	--	--	1.5	1.5	1.9	1.8	2.0	1.7	
GHB	--	--	--	--	--	--	--	--	--	1.4	1.2	1.5	1.2	1.2	1.1	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	13.5	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 11th Grade Students: 1993-2007**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
Cigarettes	51.0	50.5	53.3	54.8	54.1	53.5	51.0	47.1	42.5	39.0	35.4	35.1	33.1	33.7	32.4	-1.3
Cigars	--	--	--	--	--	37.2	35.1	30.9	31.7	24.5	22.9	23.6	22.9	23.1	24.1	1.0
Pipe	--	--	--	--	--	9.6	8.7	7.5	--	6.3	6.1	6.5	5.5	6.6	13.3	6.7
Smokeless Tobacco	23.5	23.1	23.9	22.9	21.1	18.2	15.9	16.3	12.4	12.5	11.9	13.1	11.8	13.1	13.1	
Alcohol	73.6	70.2	70.9	70.0	70.0	68.8	67.2	65.6	65.8	63.5	59.8	58.6	56.0	57.0	55.3	-1.7
Marijuana	26.3	28.7	32.8	38.6	36.3	37.5	35.2	32.9	34.1	33.7	29.9	27.5	25.7	26.5	24.8	-1.7
Cocaine	2.9	3.6	4.3	5.8	6.3	6.7	5.9	5.5	5.4	5.7	5.8	5.7	4.8	5.3	4.6	-0.7
Crack	1.6	1.9	2.5	3.2	3.6	4.2	2.9	2.3	2.6	2.9	2.5	3.0	2.3	2.4	2.2	
Inhalants	8.6	10.1	9.9	10.1	9.4	9.6	6.9	7.1	6.0	5.4	5.1	6.0	5.0	5.2	4.7	
Amphetamines	14.5	13.9	14.2	13.9	14.1	12.5	11.6	12.2	11.9	10.8	10.2	9.6	7.2	7.6	5.8	-1.8
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	3.0	2.7	2.0
Ritalin	--	--	--	--	--	5.3	4.8	5.4	4.7	4.4	4.3	3.4	3.9	4.2	6.9	2.7
Tranquilizers	9.8	10.5	11.0	13.2	13.1	13.6	12.7	13.2	13.6	13.8	11.9	11.9	9.9	10.0	8.9	-1.1
Narcotics	7.0	8.0	8.1	10.3	10.8	11.0	10.4	10.4	10.0	10.0	8.9	9.0	7.2	7.6	7.7	
Psychedelics#	8.6	8.2	11.4	14.1	12.6	13.0	12.5	10.9	--	6.6	5.2	4.8	4.1	4.2	3.6	-0.6
LSD	--	--	--	--	--	--	--	--	--	6.1	4.0	3.8	2.9	3.1	2.6	-0.5
Others	--	--	--	--	--	--	--	--	--	2.4	2.7	2.4	2.5	2.6	2.2	-0.4
Heroin	0.1	1.1	1.3	1.5	1.6	1.8	1.5	1.7	1.5	1.2	1.4	1.5	1.3	1.6	1.4	
Steroids	1.4	1.8	1.6	1.7	1.5	1.7	1.5	1.8	1.6	1.6	1.7	1.6	1.5	1.3	1.3	
Injected drugs	--	--	--	--	--	--	--	--	--	1.1	1.0	1.2	1.1	1.3	1.4	
MDMA	--	--	--	--	--	--	--	--	--	5.6	4.2	3.7	2.9	3.7	3.0	-0.7
Rohypnol	--	--	--	--	--	--	--	--	--	0.9	1.0	1.2	1.0	1.2	1.1	
GHB	--	--	--	--	--	--	--	--	--	0.9	0.8	1.0	0.7	0.8	0.8	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	9.0	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 11th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
Cigarettes	33.7	34.4	39.3	40.1	39.2	39.8	37.4	34.1	30.2	27.1	25.2	25.1	23.5	23.8	22.5	-1.3
Cigars	--	--	--	--	--	21.0	18.6	16.8	19.8	13.4	13.1	13.9	13.0	12.8	13.8	1.0
Pipe	--	--	--	--	--	4.5	4.3	3.8	--	3.2	3.2	3.2	3.0	3.5	7.4	3.9
Smokeless Tobacco	14.4	14.6	15.4	14.2	12.0	10.2	8.9	9.7	7.0	7.5	7.2	8.1	7.3	8.0	8.1	
Alcohol	49.4	47.0	47.7	48.2	45.8	47.8	45.6	43.2	42.1	41.4	38.1	36.5	34.5	35.8	33.8	-2.0
Marijuana	16.7	19.6	21.8	25.7	22.3	24.6	22.0	20.5	21.7	21.2	18.4	17.2	15.3	15.7	14.6	-1.1
Cocaine	1.2	1.6	2.0	3.0	2.9	3.3	2.4	2.2	2.6	2.7	2.5	3.1	2.1	2.7	2.1	-0.6
Crack	0.1	1.0	1.3	1.6	1.6	1.9	1.0	1.0	1.2	1.1	1.1	1.8	1.1	1.3	1.1	
Inhalants	3.8	4.2	4.3	4.3	3.6	3.9	2.7	2.8	2.3	2.1	2.2	2.7	2.1	2.3	2.2	
Amphetamines	7.2	7.0	7.0	6.4	7.0	6.0	5.8	6.6	6.4	5.6	5.3	5.4	3.5	3.9	2.8	-1.1
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	1.5	1.5	1.1	-0.4
Ritalin	--	--	--	--	--	2.4	1.9	2.2	2.0	2.0	1.8	1.6	1.8	2.0	3.2	1.2
Tranquilizers	4.6	5.3	5.7	6.9	7.0	7.1	6.9	7.3	7.7	7.6	6.7	6.5	5.1	5.6	4.3	-1.3
Narcotics	3.4	3.5	3.8	4.8	4.7	5.2	4.7	5.3	5.0	4.8	4.4	4.7	3.6	4.2	3.9	
Psychedelics#	4.5	4.1	7.6	6.4	6.9	6.9	6.0	4.9	--	2.5	2.3	2.1	1.8	2.1	1.7	-0.4
LSD	--	--	--	--	--	--	--	--	--	2.1	1.7	1.7	1.3	1.6	1.3	-0.3
Others	--	--	--	--	--	--	--	--	--	1.1	1.3	1.1	1.1	1.3	1.0	-0.3
Heroin	0.1	0.6	0.8	0.8	0.8	1.0	0.7	0.9	0.7	0.6	0.8	1.1	0.8	1.0	0.9	
Steroids	0.1	1.1	1.0	0.8	0.8	1.0	0.6	0.9	0.7	0.8	1.0	1.0	0.9	0.8	0.7	
Injected drugs	--	--	--	--	--	--	--	--	0.7	0.6	0.7	0.8	0.8	0.8	0.8	1.1
MDMA	--	--	--	--	--	--	--	--	--	2.3	1.7	1.8	1.3	1.8	1.3	-0.5
Rohypnol	--	--	--	--	--	--	--	--	--	0.6	0.6	0.7	0.5	0.7	0.7	
GHB	--	--	--	--	--	--	--	--	--	0.4	0.5	0.6	0.5	0.6	0.5	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	4.7	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 11th Grade Students: 1993-2007**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*
<b>CIGARETTES</b>																
Daily Use	22.0	22.8	26.0	27.3	26.6	27.0	25.2	23.3	20.0	17.9	16.0	16.4	14.8	14.8	13.8	-1.0
1/2+ Pack per Day	15.0	15.3	17.2	18.9	18.1	17.3	16.2	15.0	12.8	11.9	9.9	10.3	8.8	8.5	8.1	
CIGARS	--	--	--	--	--	3.5	3.4	3.3	6.8	2.8	2.8	2.6	2.7	2.6	3.5	0.9
SMOKELESS TOBACCO	5.8	6.4	6.3	6.2	4.7	4.1	3.5	3.9	2.9	3.3	2.7	3.2	2.9	3.4	3.6	
<b>ALCOHOL</b>																
Daily Use	4.1	4.5	5.0	5.1	4.5	5.6	4.9	4.9	4.3	4.7	4.2	4.3	3.6	3.5	3.6	
Binge Drinking	30.6	28.9	29.6	30.1	29.8	32.3	31.3	30.0	25.7	26.5	24.0	23.3	21.2	21.8	23.2	1.4
MARIJUANA	3.9	5.1	6.1	8.6	7.1	7.6	7.5	6.9	7.0	7.4	6.2	6.2	5.3	4.9	4.6	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates (p&lt;.05).

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 12th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National## 2006	
Cigarettes	66.4	65.6	67.9	66.4	68.6	67.2	68.1	65.8	62.9	60.5	55.9	53.9	51.9	50.9	48.4	-2.5	47.1	
Cigars	--	--	--	--	--	50.1	48.9	45.3	46.7	41.2	39.4	37.3	38.9	37.7	37.9	--	--	
Pipe	--	--	--	--	--	15.9	15.6	14.0	--	10.9	9.7	10.0	10.5	10.7	20.0	9.3	--	
Smokeless Tobacco	34.0	33.9	35.8	31.3	31.4	29.4	26.9	26.3	22.9	21.2	18.7	19.1	18.8	20.4	19.4	-1.0	15.2	
Alcohol	85.0	82.4	81.4	80.6	80.8	76.1	78.5	75.9	78.6	77.3	74.8	72.4	72.3	70.2	69.2	-1.0	72.7	
Marijuana	35.3	38.1	40.0	43.4	45.6	46.5	45.7	44.4	45.5	44.8	42.3	40.5	40.1	37.1	36.5	--	42.3	
Cocaine	5.0	6.1	7.1	7.1	8.9	9.3	8.4	8.7	9.0	8.6	8.7	8.8	8.8	8.6	7.8	-0.8	8.5	
Crack	2.0	2.8	4.3	4.1	5.6	5.6	4.5	4.2	4.1	4.1	4.0	4.4	3.9	4.0	3.2	-0.8	3.5	
Inhalants	14.0	15.1	15.5	13.7	15.5	15.3	12.8	12.3	11.6	10.3	9.6	9.5	9.6	10.1	8.5	-1.6	11.1	
Amphetamines	19.8	18.8	18.0	16.5	19.2	16.6	15.6	15.4	15.5	14.9	14.4	13.3	12.5	11.3	9.0	-2.3	12.4	
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	5.5	5.0	3.4	-1.6	
Ritalin	--	--	--	--	--	5.9	7.4	7.4	7.8	7.2	6.9	6.3	7.1	7.0	11.3	4.3	--	
Tranquilizers	12.6	13.2	14.2	16.2	16.4	15.8	15.7	16.5	17.3	16.6	15.6	15.5	14.5	12.9	-1.6	10.3	--	
Narcotics	11.3	11.5	11.4	11.7	14.5	14.7	13.6	13.8	14.7	14.5	13.4	12.6	12.5	12.0	12.1	--	--	
Psychedelics#	11.6	12.0	14.4	15.5	17.2	17.8	17.1	15.4	--	12.8	10.6	8.1	7.3	7.0	6.1	-0.9	8.3	
LSD	--	--	--	--	--	--	--	--	--	11.9	9.3	6.6	5.5	5.3	4.5	-0.8	3.3	
Others	--	--	--	--	--	--	--	--	--	4.9	4.6	4.3	4.4	4.6	3.8	-0.8	7.8	
Heroin	1.3	1.8	2.0	2.1	2.3	2.7	2.1	2.3	2.3	1.9	2.1	2.3	2.0	2.3	1.9	-0.4	1.4	
Steroids	2.4	2.8	2.6	2.1	2.5	2.6	2.5	2.5	2.3	2.4	2.3	2.3	2.1	2.3	1.9	-0.4	2.7	
Injected drugs	--	--	--	--	--	--	--	--	--	1.8	1.4	1.7	1.8	1.7	2.2	1.7	-0.5	--
MDMA	--	--	--	--	--	--	--	--	--	9.5	7.4	6.4	5.9	5.6	5.1	6.5	--	--
Rohypnol	--	--	--	--	--	--	--	--	--	1.4	1.5	1.6	1.4	1.7	1.5	--	--	
GHB	--	--	--	--	--	--	--	--	--	1.6	1.4	1.2	1.1	1.3	1.1	--	--	
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	13.5	--	--	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Monitoring the Future Study, the University of Michigan, 2006

**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 12th Grade Students: 1993-2007**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National## 2006	
Cigarettes	52.2	51.8	54.5	52.7	55.7	53.4	54.6	51.2	46.7	43.1	39.7	38.4	37.5	36.9	35.2	-1.7	--	
Cigars	--	--	--	--	--	42.9	40.4	36.8	37.8	29.4	29.1	28.4	29.7	29.1	30.5	1.4	--	
Pipe	--	--	--	--	--	11.1	10.2	9.0	--	6.9	6.4	6.7	7.1	7.8	16.6	8.8	--	
Smokeless Tobacco	23.8	24.6	25.9	21.6	21.4	20.2	18.2	18.3	15.6	14.5	12.9	13.8	13.7	15.4	14.6	--	--	
Alcohol	76.1	73.3	72.9	72.1	73.1	70.9	72.4	70.2	70.5	68.2	65.5	62.4	62.2	61.5	60.2	-1.3	66.5	
Marijuana	28.3	31.9	34.6	37.4	37.1	38.8	37.1	35.4	36.4	33.7	31.6	29.7	28.6	27.5	26.6	31.5		
Cocaine	3.3	4.1	5.6	5.7	6.9	7.1	6.2	6.5	6.1	6.1	6.0	6.0	6.0	6.1	5.3	-0.8	5.7	
Crack	1.3	2.0	3.4	3.0	4.2	4.1	2.9	2.8	2.4	2.5	2.3	2.8	2.5	2.5	2.0	-0.5	2.1	
Inhalants	7.1	8.9	9.0	7.4	8.3	8.1	6.3	6.3	5.4	4.6	4.3	4.6	4.7	4.6	4.0	-0.6	4.5	
Amphetamines	13.8	13.5	13.1	12.1	13.6	11.9	10.8	11.4	11.1	10.1	9.7	9.0	7.6	7.5	5.4	-2.1	8.1	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	--	3.1	1.9	-1.2	2.5
Ritalin	--	--	--	--	--	4.0	4.3	4.8	4.7	4.0	3.6	3.3	3.3	3.8	7.0	3.2	4.4	
Tranquilizers	9.3	9.7	9.5	11.1	12.3	12.8	11.9	12.3	12.7	11.9	11.5	11.1	9.9	9.8	8.3	-1.5	6.6	
Narcotics	7.3	7.8	8.1	8.5	10.7	11.2	9.9	10.4	10.5	9.7	8.7	8.3	7.9	7.5	7.6	--	--	
Psychedelics#	9.1	9.3	12.2	12.5	13.6	13.6	12.6	11.3	--	6.5	5.2	4.8	4.3	4.5	4.0	-0.5	4.9	
LSD	--	--	--	--	--	--	--	--	--	5.6	3.7	3.5	2.8	3.1	2.6	-0.5	1.7	
Others	--	--	--	--	--	--	--	--	--	3.1	2.9	3.0	2.8	3.2	2.7	-0.5	4.6	
Heroin	0.1	1.2	1.5	1.8	1.7	1.9	1.4	1.8	1.5	1.0	1.2	1.5	1.2	1.5	1.3	0.8		
Steroids	1.5	1.8	1.6	1.5	1.7	1.9	1.7	1.7	1.5	1.4	1.4	1.5	1.3	1.5	1.3	1.8		
Injected drugs	--	--	--	--	--	--	--	--	1.1	0.8	0.9	1.2	1.0	1.2	1.3	--	--	
MDMA	--	--	--	--	--	--	--	--	--	6.4	4.3	3.9	3.4	3.5	3.2	4.1		
Rohypnol	--	--	--	--	--	--	--	--	--	1.0	0.8	1.0	0.8	1.0	1.0	1.1		
GHB	--	--	--	--	--	--	--	--	--	0.9	0.8	0.8	0.6	0.7	0.8	1.1		
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	8.5	--		

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Monitoring the Future Study, the University of Michigan, 2006

## Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 12th Grade Students: 1993-2007

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National## 2006
Cigarettes	35.6	37.3	40.6	39.8	41.6	41.6	40.5	38.9	35.1	30.5	28.8	27.4	26.5	26.9	24.3	-2.6	21.6
Cigars	--	--	--	--	--	25.8	23.0	20.2	23.9	16.8	17.0	16.8	16.6	17.5	0.9	--	--
Pipe	--	--	--	--	--	4.8	5.2	4.4	--	3.3	3.3	3.4	3.7	4.6	9.3	4.7	--
Smokeless Tobacco	15.4	15.5	16.5	14.4	13.5	12.1	11.2	11.3	9.4	8.3	7.9	8.7	8.6	9.8	9.1	--	6.1
Alcohol	53.2	53.7	51.2	52.5	50.1	52.0	51.7	50.1	49.9	48.1	46.1	42.2	41.8	42.2	39.7	-2.5	45.3
Marijuana	17.8	21.6	24.0	25.4	23.5	24.5	23.5	22.7	23.5	20.5	19.8	18.3	17.8	17.2	15.8	-1.4	18.3
Cocaine	1.5	2.0	2.5	3.0	3.3	3.5	2.4	2.7	2.8	2.9	2.5	2.9	2.9	3.1	2.4	-0.7	2.5
Crack	0.1	0.9	1.7	1.5	1.8	2.0	1.1	1.1	1.1	1.1	1.4	1.2	1.4	1.1	-0.3	0.9	
Inhalants	2.9	3.8	3.8	3.1	3.3	3.2	2.6	2.4	2.0	1.9	1.7	1.8	1.9	2.2	1.5	-0.7	1.5
Amphetamines	6.5	7.2	6.5	6.1	6.7	5.8	5.1	5.9	5.7	5.3	4.9	4.5	3.8	4.1	2.5	-1.6	3.7
Methamphetamines	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Ritalin	--	--	--	--	--	1.7	1.5	1.9	2.0	1.4	1.5	1.4	1.7	1.8	2.9	1.1	--
Tranquilizers	4.5	4.6	4.9	6.0	6.2	6.6	5.9	6.5	6.7	6.0	5.9	5.4	5.2	5.3	4.0	-1.3	2.7
Narcotics	3.1	3.6	4.0	4.3	4.5	4.6	4.5	4.5	5.2	4.6	4.0	4.0	3.8	4.0	3.8	--	--
Psychedelics#	4.5	4.4	8.2	6.0	7.2	7.0	5.7	4.5	--	2.0	2.2	2.0	1.8	2.1	1.6	-0.5	1.5
LSD	--	--	--	--	--	--	--	--	--	1.5	1.4	1.2	1.3	1.1	-0.2	0.6	
Others	--	--	--	--	--	--	--	--	--	1.1	1.3	1.2	1.5	1.1	-0.4	1.3	
Heroin	0.1	0.7	0.9	1.1	1.0	1.1	0.7	0.9	0.8	0.6	0.6	0.8	0.7	1.0	0.8	0.4	
Steroids	0.1	0.9	0.8	0.8	1.0	1.1	0.9	0.6	0.7	0.6	0.7	0.8	0.8	1.0	0.7	-0.3	
Injected drugs	--	--	--	--	--	--	--	--	--	0.6	0.6	0.7	0.6	0.9	0.9	--	--
MDMA	--	--	--	--	--	--	--	--	--	2.1	1.6	1.4	1.3	1.7	1.4	1.3	
Rohypnol	--	--	--	--	--	--	--	--	--	0.5	0.4	0.5	0.5	0.7	0.6	--	--
GHB	--	--	--	--	--	--	--	--	--	0.4	0.4	0.4	0.5	0.5	--	--	--
Over the counter drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	4.3	--	--

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

# Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

## Monitoring the Future Study, the University of Michigan, 2006

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 12th Grade Students: 1993-2007**

DRUG	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	Change*	National##	2006
<b>CIGARETTES</b>																		
Daily Use	23.2	25.1	27.6	27.0	28.6	28.5	27.6	26.5	23.3	19.9	18.7	18.0	16.4	16.9	14.7	-2.2	12.2	
1/2+ Pack per Day	16.0	17.9	18.9	18.6	19.3	20.0	18.1	17.0	15.3	13.3	11.9	10.9	9.9	10.5	8.5	-2.0	5.9	
<b>CIGARS</b>																		
SMOKELESS TOBACCO	6.9	7.0	7.6	6.7	6.1	5.3	5.0	4.9	4.1	3.8	3.4	4.0	3.5	4.4	4.2	2.2	--	
<b>ALCOHOL</b>																		
Daily Use	6.0	6.9	6.4	7.5	5.9	8.1	7.1	6.8	6.2	6.5	6.0	5.6	5.4	5.6	4.6	-1.0	3.0	
Binge Drinking	34.3	34.8	33.3	35.6	33.2	37.5	36.3	35.4	31.9	31.5	29.3	27.0	25.9	27.3	28.6	1.3	26.5	
MARIJUANA	4.2	6.0	7.4	7.6	8.1	8.5	8.2	7.5	8.2	7.3	7.4	6.7	6.3	6.1	5.3	-0.8	5.0	

NOTES: -- Data not available

\* Statistically significant changes between 2006 and 2007 prevalence rates ( $p < .05$ ).

## Monitoring the Future Study, the University of Michigan, 2006

## Mean Age of First Time Use of Alcohol, Tobacco, and Other Drugs, 2007

DRUG	6th	7th	8th	9th	10th	11th	12th	Total
Cigarettes	9.8	10.6	11.3	12.1	12.8	13.7	14.6	12.6
Cigars	9.9	10.8	11.6	12.7	13.5	14.5	15.5	13.4
Pipe	10.2	11.1	11.9	12.9	13.7	14.5	15.6	13.8
Smokeless Tobacco	10.0	10.8	11.7	12.6	13.5	14.3	15.0	13.3
Alcohol	9.9	10.9	11.8	12.8	13.5	14.2	14.9	12.9
Marijuana	10.7	11.5	12.3	13.1	13.8	14.5	15.1	13.7
Cocaine	10.4	11.4	12.3	13.4	14.2	14.9	15.7	14.2
Crack	10.0	11.1	12.0	12.9	13.6	14.3	15.0	13.3
Inhalants	9.7	10.7	11.5	12.4	13.1	13.8	14.2	12.2
Amphetamines	10.2	11.6	12.3	13.3	14.0	14.7	15.3	14.0
Methamphetamines	10.4	11.3	12.2	13.2	13.7	14.3	15.1	13.5
Ritalin	10.0	11.1	12.3	13.4	14.2	14.8	15.5	14.0
Tranquilizers	10.4	11.5	12.4	13.3	14.1	14.8	15.4	13.5
Narcotics	10.4	11.7	12.6	13.5	14.2	14.9	15.6	14.1
LSD	10.2	11.5	12.3	13.4	13.9	14.5	15.4	13.8
Others Psychedelics	10.1	11.1	11.9	12.9	13.7	14.6	15.6	13.8
Heroin	10.2	11.0	11.7	12.6	13.3	13.9	14.3	12.9
Steroids	10.2	11.0	11.8	13.0	13.4	14.1	14.8	12.9
MDMA	10.2	11.6	12.4	13.4	14.1	14.9	15.8	14.1
Rohypnol	10.5	11.0	11.7	12.7	13.2	14.0	14.3	13.0
GHB	10.1	10.7	11.4	12.4	12.6	13.5	14.0	12.5
Over the counter drugs	9.6	10.7	11.6	12.7	13.4	14.3	14.8	12.8

### III. REGIONAL DATA

<b>Regional Prevalence Rates by Grade: Lifetime, Annual, Monthly, Daily and Special Use</b>	.....	82
• Grade 6 .....	.....	82
• Grade 7 .....	.....	86
• Grade 8 .....	.....	90
• Grade 9 .....	.....	94
• Grade 10 .....	.....	98
• Grade 11 .....	.....	102
• Grade 12 .....	.....	106

**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	11.1	10.0	12.9 ▲	8.7 ▼	13.3 ▲	8.9 ▼	17.9 ▲	8.5 ▼	14.5 ▲
Cigars	4.8	4.1	5.6	3.5 ▼	5.9 ▲	3.7 ▼	8.2 ▲	3.4 ▼	6.4 ▲
Pipe	1.3	0.8	1.3	0.5 ▼	1.7	1.0	1.9 ▲	1.1	1.9 ▲
Smokeless Tobacco	2.4	1.8	1.9	1.5 ▼	3.6 ▲	1.6 ▼	3.8 ▲	1.8 ▼	4.4 ▲
Alcohol	22.5	24.4 ▲	25.2 ▲	19.4 ▼	25.0 ▲	19.6 ▼	25.0 ▲	21.0 ▼	25.4 ▲
Marijuana	3.0	3.1	3.7	2.3	2.3	3.1	5.1 ▲	2.2 ▼	3.0
Cocaine	0.5	0.4	0.5	0.3	0.5	0.5	0.9 ▲	0.3	0.5
Crack	0.7	0.6	0.6	0.4	0.9	0.7	1.6 ▲	0.4 ▼	0.7
Inhalants	6.2	7.4 ▲	6.2	6.2	6.5	5.4 ▼	8.2 ▲	4.9 ▼	7.2 ▲
Amphetamines	0.4	0.4	0.5	0.3	0.7 ▲	0.3	0.4	0.3	0.4
Methamphetamines	0.4	0.4	0.3	0.4	0.5	0.4	0.9 ▲	0.1	0.3
Ritalin	1.0	0.8	1.2	0.6	0.9	0.9	2.2 ▲	1.0	1.2
Tranquilizers	4.0	3.0 ▼	4.5	3.7	3.5	4.1	4.7	3.5	4.5
Narcotics	0.8	0.5	0.7	0.6	0.9	0.7	1.4 ▲	0.6	1.2 ▲
Psychedelics <sup>#</sup>	0.5	0.4	0.7	0.5	0.6	0.5	0.9 ▲	0.3 ▼	0.6
LSD	0.3	0.4	0.5	0.2	0.3	0.4	0.4 ▲	0.1	0.4
Others	0.3	0.1	0.4	0.4	0.4	0.2	0.9 ▲	0.2	0.3
Heroin	0.4	0.4	0.3	0.4	0.3	0.4	0.7	0.3	0.3
Steroids	0.7	0.5	0.7	0.4	1.0	0.7	0.9	0.5	0.7
Injected Drugs	0.8	0.7	0.7	0.8	1.0	0.8	0.9	0.7	0.9
MDMA	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.1	0.3
Rohypnol	0.2	0.2	0.3	0.1	0.2	0.2	0.4	0.0	0.3
GHB	0.2	0.1	0.3	0.1	0.1	0.1	0.4	0.1	0.2
Over the counter drugs	4.7	3.7 ▼	5.1	4.0	4.4	4.7	6.4 ▲	4.1	5.7 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	5.3	4.6	6.1	3.9	6.7 ▲	4.2 ▼	8.5 ▲	3.9 ▼	7.5 ▲
Cigars	3.1	2.9	3.3	2.6	3.9	2.4 ▼	5.7 ▲	2.3 ▼	4.1 ▲
Pipe	1.0	0.9	1.1	0.6	1.3	0.9	1.4	1.0	1.2
Smokeless Tobacco	1.7	1.7	0.9 ▲	1.3	2.7 ▲	1.1 ▼	2.6 ▲	1.3	3.2 ▲
Alcohol	15.3	17.6 ▲	17.1 ▲	13.5	16.6	13.2 ▼	16.8	14.0 ▼	17.8 ▲
Marijuana	2.3	2.3	2.8	1.7	2.0	2.2	3.7 ▲	1.7 ▼	2.3
Cocaine	0.4	0.3	0.4	0.2	0.1	0.4	0.9 ▲	0.3	0.3
Crack	0.5	0.3	0.5	0.5	0.6	0.4	1.2 ▲	0.4	0.4
Inhalants	4.1	5.0 ▲	3.7	4.0	4.5	3.5 ▼	5.6 ▲	3.4	4.7
Amphetamines	0.3	0.4	0.4	0.3	0.5	0.2	0.5	0.3	0.2
Methamphetamines	0.2	0.2	0.1	0.3	0.2	0.2	0.7 ▲	0.1	0.2
Ritalin	0.7	0.5	0.7	0.3	0.7	0.6	1.4 ▲	0.9	0.7
Tranquilizers	2.8	2.2	3.3	2.6	2.4	2.9	3.5	2.5	3.0
Narcotics	0.5	0.4	0.4	0.4	0.5	0.4	0.7	0.4	0.6
Psychedelics <sup>#</sup>	0.4	0.3	0.4	0.4	0.3	0.5	0.9 ▲	0.4	0.3
LSD	0.2	0.1	0.3	0.2	0.1	0.3	0.2	0.1	0.2
Others	0.3	0.2	0.3	0.4	0.2	0.3	0.7 ▲	0.3	0.2
Heroin	0.2	0.3	0.3	0.3	0.3	0.2	0.4	0.2	0.1
Steroids	0.5	0.5	0.5	0.3	0.6	0.5	0.6	0.4	0.6
Injected Drugs	0.6	0.5	0.6	0.5	0.7	0.5	0.9	0.5	0.5
MDMA	0.2	0.2	0.1	0.2	0.0	0.2	0.2	0.1	0.1
Rohypnol	0.1	0.1	0.1	0.1	0.0	0.1	0.2	0.0	0.2
GHB	0.1	0.1	0.2	0.1	0.0	0.1	0.4 ▲	0.0	0.1
Over the counter drugs	3.8	3.0	4.0	3.4	3.1	3.9	5.9 ▲	3.4	4.4

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	2.8	2.2	3.4	2.4 ▼	3.3	2.4	3.7 ▲	1.9 ▼	3.7 ▲
Cigars	1.6	1.3	1.7	1.3	1.6	1.2 ▼	3.0 ▲	1.3	2.4 ▲
Pipe	0.5	0.7	0.5	0.3	0.5	0.4	0.6	0.6	0.8
Smokeless Tobacco	0.9	0.9	0.6	0.6	1.6 ▲	0.5 ▼	1.2	0.8	1.8 ▲
Alcohol	6.9	7.6	7.9 ▲	6.5 ▼	6.9	5.8 ▼	7.2	6.0 ▼	8.8 ▲
Marijuana	1.4	1.6	1.6	1.2	1.1	1.4	2.2 ▲	1.0	1.5
Cocaine	0.2	0.1	0.2	0.3	0.1	0.2	0.4	0.1	0.2
Crack	0.3	0.3	0.2	0.3	0.3	0.3	0.7 ▲	0.2	0.3
Inhalants	2.3	2.6	2.4	2.1	2.6	2.0	3.0	2.0	2.8
Amphetamines	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.3	0.1
Methamphetamines	0.1	0.2	0.1	0.1	0.1	0.1	0.3	0.1	0.1
Ritalin	0.4	0.4	0.4	0.4	0.4	0.3	0.9 ▲	0.6	0.4
Tranquilizers	1.5	1.3	1.6	1.5	1.3	1.7	2.0	1.4	1.5
Narcotics	0.3	0.2	0.2	0.3	0.4	0.2	0.6	0.3	0.3
Psychedelics <sup>#</sup>	0.2	0.2	0.2	0.3	0.3	0.2	0.4	0.1	0.1
LSD	0.1	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.1
Others	0.1	0.1	0.1	0.2	0.2	0.0	0.3	0.1	0.1
Heroin	0.2	0.2	0.1	0.2	0.1	0.1	0.2	0.2	0.1
Steroids	0.3	0.4	0.2	0.1	0.3	0.3	0.4	0.2	0.4
Injected Drugs	0.3	0.3	0.5	0.4	0.4	0.3	0.4	0.3	0.3
MDMA	0.1	0.2	0.1	0.1	0.0	0.1	0.1	0.1	0.0
Rohypnol	0.1	0.0	0.0	0.1	0.0	0.1	0.2	0.0	0.1
GHB	0.1	0.0	0.1	0.1	0.0	0.0	0.2	0.0	0.1
Over the counter drugs	2.4	2.0	2.3	2.1	1.9	2.6	3.5 ▲	1.9	3.1 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
<b>CIGARETTES</b>									
Daily Use	1.0	0.7	1.4	0.8	0.9	0.9	1.3	0.8	1.5 ▲
1/2+ Pack per Day	0.4	0.4	0.7	0.3	0.4	0.5	0.1	0.3	0.4
<b>CIGARS</b>									
	0.2	0.2	0.3	0.1	0.1	0.1 ▼	0.5 ▲	0.2	0.4
<b>SMOKELESS TOBACCO</b>									
	0.1	0.0	0.1	0.1	0.2	0.1	0.1	0.1	0.3
<b>ALCOHOL</b>									
Daily Use	0.3	0.2	0.5	0.1	0.3	0.2	0.5	0.2	0.5
Binge Drinking	4.9	5.3 ▲	6.1 ▲	4.4	4.7	4.5	4.6	3.9 ▼	6.2 ▲
<b>MARIJUANA</b>									
	0.2	0.1	0.2	0.2	0.1	0.2	0.3	0.1	0.2

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ( $p<.05$ ).  
 ▼ indicates that local prevalence rates are significantly lower than state rates ( $p<.05$ ).

**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 7th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	17.9	15.9 ▼	18.3	16.5	21.8 ▲	14.1 ▼	26.3 ▲	14.6 ▼	23.9 ▲
Cigars	8.6	7.5	9.2	8.1	10.3 ▲	6.5 ▼	12.6 ▲	7.5	12.4 ▲
Pipe	3.0	3.4	2.9	2.5	3.8 ▲	2.1 ▼	4.3 ▲	2.4	4.5 ▲
Smokeless Tobacco	4.3	2.7 ▼	3.7	2.4 ▼	6.9 ▲	2.6 ▼	5.7 ▲	5.0	8.1 ▲
Alcohol	32.7	33.6	31.2	28.3 ▼	35.8 ▲	29.9 ▼	34.4	33.8	38.9 ▲
Marijuana	7.7	7.8	7.6	6.6	7.0	7.2	12.1 ▲	3.5 ▼	10.3 ▲
Cocaine	1.2	1.0	1.6	1.4	1.0	1.1	1.4	0.9	1.7 ▲
Crack	1.3	0.9	1.5	1.6	0.9	1.1	2.0 ▲	0.8	1.9 ▲
Inhalants	7.9	8.8	7.3	6.0 ▼	7.4	7.2	10.7 ▲	6.8	10.0 ▲
Amphetamines	1.2	0.8	1.4	0.8	1.5	1.0	1.9 ▲	0.9	1.8 ▲
Methamphetamines	1.0	0.9	1.1	0.8	0.9	0.9	1.1	0.7	1.7 ▲
Ritalin	2.2	2.3	2.0	1.4 ▼	2.3	2.1	3.2 ▲	1.8	2.7
Tranquilizers	5.6	5.9	5.1	4.3 ▼	5.0	5.2	8.1 ▲	3.9 ▼	7.7 ▲
Narcotics	2.2	1.9	1.9	1.3 ▼	2.2	1.9	3.4 ▲	1.7	3.6 ▲
Psychedelics <sup>#</sup>	1.2	1.3	1.1	0.8	1.1	1.3	1.4	1.0	1.5
LSD	1.0	1.0	0.9	0.5	0.8	1.1	0.9	0.8	1.2
Others	0.6	0.9	0.6	0.6	0.5	0.5	0.8	0.5	0.8
Heroin	0.9	1.0	0.9	0.9	0.6	0.8	1.2	0.8	1.2
Steroids	1.2	1.0	1.2	0.9	0.9	1.1	2.0 ▲	0.9	1.8 ▲
Injected Drugs	1.4	1.3	1.5	1.1	1.7	1.4	0.8	1.1	1.9 ▲
MDMA	1.0	1.0	1.1	0.7	0.8	1.0	1.8 ▲	0.8	1.3
Rohypnol	0.6	0.7	0.7	0.3	0.4	0.5	0.8	0.6	1.0 ▲
GHB	0.4	0.5	0.4	0.4	0.2	0.4	0.6	0.2	0.4
Over the counter drugs	6.5	6.0	5.9	5.6	5.5	6.4	7.0	5.6	9.1 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 7th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	10.8	9.6	11.2	9.2	12.9 ▲	8.3 ▼	16.7 ▲	8.7 ▼	15.1 ▲
Cigars	6.2	5.6	6.3	5.6	7.3 ▲	4.5 ▼	9.2 ▲	5.4	9.5 ▲
Pipe	2.6	3.0	2.3	1.9	3.1	1.8 ▼	4.6 ▲	2.1	3.7 ▲
Smokeless Tobacco	3.3	2.1 ▼	2.5 ▼	1.5 ▼	5.2 ▲	2.2 ▼	5.0 ▲	4.0	6.0 ▲
Alcohol	24.8	26.2	23.1	21.4 ▼	27.2 ▲	22.7 ▼	25.3	25.7	29.7 ▲
Marijuana	6.3	6.9	5.9	5.5	5.9	5.7	10.7 ▲	2.6 ▼	8.1 ▲
Cocaine	1.0	1.2	1.3	1.2	0.6	0.8	1.2	0.7	1.4 ▲
Crack	0.9	0.9	1.0	1.2	0.8	0.8	1.2	0.4	1.3 ▲
Inhalants	5.4	5.9	4.7	4.4	5.4	4.8	7.7 ▲	5.0	6.8 ▲
Amphetamines	1.0	0.7	1.1	0.5	1.2	0.7	1.4	0.8	1.4 ▲
Methamphetamines	0.7	0.7	0.6	0.9	0.7	0.6	0.8	0.5	1.2 ▲
Ritalin	1.6	1.4	1.5	0.8 ▼	1.7	1.4	2.1	1.4	2.1 ▲
Tranquilizers	4.2	4.2	3.6	3.3	3.8	3.8	6.4 ▲	3.3	5.9 ▲
Narcotics	1.6	1.2	1.3	0.8 ▼	1.8	1.4	2.5 ▲	1.7	2.8 ▲
Psychedelics <sup>#</sup>	1.0	1.1	0.8	0.7	0.6	1.1	1.4	1.1	1.2
LSD	0.7	0.7	0.6	0.3	0.5	0.8	0.9	0.8	0.9
Others	0.6	0.9	0.5	0.7	0.4	0.5	0.8	0.6	0.7
Heroin	0.7	0.8	0.7	0.8	0.5	0.6	1.0	0.5	0.8
Steroids	0.9	0.6	0.8	0.7	0.6	0.8	1.8 ▲	1.0	1.5 ▲
Injected Drugs	1.0	1.0	1.2	0.8	1.0	1.0	0.8	0.8	1.3
MDMA	0.8	0.8	0.8	0.7	0.6	0.7	1.5 ▲	0.8	1.2
Rohypnol	0.5	0.4	0.5	0.3	0.3	0.4	0.8	0.5	0.7 ▲
GHB	0.3	0.4	0.4	0.4	0.2	0.3	0.3	0.3	0.4
Over the counter drugs	5.1	4.1	4.6	4.3	4.6	5.0	5.3	4.5	7.4 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).  
▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 7th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	6.2	5.6	6.8	4.7 ▼	7.4 ▲	4.4 ▼	9.8 ▲	4.6 ▼	9.9 ▲
Cigars	3.6	3.7	3.6	2.7	4.4	2.5 ▼	6.1 ▲	3.1	5.6 ▲
Pipe	1.5	1.9	1.5	1.0	1.8 ▲	1.0 ▼	2.7 ▲	1.1	2.3 ▲
Smokeless Tobacco	2.0	1.1 ▼	1.9	1.2 ▼	3.1 ▲	1.2 ▼	2.9 ▲	2.0	3.8 ▲
Alcohol	12.2	13.6 ▲	11.7	10.0 ▼	13.4	10.4 ▼	13.1	12.9	15.8 ▲
Marijuana	3.9	4.1	3.5	3.1	3.9	3.3 ▼	6.6 ▲	1.5 ▼	5.7 ▲
Cocaine	0.7	0.6	0.9	0.9	0.5	0.5	1.0	0.3	0.8
Crack	0.6	0.6	0.6	0.6	0.4	0.6	0.5	0.3	0.6
Inhalants	3.1	3.5	2.5	2.2	3.2	2.6	4.8 ▲	3.0	3.9 ▲
Amphetamines	0.6	0.5	0.6	0.2	0.8	0.5	0.8	0.5	1.0 ▲
Methamphetamines	0.5	0.5	0.5	0.4	0.5	0.4	0.5	0.4	0.7
Ritalin	1.0	0.9	1.1	0.5	1.0	0.9	1.4	0.8	1.2
Tranquilizers	2.5	2.4	1.9	2.0	2.6	2.1	3.5 ▲	2.4	3.8 ▲
Narcotics	1.0	0.6	0.7	0.4 ▼	1.1	0.8	1.4	1.3	1.7 ▲
Psychedelics <sup>#</sup>	0.6	0.6	0.5	0.4	0.5	0.6	0.8	0.5	0.7
LSD	0.5	0.4	0.3	0.1	0.3	0.6	0.6	0.5	0.5
Others	0.3	0.4	0.3	0.4	0.4	0.3	0.4	0.2	0.4
Heroin	0.4	0.5	0.5	0.3	0.3	0.4	0.4	0.5	0.6
Steroids	0.6	0.7	0.4	0.4	0.6	0.5	1.3 ▲	0.6	1.0 ▲
Injected Drugs	0.6	0.6	0.7	0.4	0.6	0.5	0.3	0.3	0.8
MDMA	0.6	0.3	0.7	0.4	0.5	0.5	1.3 ▲	0.5	0.6
Rohypnol	0.3	0.3	0.3	0.1	0.3	0.2	0.5	0.3	0.3
GHB	0.2	0.3	0.1	0.2	0.1	0.2	0.3	0.2	0.2
Over the counter drugs	3.4	2.6	3.1	3.3	3.4	3.2	3.4	3.3	5.1 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 7th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
<b>CIGARETTES</b>									
Daily Use	2.7	2.5	2.9	2.2	2.6	2.0 ▼	3.6 ▲	1.9	4.8 ▲
1/2+ Pack per Day	1.3	1.0	1.2	1.0	1.3	0.9 ▼	1.6	1.1	2.4 ▲
<b>CIGARS</b>									
	0.7	0.4	0.6	0.6	1.1	0.5 ▼	1.4 ▲	0.5	1.3 ▲
<b>SMOKELESS TOBACCO</b>									
	0.4	0.2	0.3	0.1	0.5	0.1 ▼	0.4	0.3	1.3 ▲
<b>ALCOHOL</b>									
Daily Use	0.9	0.9	0.8	0.5	1.2	0.6 ▼	1.1	0.4	1.6 ▲
Binge Drinking	8.3	9.3	7.5	7.7	9.9 ▲	7.0 ▼	8.6	7.9	10.9 ▲
<b>MARIJUANA</b>									
	0.8	0.8	0.7	0.2	1.0	0.7	1.4 ▲	0.2	1.5 ▲

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ( $p<.05$ ).  
 ▼ indicates that local prevalence rates are significantly lower than state rates ( $p<.05$ ).

**Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 8th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	28.1	27.7	30.3 ▲	23.4 ▼	29.7	25.0 ▼	36.5 ▲	24.6 ▼	33.0 ▲
Cigars	14.7	13.7	15.5	11.8 ▼	15.4	13.2 ▼	18.7 ▲	12.1 ▼	18.7 ▲
Pipe	5.8	7.8 ▲	5.4	3.9 ▼	5.9	5.2 ▼	8.0 ▲	4.3 ▼	7.4 ▲
Smokeless Tobacco	7.2	5.5 ▼	4.7 ▼	4.6 ▼	10.3 ▲	5.1 ▼	11.9 ▲	6.7	11.8 ▲
Alcohol	45.4	47.4 ▲	46.2	40.0 ▼	45.8	42.2 ▼	48.7 ▲	43.1 ▼	51.7 ▲
Marijuana	16.1	17.0	16.2	12.1 ▼	14.6 ▼	16.8	20.4 ▲	12.3 ▼	18.2 ▲
Cocaine	2.4	3.3 ▲	2.5	2.4	2.1	2.3	3.0	1.7 ▼	2.7
Crack	2.0	2.5	1.9	1.7	1.9	2.1	2.2	1.5 ▼	2.3
Inhalants	10.5	12.7 ▲	9.8	7.8 ▼	10.3	9.3 ▼	12.5 ▲	11.4	12.2 ▲
Amphetamines	3.0	3.1	2.4 ▼	2.0 ▼	3.0	2.9	4.1 ▲	2.8	3.7 ▲
Methamphetamines	1.6	2.1	1.2	1.6	1.6	1.6	2.1	1.2	2.1 ▲
Ritalin	4.5	5.3	4.0	3.7	4.1	4.2	5.6 ▲	4.7	5.0
Tranquilizers	9.1	9.3	8.1 ▼	7.2 ▼	9.3	8.9	10.8 ▲	8.5	10.4 ▲
Narcotics	5.0	4.8	3.7 ▼	3.8 ▼	5.0	5.1	6.9 ▲	4.1 ▼	6.6 ▲
Psychedelics <sup>#</sup>	2.6	3.3 ▲	1.8 ▼	1.8 ▼	2.1	2.9	3.7 ▲	2.0 ▼	3.2 ▲
LSD	2.2	2.7	1.6 ▼	1.3 ▼	1.8	2.5	3.1 ▲	1.5 ▼	2.7 ▲
Others	1.3	1.8 ▲	0.8 ▼	0.9	1.0	1.3	1.8	1.2	1.7 ▲
Heroin	1.3	1.9 ▲	1.0	1.2	1.1	1.5	1.7	1.0	1.5
Steroids	1.5	1.7	1.2	0.9 ▼	1.7	1.4	1.6	1.6	1.8
Injected Drugs	1.7	2.1	1.5	1.2	1.6	1.8	1.7	1.4	1.9
MDMA	2.3	2.9 ▲	1.9	1.6	1.9	2.4	2.6	1.7 ▼	2.8 ▲
Rohypnol	1.1	1.3	0.8	0.6	1.0	1.1	1.6 ▲	1.0	1.2
GHB	0.7	1.1 ▲	0.5	0.2	0.8	0.7	0.7	0.5	0.7
Over the counter drugs	10.2	11.0	9.5	8.4 ▼	9.8	10.2	12.1 ▲	9.2 ▼	11.6 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

## Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 2007

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	18.0	18.3	18.1	15.0 ▼	19.1	15.9 ▼	22.5 ▲	15.2 ▼	22.4 ▲
Cigars	11.0	10.6	11.5	8.7 ▼	11.2	9.8 ▼	13.6 ▲	8.8 ▼	14.4 ▲
Pipe	5.0	6.7 ▲	4.7	3.2 ▼	5.0	4.7	6.7 ▲	3.6 ▼	6.1 ▲
Smokeless Tobacco	5.8	4.7 ▼	3.8 ▼	3.6 ▼	8.1 ▲	4.0 ▼	9.5 ▲	5.6	9.3 ▲
Alcohol	36.6	38.7 ▲	36.1	31.8 ▼	37.3	33.4 ▼	40.0 ▲	34.9 ▼	43.0 ▲
Marijuana	13.1	14.5 ▲	13.7	9.1 ▼	11.6 ▼	13.5	16.1 ▲	9.6 ▼	15.3 ▲
Cocaine	1.8	2.6 ▲	1.8	1.7	1.6	1.7	2.4	1.4	2.1
Crack	1.5	1.8	1.5	1.1	1.6	1.4	1.7	1.3	1.7
Inhalants	7.1	8.1 ▲	6.9	5.1 ▼	7.2	5.8 ▼	8.2	7.9	9.0 ▲
Amphetamines	2.3	2.5	1.9	1.6	2.3	2.0	3.0	2.2	3.0 ▲
Methamphetamines	1.2	1.7 ▲	0.9	0.8	1.0	1.3	1.7	1.0	1.3
Ritalin	3.3	3.9	3.1	2.8	3.1	2.8 ▼	4.1	3.7	3.7
Tranquilizers	6.7	6.3	6.1	5.4 ▼	6.7	6.7	8.0	6.5	7.8 ▲
Narcotics	3.6	3.5	2.6 ▼	2.8	3.4	3.6	5.2 ▲	2.8 ▼	4.8 ▲
Psychedelics <sup>#</sup>	2.0	2.6 ▲	1.4 ▼	1.4	1.8	2.1	2.9 ▲	1.6	2.4
LSD	1.6	2.0	1.1 ▼	0.9 ▼	1.4	1.7	2.6 ▲	1.2	2.0
Others	1.0	1.4 ▲	0.7	0.8	0.8	0.9	1.4	0.9	1.2
Heroin	0.9	1.5 ▲	0.7	0.8	0.8	1.0	1.0	0.8	1.0
Steroids	1.2	1.4	0.8	0.9	1.3	1.0	1.3	1.2	1.5
Injected Drugs	1.3	1.7	1.1	0.8	1.0	1.3	1.5	1.2	1.4
MDMA	1.7	2.3 ▲	1.5	1.3	1.4	1.7	2.1	1.2 ▼	2.1
Rohypnol	0.8	1.2 ▲	0.6	0.5	0.7	0.7	1.3 ▲	0.7	0.9
GHB	0.5	0.7	0.4	0.3	0.6	0.5	0.6	0.5	0.6
Over the counter drugs	8.0	8.7	7.5	6.8	7.4	7.9	9.8 ▲	7.3	9.2 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).  
 ▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 8th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	10.8	11.1	10.3	8.7 ▼	11.3	9.5 ▼	14.5 ▲	9.2 ▼	14.1 ▲
Cigars	6.4	5.8	6.2	5.0 ▼	6.7	5.7 ▼	8.3 ▲	5.4 ▼	8.4 ▲
Pipe	3.1	4.5 ▲	2.8	1.8 ▼	3.2	2.7	3.9	2.2 ▼	4.0 ▲
Smokeless Tobacco	3.3	2.9	2.0 ▼	2.0 ▼	4.6 ▲	2.2 ▼	5.7 ▲	3.0	5.7 ▲
Alcohol	19.9	22.8 ▲	20.2	15.3 ▼	19.3	18.1 ▼	20.8	18.6 ▼	24.1 ▲
Marijuana	8.3	9.1	8.6	5.6 ▼	6.7 ▼	8.6	9.9 ▲	6.3 ▼	10.1 ▲
Cocaine	1.1	1.8 ▲	1.0	0.8	0.9	1.0	1.2	0.8	1.2
Crack	0.9	1.3 ▲	0.8	0.7	0.8	0.9	0.8	0.8	0.9
Inhalants	3.7	4.1	3.5	2.5 ▼	3.5	3.2 ▼	4.3	4.4	4.7 ▲
Amphetamines	1.4	1.5	1.3	0.8 ▼	1.5	1.2	1.6	1.2	1.6
Methamphetamines	0.7	0.9	0.5	0.6	0.6	0.7	1.0	0.7	0.6
Ritalin	1.9	2.5 ▲	1.6	1.5	1.8	1.8	2.2	1.9	2.2
Tranquilizers	3.9	3.8	3.8	2.7 ▼	3.9	3.8	4.2	3.7	4.6 ▲
Narcotics	2.1	2.1	1.5 ▼	1.4	1.8	2.2	2.9 ▲	1.6 ▼	2.7 ▲
Psychedelics <sup>#</sup>	1.1	1.5 ▲	0.7 ▼	0.7	1.0	1.1	1.7	1.2	1.4
LSD	0.9	1.2	0.6 ▼	0.6	0.8	0.9	1.6 ▲	0.8	1.2
Others	0.6	0.9 ▲	0.4	0.3	0.5	0.5	0.5	0.8	0.7
Heroin	0.6	1.1 ▲	0.4	0.5	0.6	0.5	0.5	0.6	0.8
Steroids	0.7	1.0	0.5	0.5	0.8	0.6	1.0	0.7	0.9
Injected Drugs	0.8	1.2 ▲	0.8	0.6	0.8	0.8	1.2	0.8	0.8
MDMA	0.9	1.1	0.7	0.6	0.8	1.0	1.1	0.9	1.1
Rohypnol	0.5	0.7	0.3	0.3	0.4	0.4	0.9 ▲	0.6	0.6
GHB	0.3	0.6	0.3	0.2	0.3	0.3	0.6	0.4	0.3
Over the counter drugs	5.2	5.5	5.2	3.9 ▼	4.8	5.1	6.2	4.8	6.1 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 8th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
<b>CIGARETTES</b>									
Daily Use	5.5	5.5	4.6 ▼	4.3 ▼	5.7	5.1	8.0 ▲	4.5 ▼	7.4 ▲
1/2+ Pack per Day	2.7	2.1	2.3	1.9 ▼	3.1	2.6	4.1 ▲	2.0 ▼	3.5 ▲
<b>CIGARS</b>									
	1.6	1.1	1.4	0.8 ▼	1.6	1.6	2.1	1.0 ▼	2.3 ▲
<b>SMOKELESS TOBACCO</b>									
	0.8	0.6	0.4 ▼	0.4	1.2 ▲	0.6	1.4 ▲	0.5	1.4 ▲
<b>ALCOHOL</b>									
Daily Use	1.8	2.1	1.6	1.3	1.6	1.7	2.4	1.0 ▼	2.5 ▲
Binge Drinking	13.2	16.6 ▲	13.9	9.3 ▼	13.4	11.6 ▼	15.0	11.4 ▼	16.1 ▲
<b>MARIJUANA</b>									
	2.0	2.3	1.9	1.2 ▼	1.6	2.3	3.0 ▲	1.1 ▼	2.5 ▲

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ( $p<.05$ ).  
 ▼ indicates that local prevalence rates are significantly lower than state rates ( $p<.05$ ).

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 2007

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	33.5	30.9 ▼	33.2	28.8 ▼	37.3 ▲	31.1 ▼	42.2 ▲	31.3	37.6 ▲
Cigars	19.4	16.0 ▼	20.5	16.2 ▼	21.7 ▲	17.9 ▼	22.0 ▲	19.6	22.9 ▲
Pipe	9.1	8.9	8.9	7.3 ▼	10.6 ▲	9.0	10.3	6.8 ▼	10.3 ▲
Smokeless Tobacco	10.8	7.0 ▼	9.4 ▼	7.1 ▼	15.0 ▲	8.4 ▼	13.6 ▲	11.9	17.8 ▲
Alcohol	51.7	51.4	50.1	48.3 ▼	54.7 ▲	49.8 ▼	55.5 ▲	53.3	54.9 ▲
Marijuana	21.2	20.5	19.7	17.5 ▼	21.1	22.7 ▲	27.3 ▲	13.3 ▼	21.9
Cocaine	3.7	3.7	4.4	2.6 ▼	3.3	3.9	4.6	2.6	3.9
Crack	2.4	2.1	3.4 ▲	1.4 ▼	2.0	2.6	3.4 ▲	1.7	1.8
Inhalants	9.8	9.5	10.1	7.1 ▼	9.5	9.6	13.7 ▲	9.9	10.3
Amphetamines	5.3	5.3	5.7	3.5 ▼	6.1	5.3	6.5 ▲	4.4	5.5
Methamphetamines	2.2	1.6	2.4	0.9 ▼	2.0	2.4	2.9	1.9	2.5
Ritalin	7.3	6.3	8.0	5.4 ▼	8.0	7.7	9.0 ▲	5.1 ▼	7.4
Tranquilizers	10.9	9.5	11.1	8.1 ▼	11.1	11.4	13.2 ▲	8.5 ▼	11.9
Narcotics	7.6	6.2 ▼	7.4	4.3 ▼	8.8 ▲	7.7	10.5 ▲	5.6 ▼	8.9 ▲
Psychedelics <sup>#</sup>	3.6	3.1	3.0	2.7 ▼	3.4	4.3 ▲	4.4	3.1	3.5
LSD	2.9	2.5	2.2 ▼	2.3	2.9	3.5 ▲	3.6	2.6	2.8
Others	1.8	1.5	1.9	1.1 ▼	1.3	2.1	2.6 ▲	1.5	1.8
Heroin	1.5	1.5	1.9	0.8 ▼	1.2	1.8	1.9	1.5	1.1
Steroids	1.7	1.3	2.1	0.9 ▼	2.0	1.6	2.1	1.9	1.6
Injected Drugs	1.7	1.3	2.3	1.3	1.4	1.8	2.1	1.8	1.7
MDMA	3.2	3.3	3.1	2.2 ▼	2.7	3.8 ▲	3.7	2.4	2.9
Rohypnol	1.4	1.3	1.3	0.9 ▼	1.4	1.6	2.3 ▲	1.5	1.1
GHB	0.9	0.8	0.8	0.8	0.6	1.1	1.5 ▲	0.9	0.6
Over the counter drugs	12.3	10.8	11.4	9.4 ▼	12.4	13.0	15.9 ▲	9.1 ▼	13.6 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

## Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 2007

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	22.6	19.0 ▼	22.1	18.7 ▼	24.8 ▲	21.4 ▼	29.1 ▲	22.5	26.1 ▲
Cigars	14.7	12.3 ▼	14.6	12.2 ▼	15.9	13.9	17.2 ▲	15.4	17.1 ▲
Pipe	7.6	7.7	7.4	5.7 ▼	8.9 ▲	7.4	8.3	5.6 ▼	8.8 ▲
Smokeless Tobacco	8.7	5.6 ▼	7.7	4.9 ▼	12.0 ▲	7.0 ▼	10.0	10.2	14.4 ▲
Alcohol	43.4	43.2	41.6	40.1 ▼	46.4 ▲	41.5 ▼	45.7	46.8	47.0 ▲
Marijuana	17.0	16.3	15.7	13.3 ▼	16.6	18.5 ▲	21.9 ▲	10.5 ▼	17.8
Cocaine	2.6	2.5	3.2	1.7 ▼	2.3	2.6	3.2	1.7	2.7
Crack	1.6	1.2	2.1 ▲	0.9 ▼	1.3	1.7	2.2 ▲	1.0	1.6
Inhalants	6.1	5.8	6.1	4.0 ▼	6.5	6.0	7.5 ▲	6.1	6.6
Amphetamines	3.7	3.4	4.1	2.4 ▼	4.4	3.7	4.6	3.1	3.9
Methamphetamines	1.4	1.0	1.8	0.7 ▼	1.1	1.6	1.8	1.1	1.8
Ritalin	5.3	4.1 ▼	6.0	3.9 ▼	5.9	5.7	6.4	4.0 ▼	5.2
Tranquilizers	7.9	6.7	7.9	5.8 ▼	8.2	8.3	9.2	6.2 ▼	9.1 ▲
Narcotics	5.3	4.0 ▼	5.3	2.8 ▼	6.3 ▲	5.6	7.0 ▲	3.5 ▼	6.6 ▲
Psychedelics <sup>#</sup>	2.5	2.1	2.3	2.0	2.0	3.1 ▲	2.9	2.0	2.6
LSD	2.0	1.6	1.6	1.7	2.4 ▲	2.3	1.7	2.0	
Others	1.3	1.0	1.5	1.1	0.8 ▼	1.5	1.7	0.9	1.1
Heroin	1.1	0.9	1.4	0.5 ▼	0.8	1.3	1.5	1.0	0.8
Steroids	1.2	1.0	1.6	0.8	1.3	1.0	1.7	1.1	1.2
Injected Drugs	1.4	1.0	1.8	0.9	1.1	1.6	1.7	1.4	1.3
MDMA	2.2	2.0	2.3	1.6	1.5 ▼	2.5	2.7	1.5	2.2
Rohypnol	1.0	0.8	1.0	0.6	0.9	1.2	1.7 ▲	0.7	0.9
GHB	0.6	0.5	0.7	0.5	0.3	0.7	1.1 ▲	0.5	0.3
Over the counter drugs	9.2	7.9	8.7	6.7 ▼	9.2	9.7	11.6 ▲	7.3 ▼	10.5 ▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).  
 ▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

## Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 2007

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	14.7	12.1 ▼	14.6	11.7 ▼	17.7 ▲	12.9 ▼	19.6 ▲	14.8	18.3 ▲
Cigars	8.7	7.1 ▲	8.5	6.8 ▼	10.4 ▲	8.1	10.2	8.4	10.6 ▲
Pipe	4.4	4.8	4.7	3.3 ▼	5.5 ▲	4.1	4.8	2.9 ▼	4.7
Smokeless Tobacco	5.3	3.6 ▼	4.7	3.0 ▼	7.3 ▲	4.0 ▼	6.2	6.3	8.8 ▲
Alcohol	24.8	24.3	23.9	21.5 ▼	25.2	23.2 ▼	25.9	28.9 ▲	29.3 ▲
Marijuana	10.3	10.1	9.4	8.0 ▼	10.5	11.0	14.1 ▲	5.7 ▼	11.1
Cocaine	1.4	1.4	1.8	1.1	1.1	1.5	1.8	0.8	1.3
Crack	0.9	0.9	1.3 ▲	0.4 ▼	0.6	1.0	1.0	0.7	0.7
Inhalants	2.9	2.7	3.2	1.7 ▼	3.1	3.1	3.9 ▲	2.6	2.9
Amphetamines	2.0	2.3	2.2	1.3 ▼	2.2	2.0	2.8	1.7	1.6
Methamphetamines	0.8	0.8	0.9	0.4 ▼	0.7	0.8	1.3 ▲	0.9	0.9
Ritalin	2.8	2.3	3.3	2.0 ▼	3.2	2.9	3.7 ▲	1.9 ▼	2.8
Tranquilizers	4.4	4.0	4.3	2.7 ▼	5.0	4.7	5.2	3.7 ▼	4.6
Narcotics	2.9	2.3	2.9	1.3 ▼	3.5	3.3	3.9 ▲	2.0 ▼	3.1
Psychedelics <sup>#</sup>	1.3	1.3	1.3	1.1	1.2	1.5	1.8	1.1	1.3
LSD	1.1	0.9	1.0	1.0	0.9	1.3	1.4	0.9	1.0
Others	0.7	0.7	0.7	0.5	0.7	0.7	1.0	0.5	0.5
Heroin	0.7	0.7	1.1 ▲	0.3 ▼	0.7	0.8	0.7	0.8	0.5
Steroids	0.8	0.7	1.1	0.6	1.0	0.6	1.4 ▲	0.9	0.9
Injected Drugs	0.9	0.7	1.1	0.7	0.8	0.9	0.8	1.5	0.8
MDMA	1.2	1.3	1.1	0.9	0.9	1.3	1.7	1.1	1.0
Rohypnol	0.6	0.4	0.5	0.4	0.6	0.6	0.9	0.4	0.5
GHB	0.4	0.4	0.4	0.3	0.3	0.4	0.9 ▲	0.3	0.3
Over the counter drugs	5.6	4.9	5.3	4.0 ▼	5.9	5.7	7.3 ▲	4.9	6.3

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 9th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
<b>CIGARETTES</b>									
Daily Use	8.1	6.1 ▼	8.5	6.4 ▼	9.8 ▲	7.2 ▼	11.6 ▲	6.3 ▼	10.0 ▲
1/2+ Pack per Day	4.4	3.0 ▼	4.8	3.4 ▼	5.3 ▲	3.6 ▼	6.4 ▲	3.5	5.7 ▲
<b>CIGARS</b>									
	2.3	1.2 ▼	2.3	2.0	3.4 ▲	1.9	3.1 ▲	1.5	3.0 ▲
<b>SMOKELESS TOBACCO</b>									
	1.6	0.8 ▼	1.2	0.9 ▼	2.2 ▲	1.1 ▼	1.8	2.0	3.6 ▲
<b>ALCOHOL</b>									
Daily Use	2.3	1.8	2.5	1.9	2.7	2.1	2.4	2.0	3.2 ▲
Binge Drinking	16.9	16.5	15.9	14.0 ▼	17.3	16.1	18.2	20.7 ▲	19.3 ▲
<b>MARIJUANA</b>									
	2.7	2.4	2.2	2.3	3.1	2.6	4.0 ▲	1.1 ▼	3.7 ▲

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ( $p<.05$ ).  
 ▼ indicates that local prevalence rates are significantly lower than state rates ( $p<.05$ ).

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 10th Grade Students: 2007

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	40.6	39.9	39.5	34.3	44.7	38.8	46.4	38.8	43.9
Cigars	26.7	25.4	26.3	22.3	29.4	25.1	31.4	27.0	28.3
Pipe	13.3	20.1	12.7	10.1	14.4	13.0	13.8	11.0	13.3
Smokeless Tobacco	15.0	12.3	10.8	10.8	20.4	11.8	20.7	15.6	19.9
Alcohol	61.0	62.8	59.5	56.4	63.2	57.7	63.4	62.1	64.4
Marijuana	29.9	31.5	29.6	23.9	29.0	32.0	35.8	27.0	30.3
Cocaine	5.8	6.0	5.4	5.2	6.0	5.8	7.1	5.1	6.3
Crack	3.1	2.7	3.2	2.3	3.3	3.0	4.3	2.7	3.5
Inhalants	10.6	10.7	9.9	8.8	11.7	9.7	13.6	12.2	10.3
Amphetamines	8.2	7.6	8.3	5.2	9.0	8.1	11.1	8.2	8.5
Methamphetamines	3.0	2.3	3.3	2.3	3.1	2.9	3.0	3.2	3.1
Ritalin	10.6	10.7	11.4	8.0	10.0	10.1	12.9	12.0	10.0
Tranquilizers	13.6	12.9	12.3	10.7	15.1	13.2	16.1	14.1	14.6
Narcotics	10.9	9.1	9.4	8.2	11.5	11.5	14.1	10.2	12.3
Psychedelics <sup>#</sup>	4.8	4.3	4.2	3.5	4.7	5.6	4.5	4.6	5.4
LSD	3.9	3.4	3.3	2.9	3.9	4.7	3.5	3.5	4.4
Others	2.6	2.4	2.4	1.9	2.5	3.0	2.9	2.8	2.7
Heroin	2.0	1.7	2.1	1.5	2.2	2.2	2.2	1.6	2.0
Steroids	2.1	1.5	2.3	1.4	2.4	2.1	2.5	1.9	2.2
Injected Drugs	2.0	1.8	2.1	1.3	2.3	2.0	2.6	1.9	2.1
MDMA	4.3	5.3	4.2	3.0	3.8	5.0	4.6	3.6	4.6
Rohypnol	1.7	1.3	1.7	0.8	2.5	1.8	2.2	1.8	1.6
GHB	1.2	0.6	1.0	0.6	1.6	1.4	1.5	1.2	1.0
Over the counter drugs	14.4	13.8	13.3	11.5	15.2	14.2	16.7	14.8	15.3

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 10th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	28.5	27.7	27.1	23.7	31.7	26.4	32.6	28.2	31.8
Cigars	20.7	19.4	20.2	17.0	22.6	19.2	23.7	21.8	22.2
Pipe	11.1	17.6	▲	10.1	8.9	▼	12.3	10.5	11.5
Smokeless Tobacco	12.0	9.6	▼	8.9	9.0	▼	16.0	▲	16.6
Alcohol	51.7	54.4	▲	50.1	46.3	▼	54.0	48.5	▼
Marijuana	23.5	25.1	23.1	18.4	▼	22.6	24.8	▲	28.0
Cocaine	4.0	4.3	3.7	3.3	4.3	4.0	5.1	3.5	4.3
Crack	2.0	2.0	1.9	1.2	▼	2.4	2.0	2.9	▲
Inhalants	5.8	6.1	5.4	4.4	▼	6.5	5.1	▼	7.5
Amphetamines	5.7	5.6	5.2	3.0	▼	6.3	5.8	8.4	6.1
Methamphetamines	1.9	1.5	1.9	1.0	▼	2.2	1.9	2.3	1.9
Ritalin	7.4	8.0	8.0	5.5	▼	6.8	6.8	9.1	8.5
Tranquilizers	9.5	8.8	8.4	▼	6.7	▼	11.1	▲	9.1
Narcotics	7.5	6.4	▼	6.0	▼	5.5	▼	8.3	7.6
Psychedelics <sup>#</sup>	3.2	2.9	3.2	2.1	▼	3.1	3.8	▲	3.1
LSD	2.4	2.2	2.3	1.5	▼	2.5	3.0	▲	2.4
Others	1.8	1.7	2.0	1.1	▼	1.8	2.1	2.0	1.7
Heroin	1.3	1.1	1.4	0.9	1.3	1.6	1.5	0.9	▼
Steroids	1.3	1.0	1.4	0.9	1.5	1.4	1.7	1.2	1.3
Injected Drugs	1.5	1.3	1.5	1.0	1.6	1.7	1.9	1.3	1.4
MDMA	2.9	3.7	▲	3.0	1.9	▼	2.5	3.5	3.3
Rohypnol	1.1	0.9	1.2	0.3	▼	1.7	▲	1.3	▲
GHB	0.7	0.5	0.7	0.3	▼	1.0	1.0	0.7	0.7
Over the counter drugs	10.3	10.4	8.9	▼	8.5	▼	11.2	10.1	11.7

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).  
 ▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 10th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	19.3	18.0	18.3	15.7	21.4	17.6	22.6	19.5	21.9
Cigars	12.3	11.7	11.9	10.3	12.6	11.1	15.1	13.4	13.7
Pipe	6.8	11.0	▲	6.2	5.6	7.5	6.4	7.0	5.5
Smokeless Tobacco	7.3	6.2	▼	5.4	5.1	10.0	▲	5.9	9.5
Alcohol	31.1	33.8	▲	29.2	▼	26.2	32.3	27.9	▼
Marijuana	14.4	16.8	▲	13.2	11.8	▼	13.6	15.0	17.2
Cocaine	1.9	2.1	1.8	1.5	2.3	2.2	1.9	1.5	1.9
Crack	1.0	0.8	1.2	0.8	1.3	1.1	1.3	0.8	1.0
Inhalants	2.5	2.3	2.2	2.1	3.0	2.2	3.3	▲	2.4
Amphetamines	2.9	2.2	▼	2.6	2.0	▼	3.6	▲	4.3
Methamphetamines	1.0	0.6	1.0	0.5	▼	1.2	1.2	1.5	0.9
Ritalin	3.7	3.6	4.7	▲	2.9	3.2	3.5	4.2	3.9
Tranquilizers	5.0	4.6	4.0	▼	3.8	▼	6.0	▲	6.7
Narcotics	3.9	3.2	2.9	▼	3.0	4.3	4.2	5.9	▲
Psychedelics <sup>#</sup>	1.6	1.3	1.4	0.8	▼	1.7	2.0	▲	1.6
LSD	1.2	0.9	1.1	0.7	▼	1.3	1.6	▲	1.1
Others	0.9	0.8	0.8	0.4	▼	1.1	1.2	1.3	0.8
Heroin	0.8	0.5	0.9	0.5	0.9	1.0	1.0	0.5	0.7
Steroids	0.9	0.8	1.1	0.6	1.2	0.9	1.1	0.8	0.8
Injected Drugs	1.0	1.0	1.0	0.6	0.9	1.3	1.5	0.9	0.8
MDMA	1.4	2.1	▲	1.3	1.0	1.3	1.7	1.7	1.0
Rohypnol	0.7	0.4	0.7	0.2	1.0	▲	0.8	1.1	▲
GHB	0.5	0.3	0.6	0.2	0.6	0.6	0.5	0.3	0.2
Over the counter drugs	5.9	5.6	5.3	4.8	▼	6.8	5.5	7.4	▲

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 10th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
<b>CIGARETTES</b>									
Daily Use	11.4	10.1	10.1 ▼	9.0 ▼	12.3	10.8	14.7 ▲	10.7	13.7 ▲
1/2+ Pack per Day	6.6	5.4 ▼	5.4 ▼	5.0 ▼	7.5 ▲	6.3	8.9 ▲	6.2	7.9 ▲
<b>CIGARS</b>									
	3.1	2.6	2.7	2.6	3.9 ▲	2.8	4.2 ▲	2.9	3.6
<b>SMOKELESS TOBACCO</b>									
	2.8	1.6 ▼	1.7 ▼	1.8 ▼	4.1 ▲	2.1 ▼	4.1 ▲	2.6	4.5 ▲
<b>ALCOHOL</b>									
Daily Use	3.4	4.4 ▲	2.8	2.7	4.3 ▲	2.9 ▼	4.5 ▲	3.4	3.6
Binge Drinking	21.7	24.5 ▲	20.3 ▼	17.6 ▼	23.2 ▲	19.8 ▼	23.1	23.5	22.9 ▲
<b>MARIJUANA</b>									
	4.6	6.6 ▲	3.8 ▼	4.2	4.0	4.3	6.7 ▲	3.7 ▼	5.2

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ( $p<.05$ ).  
 ▼ indicates that local prevalence rates are significantly lower than state rates ( $p<.05$ ).

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 11th Grade Students: 2007

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	44.6	43.4	45.1	40.4	47.8	41.3	48.1	46.7	49.2
Cigars	31.0	30.1	31.3	28.1	33.1	29.7	32.2	31.7	33.9
Pipe	16.2	21.0	14.2	13.9	16.8	16.5	15.4	12.1	17.6
Smokeless Tobacco	16.9	12.7	13.8	13.7	22.6	13.8	19.8	21.1	23.2
Alcohol	64.3	66.4	62.8	62.4	68.0	60.9	66.7	66.6	67.4
Marijuana	32.4	35.3	29.0	30.3	31.7	33.6	38.1	25.2	33.6
Cocaine	6.9	7.8	6.3	6.6	6.8	7.0	8.1	5.0	7.1
Crack	3.3	3.2	2.9	3.1	3.2	3.5	3.5	3.1	3.5
Inhalants	9.7	9.8	9.0	8.0	11.2	8.7	13.4	10.0	10.7
Amphetamines	9.1	8.8	8.4	7.5	10.8	9.1	12.2	7.1	9.3
Methamphetamines	3.3	3.0	2.9	3.4	3.6	3.1	3.2	3.8	4.1
Ritalin	10.7	11.2	12.0	10.4	12.0	10.9	11.2	8.0	8.8
Tranquilizers	13.4	12.8	12.8	10.5	14.9	13.7	16.1	11.2	14.0
Narcotics	11.7	10.4	9.9	9.7	11.9	12.8	15.6	9.1	12.8
Psychedelics <sup>#</sup>	5.4	4.6	4.5	6.5	4.9	6.4	5.3	4.2	5.2
LSD	4.4	3.4	3.8	5.1	3.7	5.3	4.2	3.6	4.5
Others	3.0	2.8	2.4	4.1	2.8	3.4	2.9	2.1	2.4
Heroin	2.0	2.3	2.1	2.5	1.5	2.2	1.9	2.0	1.7
Steroids	1.8	1.9	1.8	2.2	1.3	1.8	2.0	1.8	2.0
Injected Drugs	1.9	1.9	2.0	2.6	1.5	2.1	1.5	2.0	1.4
MDMA	4.8	5.6	4.4	4.7	3.6	5.7	4.6	3.6	4.7
Rohypnol	1.7	2.2	1.5	1.7	1.8	1.8	1.7	1.3	1.6
GHB	1.1	1.4	1.0	1.3	0.9	1.4	1.2	0.9	0.8
Over the counter drugs	13.5	14.1	12.5	11.8	15.6	14.0	15.7	9.1	13.3

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 11th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	32.4	29.6 ▼	32.9	30.3	34.7 ▲	29.7 ▼	34.2	36.9 ▲	35.8 ▲
Cigars	24.1	23.2	23.5	21.4 ▼	25.7	23.3	23.8	25.5	26.7 ▲
Pipe	13.3	19.0 ▲	11.1 ▼	11.4 ▼	13.3	13.7	12.0	9.9 ▼	14.3
Smokeless Tobacco	13.1	9.7 ▼	10.4 ▼	10.9 ▼	17.3 ▲	10.8 ▼	15.3 ▲	16.3 ▲	18.0 ▲
Alcohol	55.3	57.6	53.6	53.6	57.4	52.3 ▼	56.8	58.9 ▲	58.6 ▲
Marijuana	24.8	27.3 ▲	21.4 ▼	22.7	23.8	26.6 ▲	28.6 ▲	18.7 ▼	25.3
Cocaine	4.6	5.8 ▲	4.3	4.6	4.1	4.8	5.6	3.1 ▼	4.5
Crack	2.2	1.9	1.4 ▼	2.2	2.0	2.3	2.5	2.1	2.4
Inhalants	4.7	5.0	4.2	4.0	4.9	4.3	7.1 ▲	4.8	4.7
Amphetamines	5.8	5.4	5.0	4.5 ▼	6.6	6.1	8.4 ▲	4.1 ▼	5.9
Methamphetamines	2.0	2.0	1.6	2.0	2.0	1.9	2.3	1.9	2.5
Ritalin	6.9	6.8	7.6	7.3	7.1	7.2	7.7	4.6 ▼	5.9
Tranquilizers	8.9	8.2	7.9	6.4 ▼	10.4 ▲	9.2	11.0 ▲	7.1 ▼	9.7
Narcotics	7.7	6.8	6.1 ▼	6.4 ▼	7.8	8.3	11.1 ▲	6.1 ▼	8.6
Psychedelics <sup>#</sup>	3.6	3.2	2.8	4.7 ▲	2.9	4.0	3.6	2.2 ▼	3.7
LSD	2.6	1.9	2.0	3.5 ▲	1.9 ▼	3.1 ▲	3.0	1.7	2.7
Others	2.2	2.5	1.8	3.2 ▲	1.8	2.5	2.2	1.3 ▼	2.1
Heroin	1.4	1.5	1.4	1.7	1.1	1.5	1.3	1.2	1.2
Steroids	1.3	1.1	1.1	1.4	0.9	1.3	2.0	1.0	1.5
Injected Drugs	1.4	1.5	1.5	2.0	1.2	1.3	1.5	1.6	1.1
MDMA	3.0	3.4	2.2	3.0	2.0 ▼	3.8 ▲	2.8	2.0	2.9
Rohypnol	1.1	1.3	0.8	1.3	1.0	1.3	1.2	0.7	1.3
GHB	0.8	0.8	0.7	1.0	0.7	1.0	0.5	0.5	0.7
Over the counter drugs	9.0	9.6	7.9	8.9	10.4	9.1	10.2	6.1 ▼	9.2

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

## Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 11th Grade Students: 2007

DRUG	Indiana	Northwest	Central	North	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	22.5	20.3 ▼	22.3	21.1	25.2 ▲	20.1 ▼	25.8 ▲	25.6 ▲	24.7 ▲	24.7 ▲
Cigars	13.8	13.6	13.5	13.1	14.5	13.7	13.7	14.8	14.8	14.2
Pipe	7.4	11.2 ▲	6.1 ▼	6.8	7.1	7.8	6.2	4.1 ▼	8.0	
Smokeless Tobacco	8.1	5.9 ▼	6.5 ▼	6.3 ▼	10.6 ▲	6.6 ▼	9.8	10.1 ▲	11.3 ▲	
Alcohol	33.8	36.4	31.4 ▼	32.4	33.4	31.4 ▼	34.3	39.5 ▲	37.2 ▲	
Marijuana	14.6	16.7 ▲	12.5 ▼	13.7	13.7	15.5	18.3 ▲	9.3 ▼	14.9	
Cocaine	2.1	2.9	1.8	2.2	1.9	2.2	2.2	1.9	2.0	
Crack	1.1	1.1	0.7	1.4	1.0	1.2	0.8	1.2	1.2	
Inhalants	2.2	2.4	2.3	2.0	2.3	2.2	2.5	1.9	1.7	
Amphetamines	2.8	2.5	2.6	2.0	3.6 ▲	2.9	3.5	1.9	3.1	
Methamphetamines	1.1	1.2	1.1	1.4	1.0	0.9	1.1	1.1	1.3	
Ritalin	3.2	3.6	3.7	3.7	3.3	3.2	3.5	2.2	2.4 ▼	
Tranquilizers	4.3	3.9	4.1	2.8 ▼	5.0	4.4	5.5	3.2	4.8	
Narcotics	3.9	3.9	3.3	3.2	4.1	4.1	5.4 ▲	2.6 ▼	4.5	
Psychedelics <sup>#</sup>	1.7	1.4	1.6	2.2	1.3	1.8	1.6	1.1	2.0	
LSD	1.3	1.0	1.1	1.8	0.9	1.4	1.4	0.8	1.6	
Others	1.0	0.8	1.0	1.3	0.6	1.1	0.8	0.9	1.0	
Heroin	0.9	1.1	1.0	0.9	0.7	1.0	0.7	0.7	0.6	
Steroids	0.7	0.7	0.5	0.9	0.5	0.8	0.7	0.7	0.8	
Injected Drugs	1.1	1.2	1.3	1.7 ▲	1.0	1.0	0.8	1.1	0.7	
MDMA	1.3	1.7	0.8	1.4	0.9	1.5	1.0	1.1	1.4	
Rohypnol	0.7	0.8	0.6	0.8	0.5	0.7	0.6	0.5	0.8	
GHB	0.5	0.6	0.4	0.6	0.3	0.6	0.2	0.5	0.5	
Over the counter drugs	4.7	5.0	4.5	3.8	5.8 ▲	4.8	5.8	2.2 ▼	4.9	

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 11th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
<b>CIGARETTES</b>									
Daily Use	13.8	11.8 ▼	13.1	11.9 ▼	16.0 ▲	12.7	18.1 ▲	14.6	15.1
1/2+ Pack per Day	8.1	7.4	7.8	6.4 ▼	9.5 ▲	7.1 ▼	11.3 ▲	8.1	9.3
<b>CIGARS</b>									
	3.5	2.9	2.9	2.8	5.1	3.3 ▼	4.3	2.6	4.0
<b>SMOKELESS TOBACCO</b>									
	3.6	2.3 ▼	2.6 ▼	1.7 ▼	5.7 ▲	2.8 ▼	4.3	4.3	5.9 ▲
<b>ALCOHOL</b>									
Daily Use	3.6	4.7 ▲	2.9	2.2 ▼	3.3	3.2	4.5	4.8 ▲	4.3
Binge Drinking	23.2	25.4	22.2	21.2	22.3	21.8 ▼	23.6	28.9 ▲	24.5 ▲
<b>MARIJUANA</b>									
	4.6	5.1	3.4 ▼	3.5 ▼	4.6	5.1	7.0 ▲	2.4 ▼	5.2

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ( $p<.05$ ).  
 ▼ indicates that local prevalence rates are significantly lower than state rates ( $p<.05$ ).

## Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 12th Grade Students: 2007

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	48.4	46.5	46.6	43.6	51.6	45.2	53.0	48.6	53.3
Cigars	37.9	36.8	36.5	35.0	38.8	38.4	34.5	37.1	40.9
Pipe	20.0	29.9	▲	16.4	▼	15.6	22.7	▲	16.0
Smokeless Tobacco	19.4	16.4	▼	14.2	▼	16.8	24.7	▲	16.4
Alcohol	69.2	71.7	66.6	▼	64.4	▼	70.1	65.9	▼
Marijuana	36.5	38.7	35.2	34.4	34.2	▼	37.3	41.5	▲
Cocaine	7.8	9.0	7.1	7.7	8.5	7.7	8.5	7.1	8.0
Crack	3.2	3.6	3.2	2.2	▼	4.6	▲	3.2	4.1
Inhalants	8.5	8.2	8.2	7.9	9.0	7.5	▼	10.4	9.4
Amphetamines	9.0	8.2	7.8	6.8	▼	10.5	▲	8.4	12.3
Methamphetamines	3.4	2.6	2.6	2.9	5.5	▲	2.7	▼	2.4
Ritalin	11.3	11.6	11.5	10.8	10.9	11.1	10.8	12.2	11.3
Tranquilizers	12.9	13.3	10.3	▼	9.2	▼	15.1	▲	13.3
Narcotics	12.1	11.4	9.3	▼	9.7	▼	13.2	12.6	15.0
Psychedelics <sup>#</sup>	6.1	7.3	5.0	▼	5.8	6.5	6.6	5.7	5.6
LSD	4.5	4.9	3.5	▼	4.3	5.2	5.0	4.4	3.7
Others	3.8	5.0	▲	2.8	▼	3.6	3.7	4.1	3.0
Heroin	1.9	2.4	1.8	1.8	2.4	2.2	2.2	1.8	1.5
Steroids	1.9	2.7	2.1	1.4	2.7	▲	1.6	2.8	1.3
Injected Drugs	1.7	1.9	1.6	1.5	2.3	1.7	1.9	1.1	▼
MDMA	5.1	7.1	▲	4.2	4.6	4.6	5.6	5.5	4.3
Rohypnol	1.5	2.0	1.3	1.1	2.2	▲	1.5	1.8	1.2
GHB	1.1	1.5	1.2	1.2	1.3	1.3	1.1	1.1	0.7
Over the counter drugs	13.5	14.9	12.2	12.0	13.9	13.4	14.3	13.4	14.1

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 12th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	35.2	35.3	30.8 ▼	32.9	37.8 ▲	33.3 ▼	38.4	35.2	39.1 ▲
Cigars	30.5	29.8	29.5	27.4 ▼	31.2	31.0	26.3 ▼	29.8	33.4 ▲
Pipe	16.6	26.1 ▲	13.7 ▼	14.1 ▼	19.0 ▲	17.8	12.4 ▼	12.3 ▼	17.3
Smokeless Tobacco	14.6	11.6 ▼	11.0 ▼	13.7	18.3 ▲	12.5 ▼	11.7 ▼	15.3	19.4 ▲
Alcohol	60.2	63.3 ▲	55.8 ▼	56.5 ▼	59.9	58.3 ▼	56.9 ▼	61.9	65.3 ▲
Marijuana	26.6	29.4 ▲	24.3	25.0	25.0	28.7 ▲	29.5	23.8 ▼	27.2
Cocaine	5.3	6.3	5.0	5.3	5.9	5.4	5.2	4.3 ▼	5.3
Crack	2.0	2.4	2.0	1.4	2.7	2.2	2.3	1.4 ▼	1.9
Inhalants	4.0	4.0	3.8	4.0	4.5	3.4	5.4 ▲	4.2	4.0
Amphetamines	5.4	5.2	4.7	4.8	6.2	5.3	7.3 ▲	4.9	5.8
Methamphetamines	1.9	1.9	1.7	1.8	3.4 ▲	1.7	1.8	1.7	1.8
Ritalin	7.0	6.7	7.1	7.4	6.5	6.9	8.0	7.0	7.0
Tranquilizers	8.3	8.0	6.7 ▼	6.4 ▼	10.4 ▲	8.7	9.6	7.2	8.9
Narcotics	7.6	6.8	5.6 ▼	7.1	8.1	8.0	9.3	6.2 ▼	9.6 ▲
Psychedelics <sup>#</sup>	4.0	5.2 ▲	3.6	4.3	4.1	4.3	3.6	3.0 ▼	4.0
LSD	2.6	3.3	2.2	3.2	2.8	2.9	2.4	1.8 ▼	2.7
Others	2.7	3.9 ▲	2.4	2.6	2.6	2.9	2.2	2.1	3.0
Heroin	1.3	1.8	1.3	1.1	1.4	1.4	1.2	0.9	1.0
Steroids	1.3	2.2 ▲	1.3	1.0	1.9 ▲	1.1	2.0	0.6	1.4
Injected Drugs	1.3	1.6	1.2	1.1	1.7	1.2	1.1	0.9 ▼	1.4
MDMA	3.2	4.9 ▲	2.6	3.2	2.7	3.7	3.5	2.1	3.2
Rohypnol	1.0	1.4	0.9	1.0	0.9	1.1	1.4	0.7	0.8
GHB	0.8	1.2	0.8	0.8	0.7	0.8	1.1	0.5 ▼	0.5
Over the counter drugs	8.5	9.5	7.4	7.4	8.6	8.5	8.9	8.4	8.8

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).  
▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 12th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
Cigarettes	24.3	22.4	21.1 ▼	22.0	26.9 ▲	23.1	25.4	24.5	27.6 ▲
Cigars	17.5	16.9	17.6	13.8 ▼	17.3	17.6	13.7 ▼	16.9	20.8 ▲
Pipe	9.3	14.1 ▲	7.1 ▼	7.3 ▼	10.5	10.7 ▲	6.5 ▼	6.2 ▼	10.3
Smokeless Tobacco	9.1	7.3 ▼	6.3 ▼	7.6	11.3 ▲	7.4 ▼	6.9 ▼	10.0	13.2 ▲
Alcohol	39.7	42.6 ▲	36.2 ▼	36.8 ▼	36.6 ▼	37.6 ▼	35.7 ▼	42.9 ▲	45.3 ▲
Marijuana	15.8	17.7	14.2	14.5	15.2	17.1 ▲	17.8	13.6 ▼	16.8
Cocaine	2.4	3.3 ▲	2.7	2.7	2.7	2.5	2.1	1.4 ▼	2.1
Crack	1.1	1.6	1.2	0.8	1.4	1.1	1.2	0.5 ▼	0.9
Inhalants	1.5	1.8	1.7	1.5	1.9	1.3	2.2	1.4	1.3
Amphetamines	2.5	2.2	2.4	2.1	3.2 ▲	2.1	3.4	2.1	2.7
Methamphetamines	1.0	1.2	0.8	1.0	1.6 ▲	1.0	1.0	0.9	1.0
Ritalin	2.9	2.8	3.1	3.2	2.5	2.4	4.2 ▲	3.2	3.1
Tranquilizers	4.0	4.3	3.1	3.3	4.7	4.1	5.5 ▲	3.3	4.2
Narcotics	3.8	3.5	2.5 ▼	3.3	4.1	3.7	6.1 ▲	3.3	4.9 ▲
Psychedelics <sup>#</sup>	1.6	1.9	1.7	1.7	2.0	1.7	1.3	1.2	1.7
LSD	1.1	1.3	1.0	1.2	1.1	1.2	0.8	0.7	1.1
Others	1.1	1.7 ▲	1.2	1.2	1.3	1.1	0.8	0.7 ▼	1.1
Heroin	0.8	1.4 ▲	0.9	0.6	0.8	0.9	0.8	0.5	0.7
Steroids	0.7	1.5 ▲	0.6	0.4	1.0	0.7	1.1	0.2	0.9
Injected Drugs	0.9	1.3	0.7	0.6	1.3	0.8	0.9	0.6	1.0
MDMA	1.4	2.6 ▲	1.5	0.9	1.3	1.4	1.4	1.1	1.2
Rohypnol	0.6	1.0 ▲	0.6	0.5	0.6	0.6	0.8	0.3	0.5
GHB	0.5	1.1 ▲	0.6	0.5	0.5	0.4	0.7	0.3	0.3
Over the counter drugs	4.3	5.0	3.7	3.6	4.3	4.5	4.7	3.8	4.9

NOTES: # Psychedelic rate computed by analyses of responses indicating either LSD or Other Psychedelics use, or endorsement of both.

▲ indicates that local prevalence rates are significantly higher than state rates ( $p < .05$ ).

▼ indicates that local prevalence rates are significantly lower than state rates ( $p < .05$ ).

**Percentage Reporting Daily and Special Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 12th Grade Students: 2007**

DRUG	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
<b>CIGARETTES</b>									
Daily Use	14.7	14.1	12.7 ▼	11.6 ▼	17.7 ▲	14.3	16.3	14.1	16.5 ▲
1/2+ Pack per Day	8.5	7.5	6.3 ▼	5.7 ▼	11.6 ▲	8.6	10.9 ▲	7.3	10.0 ▲
<b>CIGARS</b>									
	3.8	3.3	3.4	2.7 ▼	4.3	4.3	2.6	3.6	4.2
<b>SMOKELESS TOBACCO</b>									
	4.2	2.7 ▼	2.6 ▼	3.2	5.4 ▲	3.1 ▼	3.3	5.3 ▲	6.5 ▲
<b>ALCOHOL</b>									
Daily Use	4.6	4.8	3.8	3.3 ▼	4.7	4.1	4.2	4.9	6.2 ▲
Binge Drinking	28.6	29.8	25.5 ▼	27.7	27.1	26.7 ▼	24.9 ▼	31.5 ▲	32.5 ▲
<b>MARIJUANA</b>									
	5.3	5.9	4.3	4.3	5.6	5.7	6.5	4.5	5.7

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ( $p<.05$ ).  
 ▼ indicates that local prevalence rates are significantly lower than state rates ( $p<.05$ ).




## IV. TRENDS AND PREVALENCE GRAPHS

**Trends Graphs .....** 112


**Prevalence Graphs .....** 126

## Trends of Annual Alcohol Use Among Indiana Students: 1993- 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Monthly Alcohol Use Among Indiana Students: 1993- 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Annual Cigarette Use Among Indiana Students: 1993- 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Monthly Cigarette Use Among Indiana Students: 1993- 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Annual Smokeless Tobacco Use Among Indiana Students: 1993- 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Monthly Smokeless Tobacco Use Among Indiana Students: 1993- 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Annual Marijuana Use Among Indiana Students: 1993- 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Monthly Marijuana Use Among Indiana Students: 1993-2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Annual Inhalent Use Among Indiana Students: 1993-2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Monthly Inhalant Use Among Indiana Students: 1993-2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Binge Drinking Among Indiana Students: 1993-2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Trends of Binge Drinking by Indiana and U.S.A. 12th Graders: 1975-2007




Source: Indiana Prevention Resource Center at Indiana University, 2007  
Johnston, et al., National Institute on Drug Abuse, 2006

## Trends of Monthly Cigarette Use by Indiana and U.S.A. 12th Graders: 1975-2007




Source: Indiana Prevention Resource Center at Indiana University, 2007  
 Johnston, et al., National Institute on Drug Abuse, 2006

## Trends of Monthly Marijuana Use by Indiana and U.S.A. 12th Graders: 1975-2007




Source: Indiana Prevention Resource Center at Indiana University, 2007  
Johnston, et al., National Institute on Drug Abuse, 2006

## Monthly Use of Selected Drugs and Binge Drinking by Indiana and U.S.A. 12th Graders




Source: Indiana Prevention Resource Center at Indiana University, 2007  
Johnston, et al., National Institute on Drug Abuse, 2006

## Alcohol Use Among Indiana Students 6th - 12th Graders, 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Cigarette Use Among Indiana Students 6th - 12th Graders, 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Smokeless Tobacco Use Among Indiana Students 6th - 12th Graders, 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Marijuana Use Among Indiana Students 6th - 12th Graders, 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Inhalant Use Among Indiana Students 6th - 12th Graders, 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Binge Drinking Among Indiana Students by Grade and Gender, 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Monthly Alcohol Use Among Indiana Students by Grade and Gender, 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007

## Monthly Cigarette Use Among Indiana Students by Grade and Gender, 2007




Source: Indiana Prevention Resource Center at Indiana University, 2007



Source: Indiana Prevention Resource Center at Indiana University, 2007

## Monthly Marijuana Use Among Indiana Students by Grade and Gender, 2007



Source: Indiana Prevention Resource Center at Indiana University, 2007

## V. FREQUENCY TABLES

<b>All, Grades 6-12 .....</b>	138
<b>Male, Grades 6-12 .....</b>	175
<b>Female, Grades 6-12 .....</b>	212

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	2.9	3.0	2.6	2.7	2.6	2.3	2.4
	Never	86.8	80.1	70.5	65.0	58.0	54.1	50.2
	7 years or younger	1.8	2.0	2.5	2.1	2.0	1.7	1.3
	8 years	1.0	1.3	1.7	1.5	1.6	1.2	0.9
	9 years	1.2	1.3	1.5	1.5	1.5	1.2	1.0
	10 years	2.1	2.3	2.5	2.3	2.2	1.7	1.3
	11 years	2.3	3.0	3.5	2.9	2.6	2.1	1.7
	12 years	1.6	4.3	5.2	4.6	4.5	4.0	3.4
	13 years	0.3	2.4	6.7	6.3	5.9	5.0	4.3
	14 years		0.3	2.9	7.3	6.3	5.5	4.6
	15 years			0.4	3.5	8.8	8.3	6.7
	16 years				0.3	3.7	9.5	9.9
	17 years					0.2	3.3	8.2
	18 years or older						0.4	4.1
Cigars	No Answer	1.8	1.9	2.0	2.5	2.6	2.7	2.6
	Never	93.9	90.1	84.5	79.5	71.9	67.7	61.1
	7 years or younger	0.6	0.9	0.9	0.8	0.7	0.6	0.5
	8 years	0.4	0.4	0.7	0.5	0.6	0.5	0.3
	9 years	0.5	0.5	0.7	0.6	0.6	0.4	0.3
	10 years	0.9	1.1	1.2	1.1	1.0	0.8	0.6
	11 years	1.0	1.4	1.6	1.4	1.0	0.8	0.6
	12 years	0.7	2.1	2.5	2.2	2.3	1.8	1.5
	13 years	0.1	1.5	3.8	3.4	3.4	2.6	2.0
	14 years		0.2	1.9	4.9	4.6	3.4	3.0
	15 years			0.2	2.9	7.5	6.8	5.0
	16 years				0.3	3.5	8.6	9.5
	17 years					0.2	3.0	7.9
	18 years or older						0.4	5.1
Pipe	No Answer	2.1	2.5	3.1	3.7	4.1	4.7	4.6
	Never	97.0	95.0	91.9	88.0	83.8	80.3	76.4
	7 years or younger	0.1	0.2	0.2	0.3	0.3	0.3	0.3
	8 years	0.1	0.2	0.2	0.1	0.2	0.2	0.1
	9 years	0.1	0.1	0.2	0.2	0.2	0.2	0.1
	10 years	0.2	0.3	0.4	0.4	0.4	0.3	0.2
	11 years	0.2	0.3	0.5	0.6	0.5	0.4	0.3
	12 years	0.2	0.7	1.0	1.0	1.1	0.9	0.7
	13 years	0.1	0.6	1.5	1.8	1.6	1.3	1.0
	14 years		0.1	0.8	2.3	2.3	1.7	1.5
	15 years			0.1	1.4	3.6	3.2	2.6
	16 years				0.1	1.7	4.8	4.5
	17 years					0.1	1.6	4.7
	18 years or older						0.1	3.0

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smokeless Tobacco	No Answer	1.6	1.8	1.8	2.1	2.5	2.6	2.5
	Never	96.2	94.2	91.3	87.6	83.0	80.9	78.5
	7 years or younger	0.3	0.5	0.5	0.6	0.7	0.5	0.5
	8 years	0.2	0.3	0.3	0.4	0.4	0.3	0.2
	9 years	0.2	0.3	0.3	0.4	0.3	0.3	0.2
	10 years	0.5	0.5	0.7	0.5	0.6	0.5	0.4
	11 years	0.6	0.6	0.7	0.7	0.5	0.4	0.4
	12 years	0.4	1.0	1.2	1.1	1.2	1.0	0.9
	13 years	0.1	0.9	1.9	2.0	1.7	1.4	1.1
	14 years		0.1	1.2	2.6	2.5	1.9	1.6
	15 years			0.1	1.9	4.3	3.7	3.1
	16 years				0.2	2.1	4.6	4.8
	17 years					0.1	1.5	3.9
	18 years or older						0.2	1.9
Alcohol	No Answer	3.1	3.4	3.8	3.8	3.8	4.1	3.8
	Never	76.7	66.8	54.5	47.9	38.4	34.9	29.8
	7 years or younger	3.7	3.6	3.3	2.3	2.4	1.9	1.8
	8 years	1.5	1.7	1.6	1.1	1.0	0.8	0.6
	9 years	2.0	1.6	1.6	1.2	1.0	0.6	0.4
	10 years	3.4	3.3	3.3	2.4	2.0	1.6	1.1
	11 years	5.0	4.5	3.8	2.8	2.1	1.5	1.1
	12 years	3.9	8.3	7.6	5.5	4.6	3.7	3.1
	13 years	0.6	6.1	12.6	10.6	8.1	6.5	5.4
	14 years		0.7	7.1	13.6	11.9	9.3	8.0
	15 years			0.8	8.0	17.1	15.1	13.1
	16 years				0.8	7.1	14.8	16.8
	17 years					0.4	4.8	10.3
	18 years or older						0.4	4.6
Marijuana	No Answer	2.0	2.2	2.3	2.8	3.2	3.4	3.1
	Never	95.4	90.7	82.9	77.5	68.7	66.0	62.1
	7 years or younger	0.2	0.3	0.5	0.4	0.5	0.3	0.3
	8 years	0.1	0.2	0.4	0.3	0.4	0.3	0.2
	9 years	0.2	0.3	0.4	0.4	0.4	0.3	0.2
	10 years	0.4	0.7	0.9	0.7	0.7	0.6	0.5
	11 years	0.8	1.1	1.4	1.2	1.2	0.8	0.7
	12 years	0.7	2.1	2.6	2.5	2.4	1.9	1.7
	13 years	0.2	2.0	5.0	4.1	4.3	3.5	3.0
	14 years		0.3	3.0	6.0	5.7	4.4	4.3
	15 years			0.5	3.7	8.4	7.5	6.7
	16 years				0.4	3.9	8.3	8.7
	17 years					0.2	2.5	6.2
	18 years or older						0.2	2.4

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	1.9	2.0	2.1	2.6	3.0	3.2	3.1
	Never	97.7	96.9	95.9	94.2	91.9	90.8	89.8
	7 years or younger	0.1	0.1	0.1	0.1	0.2	0.1	0.2
	8 years	0.0	0.0	0.1	0.0	0.0	0.1	0.0
	9 years	0.0	0.0	0.1	0.0	0.1	0.1	0.1
	10 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	11 years	0.1	0.1	0.2	0.1	0.1	0.1	0.1
	12 years	0.1	0.3	0.2	0.3	0.2	0.2	0.1
	13 years	0.0	0.3	0.7	0.5	0.5	0.4	0.2
	14 years		0.1	0.5	1.0	0.8	0.6	0.5
	15 years			0.1	1.0	1.7	1.3	0.7
	16 years				0.1	1.2	2.1	1.8
	17 years					0.1	0.9	2.2
	18 years or older						0.1	1.0
Crack	No Answer	2.1	2.2	2.3	2.8	3.2	3.5	3.1
	Never	97.4	96.9	96.1	95.3	94.3	93.8	94.1
	7 years or younger	0.1	0.1	0.1	0.2	0.2	0.1	0.2
	8 years	0.0	0.0	0.0	0.0	0.0	0.1	0.0
	9 years	0.0	0.0	0.1	0.0	0.1	0.1	0.0
	10 years	0.1	0.1	0.1	0.0	0.1	0.1	0.1
	11 years	0.1	0.2	0.1	0.1	0.1	0.1	0.0
	12 years	0.1	0.2	0.3	0.2	0.2	0.1	0.1
	13 years	0.0	0.2	0.4	0.3	0.2	0.3	0.2
	14 years		0.0	0.4	0.5	0.4	0.3	0.2
	15 years			0.1	0.5	0.7	0.6	0.3
	16 years				0.1	0.5	0.7	0.6
	17 years					0.0	0.3	0.6
	18 years or older						0.1	0.4
Inhalants	No Answer	2.7	2.4	2.5	2.9	3.1	3.4	3.4
	Never	92.1	90.8	88.4	88.6	87.7	88.1	89.2
	7 years or younger	1.2	0.9	0.8	0.5	0.5	0.3	0.3
	8 years	0.5	0.4	0.5	0.2	0.2	0.2	0.1
	9 years	0.4	0.5	0.4	0.2	0.2	0.2	0.1
	10 years	0.9	0.8	0.8	0.4	0.5	0.3	0.3
	11 years	1.2	1.1	1.1	0.9	0.5	0.3	0.2
	12 years	0.9	1.7	1.8	1.2	1.0	0.8	0.5
	13 years	0.1	1.2	2.4	1.9	1.5	1.2	0.9
	14 years		0.2	1.2	2.1	1.8	1.3	0.9
	15 years			0.1	1.0	2.2	1.9	1.2
	16 years				0.1	0.8	1.5	1.4
	17 years					0.1	0.5	1.1
	18 years or older						0.1	0.4

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Amphetamines	No Answer	3.1	2.4	2.4	2.8	3.2	3.5	3.3
	Never	96.6	96.7	95.1	92.8	89.7	88.6	88.7
	7 years or younger	0.1	0.0	0.1	0.1	0.1	0.1	0.1
	8 years	0.0	0.0	0.1	0.0	0.1	0.1	0.0
	9 years	0.0	0.0	0.1	0.1	0.0	0.1	0.0
	10 years	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	11 years	0.1	0.1	0.2	0.2	0.2	0.1	0.1
	12 years	0.1	0.3	0.5	0.6	0.4	0.3	0.2
	13 years	0.0	0.3	0.8	0.8	1.1	0.7	0.5
	14 years			0.0	0.6	1.4	1.6	1.2
	15 years				0.1	0.9	2.2	2.1
	16 years					0.1	1.1	2.2
	17 years						0.1	0.7
	18 years or older							0.1
Methamphetamines	No Answer	2.8	2.4	2.4	2.8	3.3	3.5	3.2
	Never	96.9	96.8	96.4	95.5	94.3	93.7	93.9
	7 years or younger	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	8 years	0.0	0.0	0.0	0.0	0.0	0.1	0.0
	9 years	0.0	0.0	0.1	0.0	0.0	0.0	0.0
	10 years	0.0	0.1	0.1	0.0	0.1	0.1	0.0
	11 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	12 years	0.1	0.2	0.2	0.2	0.2	0.1	0.1
	13 years	0.0	0.3	0.4	0.3	0.3	0.3	0.2
	14 years			0.0	0.3	0.5	0.5	0.3
	15 years				0.0	0.4	0.7	0.8
	16 years					0.1	0.4	0.6
	17 years						0.0	0.3
	18 years or older							0.1
Ritalin/Adderall	No Answer	3.6	2.6	2.6	3.0	3.3	3.5	3.4
	Never	95.7	95.6	93.6	90.7	87.4	87.2	86.4
	7 years or younger	0.1	0.2	0.2	0.2	0.2	0.1	0.2
	8 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	9 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	10 years	0.1	0.2	0.2	0.1	0.2	0.2	0.1
	11 years	0.2	0.2	0.4	0.2	0.2	0.1	0.1
	12 years	0.2	0.5	0.6	0.7	0.5	0.3	0.3
	13 years	0.0	0.5	1.2	1.2	1.1	0.7	0.5
	14 years			0.1	0.9	2.1	1.8	1.3
	15 years				0.1	1.6	3.3	2.4
	16 years					0.2	1.7	3.0
	17 years						0.1	1.0
	18 years or older							0.1

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Tranquilizers	No Answer	2.8	2.5	2.3	2.9	3.1	3.4	3.2
	Never	93.8	92.7	89.7	87.6	84.8	84.8	85.3
	7 years or younger	0.3	0.3	0.2	0.2	0.2	0.1	0.2
	8 years	0.2	0.2	0.1	0.1	0.1	0.1	0.0
	9 years	0.3	0.1	0.3	0.2	0.1	0.1	0.1
	10 years	0.6	0.4	0.4	0.3	0.3	0.1	0.1
	11 years	1.0	0.7	0.7	0.4	0.3	0.2	0.2
	12 years	0.8	1.5	1.5	1.1	0.7	0.5	0.4
	13 years	0.2	1.5	2.6	2.0	1.7	1.0	0.7
	14 years		0.2	1.9	3.0	2.6	1.7	1.1
	15 years			0.3	2.1	4.0	3.2	2.2
	16 years				0.2	2.0	3.4	3.3
	17 years					0.1	1.1	2.5
	18 years or older						0.1	0.9
Narcotics	No Answer	3.0	2.4	2.3	2.8	3.0	3.3	3.3
	Never	96.4	95.8	93.6	90.9	87.5	86.5	86.0
	7 years or younger	0.1	0.1	0.1	0.1	0.1	0.1	0.2
	8 years	0.1	0.0	0.1	0.1	0.1	0.1	0.0
	9 years	0.0	0.0	0.1	0.1	0.1	0.1	0.0
	10 years	0.1	0.2	0.2	0.1	0.2	0.1	0.1
	11 years	0.2	0.2	0.3	0.2	0.2	0.1	0.1
	12 years	0.1	0.5	0.7	0.6	0.5	0.3	0.3
	13 years	0.0	0.6	1.4	1.2	1.2	0.7	0.4
	14 years		0.1	1.2	2.1	1.9	1.5	0.9
	15 years			0.2	1.7	3.4	2.6	2.0
	16 years				0.2	1.8	3.3	3.2
	17 years					0.1	1.1	2.6
	18 years or older						0.1	1.1
LSD Only	No Answer	3.3	2.5	2.4	2.9	3.2	3.5	3.3
	Never	96.5	96.6	95.8	94.6	93.5	92.7	92.8
	7 years or younger	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	8 years	0.0	0.0	0.1	0.0	0.1	0.1	0.0
	9 years	0.0	0.0	0.1	0.0	0.0	0.0	0.0
	10 years	0.0	0.1	0.1	0.1	0.1	0.1	0.0
	11 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	12 years	0.1	0.2	0.3	0.2	0.2	0.2	0.1
	13 years	0.0	0.3	0.5	0.4	0.4	0.3	0.2
	14 years		0.1	0.5	0.7	0.5	0.4	0.3
	15 years			0.1	0.7	0.9	0.9	0.5
	16 years				0.1	0.7	1.1	1.1
	17 years					0.1	0.5	0.9
	18 years or older						0.1	0.5

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Psychedelics	No Answer	3.8	2.7	2.5	3.0	3.3	3.5	3.5
	Never	96.0	96.8	96.5	95.6	94.5	94.0	93.2
	7 years or younger	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	8 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	9 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	10 years	0.0	0.1	0.1	0.0	0.1	0.1	0.0
	11 years	0.0	0.1	0.1	0.1	0.1	0.1	0.0
	12 years	0.1	0.1	0.2	0.2	0.1	0.1	0.1
	13 years	0.0	0.1	0.3	0.3	0.3	0.2	0.1
	14 years			0.0	0.2	0.4	0.4	0.3
	15 years				0.0	0.3	0.7	0.6
	16 years					0.0	0.4	0.8
	17 years						0.0	0.3
	18 years or older							0.0
Heroin	No Answer	2.9	2.4	2.3	2.9	3.2	3.5	3.3
	Never	96.8	96.9	96.7	95.9	95.2	94.8	95.1
	7 years or younger	0.1	0.1	0.1	0.1	0.1	0.1	0.2
	8 years	0.0	0.0	0.1	0.0	0.1	0.0	0.0
	9 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	10 years	0.0	0.1	0.1	0.1	0.1	0.1	0.0
	11 years	0.0	0.1	0.1	0.0	0.1	0.1	0.1
	12 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	13 years	0.0	0.2	0.3	0.2	0.2	0.2	0.1
	14 years			0.0	0.2	0.3	0.2	0.1
	15 years				0.0	0.2	0.4	0.3
	16 years					0.0	0.3	0.4
	17 years						0.0	0.2
	18 years or older							0.0
Steroids	No Answer	2.6	2.5	2.4	3.0	3.3	3.5	3.5
	Never	96.9	96.6	96.4	95.7	95.0	95.0	94.9
	7 years or younger	0.1	0.1	0.1	0.1	0.2	0.1	0.1
	8 years	0.0	0.0	0.0	0.0	0.1	0.0	0.0
	9 years	0.1	0.1	0.1	0.0	0.0	0.0	0.0
	10 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	11 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	12 years	0.1	0.2	0.2	0.1	0.1	0.1	0.0
	13 years	0.0	0.3	0.3	0.2	0.2	0.1	0.1
	14 years			0.0	0.3	0.3	0.2	0.1
	15 years				0.0	0.4	0.3	0.2
	16 years					0.1	0.4	0.4
	17 years						0.0	0.2
	18 years or older							0.1

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	3.7	2.6	2.5	2.9	3.3	3.4	3.5
	Never	96.1	96.5	95.6	94.4	93.1	92.4	92.0
	7 years or younger	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	8 years	0.0	0.0	0.0	0.0	0.0	0.1	0.0
	9 years	0.0	0.0	0.1	0.0	0.1	0.0	0.0
	10 years	0.0	0.0	0.1	0.0	0.1	0.1	0.1
	11 years	0.0	0.1	0.1	0.1	0.1	0.1	0.0
	12 years	0.0	0.2	0.3	0.2	0.2	0.1	0.1
	13 years	0.0	0.3	0.6	0.5	0.4	0.3	0.2
	14 years		0.1	0.5	0.8	0.6	0.4	0.3
	15 years			0.1	0.7	1.2	0.9	0.5
	16 years				0.1	0.9	1.4	1.0
	17 years					0.1	0.7	1.2
	18 years or older						0.1	0.9
Rohypnol	No Answer	3.9	2.7	2.5	2.9	3.3	3.5	3.4
	Never	96.0	96.8	96.7	95.9	95.3	95.0	95.3
	7 years or younger	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	8 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	9 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	10 years	0.0	0.0	0.1	0.0	0.0	0.1	0.0
	11 years	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	12 years	0.0	0.1	0.1	0.2	0.1	0.1	0.1
	13 years	0.0	0.1	0.2	0.2	0.1	0.2	0.1
	14 years		0.0	0.1	0.3	0.3	0.2	0.1
	15 years			0.0	0.2	0.4	0.2	0.2
	16 years				0.0	0.2	0.4	0.2
	17 years					0.0	0.1	0.2
	18 years or older						0.0	0.2
GHB	No Answer	4.1	2.8	2.6	3.0	3.3	3.5	3.5
	Never	95.8	96.9	96.9	96.3	95.7	95.5	95.5
	7 years or younger	0.0	0.0	0.1	0.1	0.1	0.0	0.1
	8 years	0.0	0.0	0.0	0.0	0.0	0.1	0.0
	9 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	10 years	0.0	0.0	0.0	0.1	0.1	0.1	0.0
	11 years	0.0	0.1	0.1	0.0	0.0	0.1	0.0
	12 years	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	13 years	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	14 years		0.0	0.1	0.2	0.2	0.1	0.1
	15 years			0.0	0.1	0.2	0.2	0.1
	16 years				0.0	0.1	0.1	0.1
	17 years					0.0	0.1	0.1
	18 years or older						0.0	0.1

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Over the counter drugs	No Answer	3.7	3.0	2.9	3.3	3.6	3.9	3.7
	Never	92.4	91.5	88.5	86.2	84.2	84.5	84.6
	7 years or younger	1.0	0.9	1.1	0.8	0.7	0.4	0.6
	8 years	0.4	0.3	0.3	0.3	0.2	0.2	0.1
	9 years	0.3	0.3	0.3	0.2	0.2	0.1	0.1
	10 years	0.6	0.5	0.6	0.4	0.4	0.3	0.2
	11 years	0.7	0.7	0.8	0.6	0.4	0.2	0.2
	12 years	0.8	1.3	1.4	1.2	1.1	0.7	0.5
	13 years	0.1	1.2	2.3	2.0	1.6	1.1	0.7
	14 years		0.2	1.7	3.1	2.5	1.7	1.2
	15 years			0.2	1.8	3.3	2.6	2.1
	16 years				0.2	1.6	3.1	2.8
	17 years					0.1	0.9	2.2
	18 years or older						0.1	0.9

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	0.5	0.5	0.5	0.4	0.3	0.4	0.3
	Never	88.4	81.7	71.4	66.1	59.1	55.0	51.3
	Once or Twice	8.3	10.6	14.6	14.5	15.8	16.0	17.3
	Occasionally	1.3	3.4	6.2	8.4	10.2	12.0	13.6
	Past Regularly	1.0	2.0	3.1	4.2	5.0	5.2	5.1
	Current Regularly	0.5	1.9	4.2	6.4	9.6	11.4	12.3
Cigars	No Answer	0.3	0.3	0.3	0.4	0.5	0.5	0.3
	Never	95.0	91.0	85.0	80.2	72.8	68.5	61.8
	1-5 times	3.8	6.1	9.9	11.9	14.8	16.3	19.0
	6-19 times	0.4	0.9	1.9	2.9	4.7	6.1	7.5
	20-40 times	0.2	0.6	1.0	1.6	2.7	3.4	4.6
	40+ times	0.3	1.0	1.9	3.0	4.6	5.3	6.8
Pipe	No Answer	0.5	0.6	0.8	1.0	1.1	1.3	1.2
	Never	98.3	96.4	93.4	89.8	85.6	82.5	78.8
	1-5 times	1.0	1.9	3.4	4.9	6.3	7.7	9.5
	6-19 times	0.2	0.5	1.1	1.7	2.4	3.2	4.1
	20-40 times	0.0	0.2	0.5	1.0	1.5	1.9	2.3
	40+ times	0.1	0.3	0.8	1.6	3.1	3.4	4.0
Smokeless Tobacco	No Answer	0.3	0.3	0.3	0.3	0.4	0.4	0.4
	Never	97.3	95.4	92.5	88.8	84.6	82.7	80.2
	1-5 times	1.9	3.1	4.7	6.3	8.0	8.6	9.5
	6-19 times	0.2	0.5	1.0	1.4	1.9	2.2	2.4
	20-40 times	0.1	0.2	0.5	0.9	1.3	1.3	1.4
	40+ times	0.2	0.5	1.1	2.2	3.8	4.8	6.1
Alcohol	No Answer	0.7	0.7	0.8	0.9	0.9	1.0	1.0
	Never	76.8	66.6	53.9	47.5	38.2	34.7	29.8
	1-5 times	18.5	23.4	27.6	26.6	26.2	24.3	21.6
	6-19 times	2.7	5.5	9.5	11.9	14.2	15.0	15.5
	20-40 times	0.6	1.9	3.9	5.9	8.6	9.7	11.3
	40+ times	0.6	1.8	4.4	7.3	12.0	15.3	20.8
Marijuana	No Answer	0.5	0.6	0.7	0.7	0.8	0.9	0.9
	Never	96.5	91.7	83.2	78.2	69.3	66.7	62.6
	1-5 times	2.1	4.5	7.9	9.3	11.6	12.6	13.5
	6-19 times	0.5	1.3	2.9	3.6	5.0	5.4	6.3
	20-40 times	0.2	0.6	1.7	2.5	3.3	3.6	4.5
	40+ times	0.3	1.3	3.7	5.8	10.0	10.8	12.3
Cocaine	No Answer	0.4	0.4	0.5	0.5	0.6	0.7	0.7
	Never	99.1	98.3	97.1	95.7	93.6	92.4	91.4
	1-5 times	0.3	0.8	1.5	2.5	3.5	3.9	4.2
	6-19 times	0.1	0.2	0.3	0.4	0.9	1.3	1.4
	20-40 times	0.0	0.1	0.2	0.3	0.6	0.8	1.0
	40+ times	0.1	0.2	0.3	0.5	0.8	0.9	1.2

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Crack	No Answer	0.5	0.4	0.5	0.5	0.6	0.6	0.6
	Never	98.8	98.3	97.4	97.1	96.3	96.1	96.1
	1-5 times	0.5	0.9	1.3	1.5	1.8	2.0	1.9
	6-19 times	0.1	0.2	0.2	0.3	0.5	0.4	0.4
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.3	0.4
	40+ times	0.1	0.1	0.3	0.4	0.6	0.6	0.6
Inhalants	No Answer	0.9	0.6	0.5	0.7	0.6	0.6	0.7
	Never	92.9	91.5	88.9	89.5	88.8	89.6	90.8
	1-5 times	4.8	5.7	7.2	6.5	6.8	6.4	5.4
	6-19 times	0.7	1.2	1.8	1.8	1.9	1.7	1.6
	20-40 times	0.3	0.4	0.7	0.7	0.8	0.8	0.8
	40+ times	0.3	0.6	0.8	0.8	1.0	0.9	0.7
Amphetamines	No Answer	1.8	0.9	0.7	0.7	0.8	0.7	0.8
	Never	97.8	97.9	96.3	94.0	91.0	90.2	90.2
	1-5 times	0.2	0.7	1.7	2.7	4.0	4.2	3.9
	6-19 times	0.1	0.2	0.7	1.2	1.9	2.3	2.2
	20-40 times	0.0	0.1	0.3	0.7	1.1	1.2	1.3
	40+ times	0.0	0.1	0.3	0.8	1.2	1.5	1.6
Methamphetamine	No Answer	1.2	0.6	0.6	0.6	0.7	0.7	0.8
	Never	98.4	98.3	97.8	97.2	96.3	96.0	95.9
	1-5 times	0.2	0.6	0.9	1.3	1.6	1.7	1.7
	6-19 times	0.0	0.2	0.3	0.3	0.5	0.5	0.6
	20-40 times	0.0	0.1	0.2	0.2	0.4	0.4	0.3
	40+ times	0.1	0.2	0.3	0.3	0.5	0.7	0.8
Ritalin/Adderall	No Answer	1.9	1.0	0.8	0.8	0.8	0.9	0.9
	Never	97.0	96.8	94.7	91.9	88.6	88.5	87.8
	1-5 times	0.7	1.3	2.7	4.1	5.5	5.4	5.6
	6-19 times	0.1	0.4	0.8	1.5	2.3	2.6	2.8
	20-40 times	0.1	0.1	0.4	0.7	1.2	1.2	1.3
	40+ times	0.1	0.3	0.5	1.0	1.6	1.6	1.6
Tranquilizers	No Answer	1.1	0.7	0.6	0.7	0.8	0.8	0.8
	Never	95.0	93.6	90.3	88.4	85.6	85.8	86.3
	1-5 times	3.2	4.0	6.0	6.5	7.5	7.2	6.3
	6-19 times	0.5	1.0	1.7	2.2	2.9	3.0	3.2
	20-40 times	0.2	0.3	0.7	1.0	1.6	1.4	1.6
	40+ times	0.2	0.4	0.7	1.2	1.7	1.8	1.8
Narcotics	No Answer	1.6	0.8	0.7	0.7	0.7	0.7	0.9
	Never	97.6	97.0	94.3	91.8	88.4	87.6	87.0
	1-5 times	0.5	1.3	3.0	4.0	5.2	5.3	5.4
	6-19 times	0.1	0.4	1.0	1.6	2.4	2.9	3.0
	20-40 times	0.0	0.2	0.5	1.0	1.4	1.6	1.7
	40+ times	0.1	0.2	0.6	1.1	1.8	2.0	2.0

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Psychedelics (All)	No Answer	1.4	0.7	0.4	0.5	0.5	0.4	0.5
	Never	98.1	98.1	96.9	95.9	94.7	94.1	93.4
	1-5 times	0.3	0.8	1.5	2.2	2.9	3.3	3.8
	6-19 times	0.1	0.2	0.5	0.6	0.8	0.9	1.1
	20-40 times	0.0	0.1	0.2	0.3	0.5	0.4	0.4
	40+ times	0.1	0.1	0.4	0.5	0.6	0.8	0.7
LSD Only	No Answer	1.7	0.9	0.7	0.7	0.8	0.8	0.7
	Never	98.0	98.1	97.1	96.4	95.3	94.8	94.8
	1-5 times	0.2	0.6	1.3	1.9	2.5	2.8	2.9
	6-19 times	0.0	0.2	0.4	0.5	0.6	0.6	0.7
	20-40 times	0.0	0.1	0.2	0.2	0.4	0.3	0.3
	40+ times	0.1	0.1	0.3	0.4	0.5	0.7	0.5
Other Psychedelics	No Answer	2.3	1.1	0.8	0.8	0.9	0.9	0.9
	Never	97.3	98.3	97.9	97.4	96.5	96.2	95.3
	1-5 times	0.2	0.3	0.8	1.0	1.5	1.6	2.4
	6-19 times	0.1	0.1	0.2	0.3	0.5	0.6	0.7
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.2	0.3
	40+ times	0.1	0.1	0.2	0.3	0.4	0.5	0.5
Heroin	No Answer	1.3	0.7	0.6	0.7	0.8	0.8	0.8
	Never	98.3	98.4	98.1	97.8	97.2	97.2	97.3
	1-5 times	0.2	0.5	0.7	0.8	1.0	1.0	0.8
	6-19 times	0.0	0.2	0.2	0.2	0.4	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.3	0.3
	40+ times	0.1	0.2	0.3	0.3	0.4	0.5	0.5
Steroids	No Answer	1.1	0.9	0.8	0.8	0.9	0.9	0.8
	Never	98.2	97.9	97.7	97.6	97.1	97.3	97.2
	1-5 times	0.5	0.8	0.9	0.9	0.9	0.8	0.7
	6-19 times	0.1	0.2	0.2	0.2	0.3	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.2	0.3
	40+ times	0.1	0.2	0.3	0.4	0.6	0.5	0.6
Injected Drugs	No Answer	2.7	3.1	3.4	4.3	4.3	4.4	4.4
	Never	96.5	95.4	94.9	94.0	93.7	93.7	93.9
	1-5 times	0.5	0.8	0.9	0.9	1.0	0.8	0.7
	6-19 times	0.1	0.2	0.3	0.3	0.3	0.3	0.3
	20-40 times	0.0	0.2	0.1	0.2	0.2	0.2	0.2
	40+ times	0.1	0.2	0.4	0.4	0.5	0.6	0.5
MDMA	No Answer	2.3	1.0	0.7	0.7	0.8	0.7	0.8
	Never	97.5	97.9	97.0	96.1	94.9	94.4	94.1
	1-5 times	0.2	0.6	1.5	2.1	2.7	3.2	3.3
	6-19 times	0.0	0.2	0.4	0.5	0.7	0.7	0.8
	20-40 times	0.0	0.1	0.2	0.3	0.4	0.3	0.4
	40+ times	0.1	0.2	0.3	0.3	0.6	0.6	0.6

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Rohypnol	No Answer	2.4	1.1	0.8	0.8	0.9	0.8	0.9
	Never	97.4	98.3	98.1	97.8	97.4	97.5	97.6
	1-5 times	0.1	0.4	0.6	0.8	1.0	0.9	0.8
	6-19 times	0.0	0.1	0.2	0.3	0.3	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.2	0.2
	40+ times	0.0	0.1	0.2	0.2	0.3	0.3	0.3
GHB	No Answer	2.7	1.3	0.9	0.8	0.9	0.9	0.9
	Never	97.2	98.4	98.4	98.3	97.9	97.9	98.0
	1-5 times	0.1	0.2	0.4	0.5	0.5	0.5	0.5
	6-19 times	0.0	0.1	0.1	0.2	0.3	0.2	0.1
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.1	0.2	0.2	0.3	0.3	0.3
Over the counter drugs	No Answer	1.9	1.2	1.1	1.0	1.0	1.0	1.0
	Never	93.4	92.3	88.7	86.7	84.6	85.5	85.5
	1-5 times	3.0	3.9	5.9	6.8	7.5	7.2	7.0
	6-19 times	0.9	1.2	2.1	2.5	3.2	3.1	3.2
	20-40 times	0.3	0.6	0.9	1.2	1.5	1.1	1.2
	40+ times	0.5	0.8	1.4	1.8	2.2	2.2	2.1

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	1.2	1.1	0.9	1.0	0.9	1.0	0.9
	None	93.5	88.2	81.1	76.4	70.6	66.6	63.9
	Few times	3.8	6.9	10.6	12.3	14.8	16.2	18.3
	1-5 cigarettes/day	1.0	2.3	3.9	5.2	6.3	7.1	7.6
	1/2 pack/day	0.3	0.8	1.8	2.6	3.7	4.6	4.8
	1 pack/day	0.1	0.5	1.0	1.7	2.5	3.3	3.4
	1 & 1/2 pack/day	0.1	0.1	0.3	0.4	0.5	0.6	0.5
	2+ pack/day	0.1	0.2	0.3	0.5	0.6	0.6	0.6
Cigars	No Answer	2.2	2.6	2.8	3.5	3.9	4.0	3.8
	Never	94.7	91.2	86.2	81.8	75.4	72.0	65.7
	1-5 times	2.2	3.8	6.4	8.2	10.8	12.7	16.3
	6-19 times	0.5	1.0	1.8	2.4	4.1	4.8	6.1
	20-40 times	0.2	0.5	0.9	1.4	2.1	2.2	2.9
Pipe	40+ times	0.3	0.9	1.8	2.7	3.8	4.4	5.2
	No Answer	2.4	2.8	3.4	4.1	4.5	4.7	4.7
	Never	96.6	94.6	91.6	88.3	84.4	82.0	78.6
	1-5 times	0.7	1.4	2.7	3.8	5.0	6.3	8.4
	6-19 times	0.2	0.6	1.0	1.6	2.1	2.7	3.4
Smokeless Tobacco	20-40 times	0.1	0.2	0.5	0.9	1.5	1.6	2.0
	40+ times	0.1	0.3	0.8	1.4	2.5	2.8	3.0
	No Answer	2.1	2.5	2.8	3.5	3.8	3.8	3.9
	Never	96.2	94.2	91.4	87.8	84.2	83.1	81.5
	1-5 times	1.2	2.0	3.4	4.6	5.6	5.9	6.3
Alcohol	6-19 times	0.3	0.6	1.0	1.4	1.9	1.8	1.9
	20-40 times	0.1	0.2	0.5	0.8	1.2	1.1	1.3
	40+ times	0.1	0.5	1.0	1.9	3.3	4.4	5.1
	No Answer	2.4	2.8	3.1	3.6	4.0	4.0	3.9
	Never	82.3	72.4	60.3	53.0	44.3	40.7	35.9
Marijuana	1-5 times	12.4	17.6	23.4	24.4	25.8	26.1	24.6
	6-19 times	2.1	4.5	7.8	10.4	12.5	13.6	15.0
	20-40 times	0.5	1.5	2.8	4.3	6.5	7.4	8.9
	40+ times	0.4	1.2	2.6	4.2	6.9	8.3	11.7
	No Answer	2.4	2.7	3.1	3.9	4.2	4.3	4.4
	Never	95.4	91.0	83.9	79.2	72.3	70.9	69.1
	1-5 times	1.4	3.4	6.3	7.7	9.7	10.4	10.9
	6-19 times	0.4	1.2	2.6	3.3	4.3	4.4	4.7
	20-40 times	0.2	0.6	1.6	2.0	2.9	2.9	3.2
	40+ times	0.2	1.0	2.6	4.0	6.7	7.0	7.8

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	2.3	2.7	3.1	3.8	4.3	4.4	4.4
	Never	97.3	96.3	95.0	93.6	91.7	91.0	90.3
	1-5 times	0.2	0.6	1.1	1.6	2.4	2.7	3.0
	6-19 times	0.1	0.2	0.3	0.4	0.7	0.9	1.0
	20-40 times	0.0	0.1	0.2	0.2	0.5	0.5	0.7
	40+ times	0.0	0.1	0.2	0.3	0.4	0.5	0.6
Crack	No Answer	2.4	2.8	3.1	3.9	4.3	4.3	4.4
	Never	97.1	96.3	95.4	94.6	93.7	93.5	93.6
	1-5 times	0.3	0.5	0.8	0.9	1.1	1.2	1.1
	6-19 times	0.1	0.2	0.3	0.3	0.4	0.4	0.4
	20-40 times	0.1	0.1	0.1	0.1	0.3	0.2	0.3
	40+ times	0.0	0.1	0.2	0.2	0.2	0.3	0.3
Inhalants	No Answer	2.8	2.8	3.2	4.0	4.2	4.4	4.5
	Never	93.2	91.8	89.7	90.0	89.9	90.9	91.5
	1-5 times	2.8	3.4	4.7	3.9	3.8	2.9	2.6
	6-19 times	0.8	1.2	1.5	1.3	1.2	1.0	0.8
	20-40 times	0.3	0.4	0.5	0.5	0.5	0.4	0.4
	40+ times	0.2	0.3	0.4	0.4	0.4	0.4	0.3
Amphetamines	No Answer	3.5	3.0	3.3	4.0	4.4	4.4	4.5
	Never	96.2	96.1	94.5	92.3	89.9	89.8	90.1
	1-5 times	0.2	0.5	1.3	2.0	3.0	3.1	2.8
	6-19 times	0.0	0.2	0.6	0.9	1.5	1.3	1.4
	20-40 times	0.0	0.1	0.2	0.4	0.7	0.7	0.6
	40+ times	0.0	0.1	0.2	0.4	0.5	0.7	0.6
Methamphetamines	No Answer	3.1	2.9	3.2	3.9	4.4	4.5	4.6
	Never	96.7	96.4	95.6	94.6	93.8	93.5	93.5
	1-5 times	0.1	0.4	0.6	0.8	1.0	1.0	0.9
	6-19 times	0.0	0.1	0.3	0.4	0.4	0.4	0.4
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.3	0.3
	40+ times	0.0	0.1	0.2	0.2	0.2	0.3	0.3
Ritalin/Adderall	No Answer	3.6	3.1	3.3	4.0	4.4	4.4	4.5
	Never	95.7	95.4	93.4	90.7	88.2	88.7	88.5
	1-5 times	0.5	0.8	1.9	3.0	4.1	3.8	4.1
	6-19 times	0.1	0.3	0.7	1.2	1.8	1.6	1.6
	20-40 times	0.0	0.2	0.3	0.6	0.8	0.8	0.6
	40+ times	0.1	0.2	0.3	0.5	0.7	0.7	0.6
Tranquilizers	No Answer	3.0	2.9	3.3	4.0	4.4	4.5	4.5
	Never	94.2	92.9	90.0	88.1	86.0	86.6	87.2
	1-5 times	2.1	2.7	4.1	4.8	5.3	5.0	4.7
	6-19 times	0.5	0.9	1.5	1.6	2.3	2.0	2.0
	20-40 times	0.2	0.3	0.6	0.8	1.1	1.0	0.8
	40+ times	0.1	0.3	0.5	0.7	0.8	0.9	0.8

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	3.5	3.0	3.3	4.0	4.4	4.5	4.6
	Never	96.0	95.4	93.1	90.6	88.1	87.8	87.8
	1-5 times	0.3	0.9	2.0	3.0	4.0	4.0	3.9
	6-19 times	0.1	0.4	0.8	1.2	1.8	1.8	1.8
	20-40 times	0.0	0.2	0.4	0.6	0.9	1.0	1.1
	40+ times	0.0	0.1	0.3	0.5	0.9	0.9	0.8
Psychedelics (All)	No Answer	3.2	2.8	3.0	3.7	3.9	4.0	4.1
	Never	96.4	96.2	95.0	93.7	92.9	92.4	91.9
	1-5 times	0.2	0.5	1.1	1.5	1.9	2.0	2.6
	6-19 times	0.1	0.2	0.4	0.5	0.6	0.6	0.6
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.4	0.3
	40+ times	0.2	0.2	0.3	0.4	0.4	0.6	0.5
LSD Only	No Answer	3.6	3.1	3.3	4.0	4.3	4.5	4.5
	Never	96.2	96.2	95.2	94.0	93.2	92.9	92.9
	1-5 times	0.1	0.4	0.9	1.2	1.6	1.6	1.7
	6-19 times	0.0	0.1	0.3	0.4	0.4	0.5	0.4
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.2	0.2
	40+ times	0.0	0.1	0.2	0.2	0.2	0.4	0.3
Other Psychedelics	No Answer	4.0	3.3	3.5	4.2	4.5	4.4	4.5
	Never	95.7	96.1	95.5	94.5	93.7	93.4	92.7
	1-5 times	0.1	0.2	0.5	0.6	1.0	1.2	1.8
	6-19 times	0.0	0.1	0.2	0.3	0.4	0.4	0.4
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.3	0.2
	40+ times	0.1	0.2	0.2	0.2	0.3	0.4	0.4
Heroin	No Answer	3.2	2.9	3.3	4.0	4.4	4.5	4.6
	Never	96.6	96.4	95.8	94.9	94.4	94.1	94.2
	1-5 times	0.2	0.3	0.5	0.6	0.6	0.6	0.5
	6-19 times	0.0	0.1	0.2	0.2	0.2	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.2	0.2
	40+ times	0.0	0.1	0.1	0.2	0.2	0.3	0.3
Steroids	No Answer	2.9	2.9	3.4	4.2	4.5	4.5	4.6
	Never	96.6	96.1	95.5	94.7	94.2	94.2	94.1
	1-5 times	0.3	0.6	0.6	0.6	0.6	0.5	0.6
	6-19 times	0.1	0.2	0.2	0.2	0.2	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.2	0.1
	40+ times	0.1	0.1	0.2	0.2	0.3	0.3	0.4
Injected Drugs	No Answer	4.0	4.3	4.5	5.2	5.5	5.5	5.6
	Never	95.5	94.6	94.3	93.4	93.0	93.1	93.2
	1-5 times	0.4	0.6	0.7	0.8	0.7	0.5	0.5
	6-19 times	0.1	0.2	0.2	0.3	0.3	0.4	0.3
	20-40 times	0.0	0.1	0.2	0.1	0.2	0.2	0.2
	40+ times	0.1	0.1	0.2	0.2	0.3	0.3	0.3

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	3.9	3.2	3.3	4.1	4.3	4.4	4.5
	Never	95.9	95.9	95.0	93.8	92.8	92.6	92.3
	1-5 times	0.1	0.5	1.0	1.4	1.9	1.9	2.2
	6-19 times	0.0	0.2	0.4	0.4	0.5	0.5	0.5
	20-40 times	0.0	0.1	0.1	0.2	0.3	0.2	0.2
	40+ times	0.0	0.1	0.2	0.2	0.2	0.3	0.3
Rohypnol	No Answer	4.0	3.3	3.4	4.1	4.5	4.4	4.5
	Never	95.9	96.3	95.8	94.9	94.4	94.5	94.5
	1-5 times	0.1	0.3	0.4	0.6	0.6	0.6	0.5
	6-19 times	0.0	0.1	0.2	0.2	0.2	0.2	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.2	0.1
	40+ times	0.0	0.1	0.1	0.1	0.1	0.2	0.2
GHB	No Answer	4.2	3.3	3.5	4.2	4.5	4.5	4.6
	Never	95.7	96.3	96.0	95.2	94.8	94.7	94.7
	1-5 times	0.0	0.1	0.2	0.3	0.3	0.3	0.3
	6-19 times	0.0	0.1	0.1	0.2	0.2	0.1	0.1
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.2	0.1
	40+ times	0.0	0.1	0.1	0.1	0.1	0.2	0.2
Over the counter drugs	No Answer	3.6	3.4	3.7	4.5	4.8	5.0	5.0
	Never	92.6	91.4	88.2	86.3	84.9	86.0	86.6
	1-5 times	2.4	2.9	4.4	4.9	5.4	5.1	4.7
	6-19 times	0.9	1.2	2.0	2.3	2.6	2.1	2.0
	20-40 times	0.3	0.5	0.7	1.0	1.1	0.8	0.8
	40+ times	0.2	0.5	0.8	0.9	1.1	1.0	0.9

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	1.0	1.0	0.8	0.8	0.9	0.9	0.8
	None	96.2	92.8	88.4	84.4	79.8	76.6	74.9
	Few times	1.7	3.5	5.3	6.7	7.9	8.7	9.6
	1-5 cigarettes/day	0.6	1.4	2.8	3.7	4.8	5.7	6.2
	1/2 pack/day	0.2	0.5	1.2	1.9	3.0	3.8	4.1
	1 pack/day	0.1	0.4	0.9	1.5	2.4	3.2	3.3
	1 & 1/2 pack/day	0.1	0.1	0.2	0.4	0.5	0.5	0.5
	2+ pack/day	0.1	0.3	0.3	0.5	0.6	0.7	0.7
Cigars	No Answer	2.8	3.3	3.7	4.5	4.9	4.9	4.8
	Never	95.6	93.1	90.0	86.8	82.8	81.3	77.7
	1-5 times	1.2	2.3	3.9	5.0	7.2	8.4	11.0
	6-19 times	0.2	0.6	0.9	1.4	2.0	2.0	2.7
	20-40 times	0.1	0.3	0.6	0.8	1.0	1.1	1.1
Pipe	40+ times	0.1	0.4	0.9	1.4	2.1	2.4	2.7
	No Answer	3.0	3.6	3.9	4.8	5.3	5.3	5.3
	Never	96.5	94.8	93.0	90.8	88.0	87.3	85.4
	1-5 times	0.4	0.9	1.8	2.3	3.4	4.1	5.6
	6-19 times	0.1	0.3	0.6	1.0	1.4	1.3	1.7
Smokeless Tobacco	20-40 times	0.0	0.1	0.3	0.4	0.9	0.9	0.8
	40+ times	0.0	0.2	0.4	0.6	1.1	1.1	1.2
	No Answer	2.8	3.3	3.6	4.3	4.8	4.7	4.8
	Never	96.3	94.7	93.0	90.5	87.9	87.2	86.1
	1-5 times	0.7	1.2	2.0	2.8	3.2	3.3	3.5
Alcohol	6-19 times	0.1	0.4	0.5	0.8	1.2	1.2	1.4
	20-40 times	0.1	0.2	0.4	0.6	0.9	1.0	1.1
	40+ times	0.1	0.2	0.5	1.1	1.9	2.6	3.1
	No Answer	3.1	3.5	3.7	4.4	4.8	4.7	4.7
	Never	90.1	84.3	76.4	70.9	64.1	61.5	55.6
Marijuana	1-5 times	5.9	9.7	15.1	18.2	21.6	23.4	25.7
	6-19 times	0.7	1.6	2.9	4.2	6.1	6.8	9.4
	20-40 times	0.2	0.5	1.0	1.4	1.8	2.0	2.5
	40+ times	0.1	0.4	0.7	0.9	1.6	1.6	2.1
	No Answer	3.1	3.5	3.9	4.7	5.1	5.1	5.1
	Never	95.5	92.7	87.9	85.0	80.5	80.3	79.1
	1-5 times	1.0	2.3	4.6	5.6	6.9	6.8	7.5
	6-19 times	0.3	0.7	1.6	2.1	3.0	3.1	3.0
	20-40 times	0.1	0.4	0.9	1.3	1.8	1.8	1.9
	40+ times	0.1	0.4	1.1	1.5	2.7	2.8	3.4

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	3.1	3.6	3.9	4.9	5.3	5.3	5.3
	Never	96.7	95.8	95.0	93.7	92.8	92.6	92.3
	1-5 times	0.1	0.4	0.7	1.0	1.2	1.4	1.6
	6-19 times	0.0	0.1	0.2	0.2	0.4	0.5	0.5
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.1	0.2
	40+ times	0.0	0.0	0.1	0.1	0.1	0.2	0.2
Crack	No Answer	3.2	3.6	4.0	4.9	5.3	5.2	5.3
	Never	96.5	95.9	95.1	94.3	93.7	93.7	93.6
	1-5 times	0.2	0.3	0.5	0.6	0.6	0.7	0.6
	6-19 times	0.0	0.1	0.2	0.2	0.3	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.0	0.1
	40+ times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
Inhalants	No Answer	3.5	3.6	4.1	4.9	5.3	5.2	5.3
	Never	94.1	93.3	92.2	92.1	92.2	92.6	93.1
	1-5 times	1.8	2.2	2.8	2.2	1.8	1.6	1.1
	6-19 times	0.4	0.6	0.6	0.5	0.5	0.3	0.3
	20-40 times	0.1	0.2	0.2	0.2	0.1	0.1	0.1
	40+ times	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Amphetamines	No Answer	4.2	3.8	4.1	4.9	5.3	5.3	5.3
	Never	95.6	95.5	94.5	93.1	91.7	91.9	92.2
	1-5 times	0.2	0.4	0.9	1.3	1.9	1.9	1.6
	6-19 times	0.0	0.1	0.3	0.5	0.7	0.6	0.6
	20-40 times	0.0	0.0	0.1	0.1	0.2	0.2	0.2
	40+ times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
Methamphetamines	No Answer	3.9	3.7	4.1	5.0	5.3	5.2	5.4
	Never	96.0	95.8	95.2	94.2	93.7	93.7	93.6
	1-5 times	0.1	0.3	0.4	0.6	0.6	0.7	0.6
	6-19 times	0.0	0.1	0.2	0.1	0.2	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.0	0.0	0.1	0.1	0.1
Ritalin/Adderall	No Answer	4.3	3.9	4.1	4.9	5.4	5.3	5.3
	Never	95.3	95.1	93.9	92.2	91.0	91.5	91.7
	1-5 times	0.3	0.6	1.2	1.9	2.5	2.1	2.0
	6-19 times	0.1	0.2	0.4	0.6	0.7	0.7	0.6
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.2	0.2
	40+ times	0.0	0.1	0.1	0.1	0.1	0.2	0.1
Tranquilizers	No Answer	3.6	3.8	4.1	4.9	5.3	5.3	5.5
	Never	94.8	93.7	92.0	90.7	89.7	90.4	90.5
	1-5 times	1.1	1.8	2.8	3.0	3.4	2.9	2.7
	6-19 times	0.3	0.5	0.7	0.9	1.1	0.8	0.9
	20-40 times	0.1	0.1	0.3	0.3	0.4	0.4	0.3
	40+ times	0.0	0.1	0.1	0.2	0.2	0.2	0.1

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	4.1	3.9	4.1	4.9	5.3	5.3	5.5
	Never	95.6	95.1	93.8	92.2	90.9	90.7	90.7
	1-5 times	0.2	0.7	1.3	1.9	2.4	2.5	2.4
	6-19 times	0.1	0.2	0.5	0.6	0.9	0.8	0.9
	20-40 times	0.0	0.0	0.2	0.2	0.3	0.3	0.3
	40+ times	0.0	0.0	0.1	0.2	0.2	0.3	0.2
Psychedelics (All)	No Answer	4.0	3.7	3.9	4.7	5.1	5.1	5.2
	Never	95.8	95.7	94.9	93.9	93.4	93.2	93.1
	1-5 times	0.1	0.4	0.7	0.8	1.0	1.1	1.0
	6-19 times	0.0	0.1	0.2	0.3	0.3	0.3	0.4
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
LSD Only	No Answer	4.2	4.0	4.1	5.0	5.4	5.4	5.4
	Never	95.6	95.5	95.0	93.9	93.4	93.3	93.5
	1-5 times	0.1	0.3	0.6	0.7	0.8	0.9	0.6
	6-19 times	0.0	0.1	0.2	0.2	0.2	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
Other Psychedelics	No Answer	4.6	4.0	4.3	5.1	5.4	5.3	5.6
	Never	95.3	95.6	95.1	94.3	93.7	93.7	93.3
	1-5 times	0.1	0.2	0.3	0.4	0.6	0.7	0.7
	6-19 times	0.0	0.1	0.1	0.2	0.2	0.2	0.3
	20-40 times	0.0	0.1	0.1	0.0	0.1	0.1	0.1
	40+ times	0.0	0.0	0.0	0.0	0.1	0.0	0.1
Heroin	No Answer	3.9	3.8	4.1	5.0	5.4	5.4	5.5
	Never	96.0	95.7	95.3	94.3	93.9	93.8	93.7
	1-5 times	0.1	0.2	0.3	0.4	0.4	0.4	0.3
	6-19 times	0.0	0.1	0.2	0.1	0.2	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.0	0.1	0.1	0.1	0.1
Steroids	No Answer	3.6	3.8	4.2	5.1	5.5	5.4	5.5
	Never	96.1	95.6	95.1	94.1	93.6	93.9	93.8
	1-5 times	0.2	0.4	0.3	0.4	0.5	0.3	0.3
	6-19 times	0.1	0.1	0.2	0.2	0.2	0.2	0.2
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
Injected Drugs	No Answer	4.0	4.4	4.6	5.3	5.6	5.6	5.6
	Never	95.6	95.0	94.6	93.8	93.4	93.3	93.5
	1-5 times	0.2	0.3	0.5	0.5	0.4	0.4	0.4
	6-19 times	0.0	0.1	0.1	0.2	0.2	0.3	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.1	0.1	0.2	0.1	0.3	0.3	0.2

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	4.6	4.1	4.2	5.0	5.4	5.4	5.5
	Never	95.3	95.4	94.9	93.8	93.1	93.3	93.2
	1-5 times	0.1	0.4	0.6	0.8	1.0	0.9	0.9
	6-19 times	0.0	0.1	0.2	0.2	0.3	0.2	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.1	0.0	0.1	0.1	0.1
Rohypnol	No Answer	4.7	4.0	4.2	5.1	5.4	5.4	5.6
	Never	95.2	95.8	95.3	94.3	93.9	94.0	93.8
	1-5 times	0.0	0.2	0.3	0.4	0.4	0.4	0.3
	6-19 times	0.0	0.1	0.1	0.1	0.2	0.2	0.1
	20-40 times	0.0	0.0	0.0	0.1	0.1	0.0	0.1
	40+ times	0.0	0.0	0.0	0.0	0.0	0.0	0.1
GHB	No Answer	4.9	4.1	4.3	5.1	5.5	5.5	5.7
	Never	95.0	95.7	95.4	94.5	94.1	94.0	93.9
	1-5 times	0.0	0.1	0.2	0.3	0.2	0.3	0.2
	6-19 times	0.1	0.1	0.1	0.1	0.1	0.1	0.2
	20-40 times	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	40+ times	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Over the counter drugs	No Answer	4.3	4.2	4.6	5.5	5.9	5.8	5.8
	Never	93.3	92.4	90.2	89.0	88.2	89.5	89.9
	1-5 times	1.9	2.4	3.6	3.8	4.1	3.3	3.0
	6-19 times	0.4	0.6	1.0	1.1	1.1	0.8	0.8
	20-40 times	0.1	0.2	0.4	0.4	0.4	0.3	0.3
	40+ times	0.1	0.2	0.3	0.3	0.3	0.3	0.3

**Binge Drinking in the Past Two Weeks  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Binge drinking	No Answer	5.9	5.1	4.4	5.0	5.1	4.9	4.6
	None	89.2	86.6	82.4	78.1	73.2	71.9	66.8
	Once	2.7	4.0	6.0	7.4	8.8	9.2	10.6
	Twice	1.0	2.0	3.0	4.0	5.1	5.8	7.3
	3-5 times	0.8	1.3	2.3	3.2	4.6	4.8	6.8
	6-9 times	0.2	0.4	0.8	1.1	1.3	1.6	1.8
	10+ times	0.3	0.6	1.1	1.2	1.8	1.9	2.2

**Main Sources of Alcoholic Beverages in the Past Month  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Main sources of alcoholic beverages	No Answer	11.1	10.1	9.7	11.3	11.6	12.3	12.7
	No drink	81.9	77.9	71.3	66.1	60.2	57.7	51.9
	Liquor stores/ convenience stores/ supermarkets	0.1	0.1	0.2	0.2	0.7	0.9	1.6
	Restaurants/bars/ clubs	0.0	0.0	0.1	0.1	0.1	0.2	0.5
	Public events	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	Had someone else buy it	0.3	0.9	2.5	4.3	8.0	10.1	13.0
	Received from person 21 or older	0.9	1.5	3.0	4.0	5.5	6.5	8.8
	Took it from a store	0.1	0.2	0.3	0.4	0.5	0.4	0.3
	Family members	3.0	4.1	5.3	4.9	4.3	3.6	3.5
	Other ways	2.5	5.0	7.6	8.7	9.0	8.1	7.6

**Most Important Reasons for Drinking  
by Indiana 6th - 12th Graders, 2007**  
**(Values expressed as percentages, multiple responses allowed)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
Did not drink alcoholic beverages	80.6	70.8	58.5	51.8	43.1	40.1	34.7
To experiment	9.4	10.9	11.4	9.6	8.3	6.7	5.7
To relax or relieve tension	0.3	0.6	1.0	1.2	1.6	1.7	2.3
To feel good or get high	0.3	0.5	0.9	1.0	1.3	1.3	1.3
To seek deeper insights or understanding	0.1	0.1	0.2	0.3	0.3	0.3	0.4
To have a good time with friends	1.0	2.4	4.8	8.2	12.1	14.9	18.2
To fit in with a group I like	0.4	0.5	0.7	0.7	0.9	0.9	0.9
To get away from my problems	0.7	1.0	1.8	2.2	2.8	2.8	3.2
Because of boredom	1.0	1.8	3.2	3.9	5.2	5.1	5.4
Because of anger	0.9	1.6	2.5	3.0	3.5	3.4	3.1
To get through the day	0.2	0.4	0.6	0.6	0.7	0.5	0.6
To increase the effects of other drugs	0.1	0.2	0.4	0.7	1.0	1.0	1.1
To decrease the effects of other drugs	0.1	0.0	0.1	0.1	0.1	0.1	0.1
To get to sleep	0.3	0.5	0.7	0.8	1.1	1.2	1.4
Because it tastes good	4.1	7.6	12.0	14.4	16.4	18.3	20.1
Because I am hooked	0.6	1.1	1.4	1.5	1.8	1.7	1.6

**Consequences of ATOD Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had a hangover	No Answer	4.7	4.2	4.0	4.3	4.4	4.3	4.2
	Never	91.4	86.4	78.1	72.8	65.6	62.1	57.8
	Once	2.0	4.4	7.4	9.0	10.9	12.1	12.7
	Twice	0.9	1.9	3.9	4.9	6.6	7.3	7.7
	3-5 times	0.6	1.6	3.3	4.4	6.3	7.1	8.5
	6-10 times	0.2	0.5	1.3	1.7	2.4	2.7	3.9
	11 or more times	0.3	1.0	2.0	2.9	3.8	4.4	5.3
Had nausea or vomited	No Answer	5.1	4.6	4.4	4.7	4.8	4.6	4.5
	Never	91.3	87.9	81.9	76.8	69.3	64.6	59.9
	Once	2.1	4.1	6.8	8.9	11.4	13.0	14.2
	Twice	0.7	1.6	3.1	4.0	5.5	6.8	7.7
	3-5 times	0.4	0.9	2.1	2.9	5.2	6.4	8.4
	6-10 times	0.2	0.4	0.8	1.3	1.9	2.3	3.0
	11 or more times	0.2	0.5	0.9	1.4	1.9	2.3	2.4
Had a memory loss	No Answer	5.3	4.8	4.7	5.1	5.2	5.0	5.0
	Never	92.6	90.1	85.4	81.1	75.0	73.3	71.1
	Once	1.2	2.6	4.5	5.6	7.4	7.9	8.2
	Twice	0.4	1.0	2.0	2.7	3.9	4.1	4.6
	3-5 times	0.2	0.7	1.5	2.6	4.0	4.7	5.5
	6-10 times	0.1	0.3	0.8	1.1	2.0	2.0	2.6
	11 or more times	0.2	0.4	1.1	1.7	2.6	3.0	3.0
Poor on school test	No Answer	5.4	4.9	4.7	5.1	5.2	4.9	4.9
	Never	92.2	90.6	88.5	87.3	86.3	87.6	88.1
	Once	1.1	1.8	2.6	2.5	2.4	2.2	2.1
	Twice	0.4	0.8	1.1	1.4	1.6	1.3	1.2
	3-5 times	0.4	0.8	1.2	1.4	1.8	1.7	1.6
	6-10 times	0.1	0.3	0.7	0.7	0.8	0.6	0.7
	11 or more times	0.3	0.7	1.2	1.5	1.8	1.7	1.3
Missed school	No Answer	5.5	5.0	4.7	5.1	5.2	5.0	4.9
	Never	92.6	91.3	89.8	88.6	87.1	87.9	87.2
	Once	0.7	1.4	2.0	2.2	2.7	2.2	2.6
	Twice	0.4	0.7	1.0	1.2	1.3	1.3	1.5
	3-5 times	0.4	0.7	1.0	1.2	1.7	1.5	1.6
	6-10 times	0.2	0.4	0.6	0.7	0.8	0.8	0.9
	11 or more times	0.2	0.5	1.0	1.0	1.2	1.2	1.2
Got into a fight	No Answer	5.4	4.8	4.8	5.2	5.2	5.0	5.0
	Never	90.5	88.0	83.9	82.2	79.0	79.1	78.7
	Once	1.9	3.2	4.7	5.3	6.6	6.5	6.6
	Twice	0.7	1.3	2.2	2.7	3.5	3.9	3.7
	3-5 times	0.6	1.2	2.0	2.1	2.9	2.9	3.4
	6-10 times	0.3	0.5	0.9	1.0	1.1	1.0	1.1
	11 or more times	0.6	0.9	1.5	1.6	1.7	1.6	1.4

**Consequences of ATOD Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Damaged property	No Answer	5.5	5.0	4.8	5.3	5.3	5.2	5.0
	Never	93.3	92.6	90.6	89.8	88.8	89.9	90.0
	Once	0.6	1.0	1.9	1.8	2.2	2.0	2.0
	Twice	0.3	0.4	0.8	1.0	1.3	0.9	1.0
	3-5 times	0.2	0.4	0.8	0.9	1.1	1.1	0.9
	6-10 times	0.1	0.2	0.4	0.4	0.4	0.3	0.4
	11 or more times	0.1	0.4	0.7	0.7	0.8	0.7	0.6
Had trouble with parents	No Answer	5.6	5.0	4.8	5.2	5.2	5.0	5.0
	Never	91.0	88.8	85.3	83.2	80.2	80.2	80.8
	Once	1.7	3.0	4.7	5.6	7.4	7.8	7.4
	Twice	0.5	1.1	1.7	2.1	2.7	2.7	3.0
	3-5 times	0.4	0.9	1.3	1.6	2.1	2.1	2.1
	6-10 times	0.1	0.3	0.6	0.7	0.8	0.7	0.7
	11 or more times	0.6	1.0	1.6	1.6	1.6	1.5	1.1
Had trouble with police	No Answer	5.6	5.0	4.9	5.3	5.4	5.4	5.1
	Never	92.8	91.8	90.0	88.9	87.9	87.9	88.3
	Once	0.7	1.5	2.2	2.8	3.7	4.0	4.4
	Twice	0.4	0.6	1.0	1.2	1.2	1.2	1.0
	3-5 times	0.3	0.5	1.0	0.9	1.0	0.7	0.6
	6-10 times	0.1	0.1	0.4	0.3	0.3	0.3	0.2
	11 or more times	0.2	0.4	0.6	0.6	0.6	0.6	0.4
Driven under the influence	No Answer	5.8	5.2	5.3	5.6	5.7	5.5	5.4
	Never	93.4	93.0	91.2	90.0	85.6	80.5	76.5
	Once	0.4	0.8	1.5	1.9	3.7	5.2	5.9
	Twice	0.2	0.3	0.7	0.9	1.7	2.6	3.2
	3-5 times	0.1	0.3	0.6	0.8	1.4	2.6	3.7
	6-10 times	0.1	0.1	0.3	0.3	0.6	1.3	1.8
	11 or more times	0.1	0.3	0.6	0.6	1.3	2.4	3.5
Rode with a drunk driver	No Answer	5.6	4.8	4.8	5.1	5.3	5.2	5.0
	Never	90.1	87.7	81.4	77.7	71.6	69.7	68.3
	Once	1.6	2.5	4.0	4.9	6.4	6.9	6.8
	Twice	0.7	1.3	2.4	3.0	4.1	4.2	4.3
	3-5 times	0.8	1.3	2.7	3.8	4.8	5.2	5.7
	6-10 times	0.3	0.7	1.3	1.5	2.2	2.4	2.9
	11 or more times	0.9	1.7	3.4	4.0	5.6	6.4	6.9
Forgot things you did while high	No Answer	5.7	5.1	5.1	5.4	5.5	5.4	5.2
	Never	91.9	88.6	82.8	77.9	71.6	69.8	68.3
	Once	1.4	3.0	5.1	6.7	8.0	8.5	8.5
	Twice	0.4	1.4	2.4	3.3	4.5	4.8	5.3
	3-5 times	0.2	0.9	2.3	3.1	4.8	5.4	5.8
	6-10 times	0.1	0.4	0.9	1.5	2.1	2.3	2.9
	11 or more times	0.2	0.6	1.5	2.1	3.4	3.8	4.0

**Consequences of ATOD Use  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had been told to cut down	No Answer	5.7	5.1	5.0	5.4	5.4	5.1	5.1
	Never	92.9	92.0	89.6	87.5	85.2	85.4	84.8
	Once	0.6	1.2	1.9	2.5	3.1	3.0	3.1
	Twice	0.2	0.6	0.9	1.3	1.8	1.8	2.1
	3-5 times	0.2	0.4	0.9	1.2	1.9	1.8	2.3
	6-10 times	0.1	0.2	0.5	0.6	0.8	0.8	0.9
	11 or more times	0.3	0.5	1.2	1.4	1.9	2.1	1.7
Used alcohol or drugs to relax or fit in	No Answer	5.6	5.0	4.9	5.4	5.3	5.2	5.2
	Never	91.3	88.7	84.0	80.5	76.7	75.5	74.4
	Once	1.5	2.6	4.3	5.2	5.9	5.9	5.7
	Twice	0.7	1.1	2.0	2.5	3.2	3.2	3.5
	3-5 times	0.5	1.1	2.0	2.7	3.6	4.1	4.4
	6-10 times	0.2	0.4	0.9	1.2	1.6	1.9	2.3
	11 or more times	0.4	0.9	1.9	2.6	3.6	4.3	4.5
Used alcohol or drugs alone	No Answer	5.6	5.0	4.8	5.3	5.3	5.2	5.0
	Never	90.5	87.2	81.1	78.2	74.5	75.1	75.1
	Once	1.8	3.0	5.0	5.5	6.2	5.7	5.7
	Twice	0.8	1.6	2.9	3.1	4.0	3.6	3.7
	3-5 times	0.6	1.5	2.7	3.5	4.2	4.1	4.3
	6-10 times	0.3	0.6	1.3	1.7	1.9	1.9	1.9
	11 or more times	0.4	1.0	2.1	2.7	3.9	4.3	4.2

**Perceived Risk of Selected Substance Use Behavior  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	4.0	4.0	4.2	5.0	5.1	5.1	5.2
	No Risk	8.8	8.2	7.7	6.8	5.8	5.2	5.0
	Slight Risk	14.2	12.5	12.5	10.8	9.7	9.0	7.9
	Moderate Risk	28.1	28.6	28.4	27.2	26.1	26.1	24.5
	Great Risk	44.9	46.7	47.2	50.1	53.3	54.6	57.4
Smoke marijuana occasionally	No Answer	4.3	4.3	4.4	5.2	5.3	5.2	5.3
	No Risk	8.8	9.6	11.6	13.7	15.6	16.3	17.5
	Slight Risk	11.1	12.8	17.5	20.0	23.9	26.2	27.4
	Moderate Risk	31.2	32.1	30.6	28.5	26.8	25.5	25.6
	Great Risk	44.5	41.3	35.9	32.6	28.4	26.7	24.1
Smoke marijuana regularly	No Answer	4.4	4.4	4.5	5.3	5.4	5.3	5.5
	No Risk	8.6	8.3	8.7	9.0	9.4	9.1	8.8
	Slight Risk	4.0	4.3	7.1	9.2	11.1	12.7	13.9
	Moderate Risk	13.9	14.2	17.2	18.8	21.3	23.2	24.6
	Great Risk	69.1	68.8	62.5	57.8	52.8	49.8	47.2
Use cocaine occasionally	No Answer	4.6	4.6	4.6	5.4	5.5	5.4	5.5
	No Risk	8.5	7.5	6.7	5.8	5.1	4.6	4.6
	Slight Risk	8.8	8.0	7.5	7.0	6.4	5.8	5.7
	Moderate Risk	31.1	31.4	30.2	28.7	25.8	25.1	22.5
	Great Risk	47.1	48.6	51.0	53.1	57.2	59.1	61.7
Use cocaine regularly	No Answer	5.7	5.7	6.0	6.5	6.6	6.6	6.6
	No Risk	8.5	7.3	6.5	5.4	4.8	4.2	4.1
	Slight Risk	3.4	2.5	2.5	2.2	1.9	1.7	1.5
	Moderate Risk	11.7	10.0	9.3	9.2	8.0	7.3	6.5
	Great Risk	70.7	74.5	75.7	76.6	78.8	80.3	81.2
Occasionally consume 1-2 alcoholic drinks	No Answer	5.2	5.0	5.2	5.8	6.0	5.7	5.7
	No Risk	21.0	22.8	27.4	29.2	30.5	33.6	36.2
	Slight Risk	35.9	37.0	36.7	35.5	36.4	36.1	35.9
	Moderate Risk	19.4	18.3	15.7	15.1	13.8	12.3	11.1
	Great Risk	18.5	16.8	15.0	14.4	13.5	12.2	11.1
Binge drink weekly	No Answer	4.5	4.5	4.8	5.5	5.6	5.4	5.6
	No Risk	9.8	9.3	9.7	9.3	8.7	8.7	8.8
	Slight Risk	11.3	12.2	14.9	16.1	16.7	18.1	19.2
	Moderate Risk	28.1	28.7	29.5	29.5	29.9	30.8	31.3
	Great Risk	46.3	45.3	41.1	39.7	39.1	37.1	35.1

**Perceived Peer Approval of Using Selected Substances  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	6.6	6.0	5.7	6.2	6.3	6.3	6.4
	Strongly Approve	3.5	3.5	3.4	3.5	3.7	3.3	3.0
	Approve	1.4	2.4	4.2	5.4	6.6	7.8	8.0
	Do Not Know	12.9	14.2	17.0	17.4	18.3	18.1	18.8
	Disapprove	13.6	14.6	16.5	17.7	18.9	19.9	21.7
Smoke marijuana occasionally	Strongly Disapprove	62.0	59.3	53.1	49.7	46.3	44.6	42.1
	No Answer	6.8	6.2	5.8	6.4	6.4	6.4	6.6
	Strongly Approve	3.6	3.8	4.5	4.8	6.0	5.3	5.7
	Approve	1.4	3.3	6.8	9.2	12.3	14.0	14.8
	Do Not Know	10.5	11.7	14.9	16.0	17.1	17.2	18.0
Smoke marijuana regularly	Disapprove	10.1	10.8	12.7	13.7	14.6	15.5	16.6
	Strongly Disapprove	67.6	64.2	55.3	50.0	43.6	41.6	38.4
	No Answer	6.9	6.3	5.9	6.4	6.6	6.6	6.7
	Strongly Approve	3.5	3.8	4.5	4.9	5.9	5.1	5.3
	Approve	1.0	1.9	4.2	5.7	7.6	8.3	8.2
Use cocaine occasionally	Do Not Know	9.9	11.2	13.9	14.9	15.5	15.5	15.5
	Disapprove	7.3	7.9	10.1	11.5	12.7	13.7	15.3
	Strongly Disapprove	71.4	68.8	61.4	56.6	51.8	50.7	49.0
	No Answer	7.0	6.3	6.0	6.5	6.6	6.6	6.7
	Strongly Approve	3.4	3.0	2.8	2.7	2.7	2.3	2.3
Use cocaine regularly	Approve	0.8	1.0	1.5	1.6	1.9	1.9	1.9
	Do Not Know	9.7	10.7	12.3	12.5	12.0	11.7	11.1
	Disapprove	9.8	10.0	11.1	11.7	11.5	11.5	11.3
	Strongly Disapprove	69.4	68.9	66.4	65.0	65.2	66.1	66.6
	No Answer	7.3	6.7	6.5	7.2	7.2	7.3	7.4
Occasionally consume 1-2 alcoholic drinks	Strongly Approve	3.4	3.0	2.9	2.7	2.7	2.2	2.2
	Approve	0.6	0.9	1.2	1.2	1.4	1.2	1.0
	Do Not Know	9.6	10.3	11.8	12.1	11.5	11.3	10.4
	Disapprove	6.6	6.9	8.1	8.8	8.5	8.4	8.3
	Strongly Disapprove	72.5	72.2	69.5	68.0	68.8	69.7	70.7
Binge drink weekly	No Answer	7.2	6.4	6.1	6.5	6.6	6.6	6.7
	Strongly Approve	4.0	5.0	6.4	7.9	9.6	10.1	12.2
	Approve	5.5	9.6	15.9	20.7	26.8	29.8	32.9
	Do Not Know	16.8	18.8	22.4	22.8	22.6	21.8	21.0
	Disapprove	14.1	14.0	13.5	13.1	11.8	11.1	10.0
	Strongly Disapprove	52.5	46.3	35.7	29.0	22.7	20.6	17.2
	No Answer	7.2	6.5	6.1	6.6	6.6	6.7	6.8
	Strongly Approve	3.8	4.3	5.2	6.6	7.8	7.8	9.1
	Approve	1.8	4.0	7.6	10.6	14.8	16.6	19.1
	Do Not Know	12.0	14.3	18.3	20.4	21.0	21.4	21.1
	Disapprove	10.1	11.1	13.4	13.7	14.1	14.2	14.6
	Strongly Disapprove	65.1	59.9	49.4	42.1	35.6	33.2	29.4

**Perceived Parental Approval of Using Selected Substances  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	6.6	6.0	5.7	6.4	6.4	6.5	6.6
	Strongly Approve	3.3	2.9	2.6	2.3	2.2	1.8	1.9
	Approve	0.5	0.6	1.0	1.3	1.9	2.4	2.2
	Do Not Know	4.3	4.5	4.7	5.1	5.6	5.7	6.1
	Disapprove	3.7	3.8	4.9	5.8	7.6	8.6	11.0
	Strongly Disapprove	81.6	82.1	81.1	79.1	76.3	74.9	72.2
Smoke marijuana occasionally	No Answer	6.8	6.2	5.8	6.5	6.5	6.5	6.7
	Strongly Approve	3.2	2.8	2.6	2.3	2.3	1.8	1.9
	Approve	0.3	0.6	1.1	1.3	2.1	2.2	2.3
	Do Not Know	3.9	4.3	4.8	5.2	5.8	5.8	6.2
	Disapprove	2.7	2.8	4.0	4.9	6.4	7.1	8.3
	Strongly Disapprove	83.1	83.3	81.6	79.8	76.8	76.5	74.6
Smoke marijuana regularly	No Answer	6.8	6.3	5.8	6.6	6.5	6.7	6.7
	Strongly Approve	3.2	2.8	2.6	2.3	2.3	1.8	1.9
	Approve	0.2	0.4	0.7	0.8	1.3	1.3	1.3
	Do Not Know	3.9	4.1	4.7	4.9	5.5	5.4	5.6
	Disapprove	2.2	2.2	3.5	4.1	5.2	5.6	6.2
	Strongly Disapprove	83.7	84.2	82.7	81.4	79.2	79.3	78.2
Use cocaine occasionally	No Answer	6.9	6.3	5.9	6.7	6.7	6.7	6.8
	Strongly Approve	3.2	2.7	2.5	2.1	2.0	1.6	1.7
	Approve	0.1	0.2	0.3	0.2	0.3	0.3	0.3
	Do Not Know	3.7	3.9	3.9	4.0	4.4	4.2	4.1
	Disapprove	2.5	2.3	2.8	3.0	3.6	3.6	3.8
	Strongly Disapprove	83.6	84.5	84.6	83.8	83.0	83.6	83.3
Use cocaine regularly	No Answer	7.0	6.4	6.1	6.8	6.8	6.8	7.0
	Strongly Approve	3.2	2.7	2.5	2.2	2.1	1.6	1.7
	Approve	0.1	0.2	0.3	0.2	0.3	0.3	0.2
	Do Not Know	3.8	3.9	3.9	4.0	4.3	4.1	4.1
	Disapprove	2.0	1.9	2.6	2.8	3.4	3.3	3.5
	Strongly Disapprove	83.9	84.9	84.8	84.0	83.2	83.8	83.4
Occasionally consume 1-2 alcoholic drinks	No Answer	7.0	6.4	6.0	6.7	6.6	6.6	6.8
	Strongly Approve	3.3	2.9	2.8	2.5	2.5	2.3	2.4
	Approve	3.3	4.1	5.5	6.0	7.7	9.2	12.7
	Do Not Know	7.3	8.3	9.4	10.0	11.0	11.1	13.0
	Disapprove	6.3	7.1	9.0	10.9	12.4	13.7	15.4
	Strongly Disapprove	72.8	71.2	67.3	63.9	59.7	57.1	49.8
Binge drink weekly	No Answer	7.1	6.5	6.2	7.0	6.9	7.0	6.9
	Strongly Approve	3.2	2.8	2.7	2.4	2.2	1.9	1.9
	Approve	0.7	0.9	1.3	1.5	2.3	2.7	3.5
	Do Not Know	4.8	5.4	6.2	7.0	7.7	7.9	9.4
	Disapprove	3.6	4.3	5.6	6.8	8.3	9.6	12.2
	Strongly Disapprove	80.6	80.2	78.0	75.4	72.5	70.9	66.0

**Days being Absent from School in the Past Year  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No Answer	2.1	1.4	1.0	0.7	0.6	0.5	0.4
Never	12.9	12.9	11.4	13.9	12.0	10.8	8.8
Once	10.8	11.7	10.6	11.9	10.9	10.1	8.7
Twice	17.4	17.0	15.6	15.5	15.4	14.5	13.4
3-5 days	33.8	33.2	33.2	31.9	32.4	33.5	33.2
6-10 days	15.6	16.0	17.9	16.9	17.8	19.0	21.1
11-19 days	5.2	5.3	6.7	5.7	6.7	7.5	9.1
20 or more days	2.1	2.5	3.6	3.5	4.3	4.1	5.3

**Participation in Some Form of Camp  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No participation in camps	72.9	76.9	78.8	81.3	82.2	81.2	83.0
Afternoons R.O.C.K. in Indiana	7.8	5.6	4.6	2.1	1.2	0.8	0.7
Youth Leadership Programs	8.3	10.0	10.2	9.8	9.1	9.2	8.6
S.A.D.D., S.T.A.N.D. or other prevention programs	9.2	6.0	5.2	5.5	5.9	6.6	6.0
Participated in 2+ types of camps	1.7	1.4	1.2	1.3	1.6	2.2	1.7

**Participation in Organized Family Events  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No Answer	4.3	3.7	3.1	2.9	2.6	2.3	2.4
Never	16.5	17.7	20.0	19.9	20.3	20.6	21.2
Once per Week	38.4	40.9	41.9	43.3	45.4	46.7	47.6
Twice per Week	19.8	19.2	18.4	18.5	17.5	17.3	16.1
3+ Times per Week	21.1	18.6	16.7	15.5	14.2	13.1	12.8

**Number of Days per School Year Spent in Various Afterschool Activities  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
School sports team	No Answer	6.4	6.0	6.0	7.4	7.4	7.2	7.6
	None	34.6	33.2	35.7	38.6	43.6	47.5	51.5
	1-9 Days	15.2	11.1	9.5	6.7	5.2	4.3	3.8
	10-29 Days	9.1	6.5	5.6	3.1	2.4	2.2	2.0
	30-59 Days	11.0	11.4	10.4	7.7	6.0	6.2	5.6
	60-89 Days	8.6	11.1	10.8	10.6	10.0	9.5	9.5
	90-129 Days	7.5	11.1	11.1	12.6	12.2	11.4	10.3
	130-180 Days	7.6	9.6	10.7	13.2	13.3	11.8	9.7
School arts programs	No Answer	7.6	7.3	7.0	8.2	8.3	8.0	8.5
	None	56.7	57.2	56.2	58.0	59.3	60.0	60.3
	1-9 Days	17.3	17.0	16.5	13.2	11.7	10.3	9.1
	10-29 Days	6.5	6.4	6.7	5.6	5.1	5.2	5.2
	30-59 Days	4.0	4.3	4.7	4.3	4.4	4.6	4.7
	60-89 Days	2.5	2.5	3.0	3.7	3.8	4.1	3.9
	90-129 Days	1.9	1.8	2.2	2.9	3.4	3.5	3.8
	130-180 Days	3.4	3.6	3.7	4.0	4.0	4.3	4.5
School club/intramurals	No Answer	8.7	8.4	8.1	9.3	9.0	8.5	9.0
	None	55.8	57.1	57.7	55.3	53.0	50.2	48.5
	1-9 Days	15.5	15.2	14.4	14.4	14.1	14.6	13.9
	10-29 Days	8.5	8.1	7.7	8.6	9.9	11.6	12.5
	30-59 Days	4.9	4.8	5.1	5.5	6.1	7.0	7.6
	60-89 Days	2.4	2.7	2.9	3.1	3.4	4.0	3.9
	90-129 Days	1.9	1.8	1.9	2.0	2.2	2.3	2.4
	130-180 Days	2.3	1.9	2.1	1.9	2.1	1.8	2.2
Youth centers (YMCA, etc)	No Answer	9.0	8.1	7.8	9.1	8.9	8.4	8.8
	None	66.0	67.0	68.2	70.2	71.4	72.7	74.5
	1-9 Days	11.8	11.3	10.9	10.0	9.1	8.7	7.6
	10-29 Days	4.7	4.9	4.7	4.0	4.0	3.9	3.7
	30-59 Days	2.9	3.6	3.2	2.8	2.9	2.8	2.4
	60-89 Days	2.0	2.0	2.0	1.6	1.6	1.6	1.5
	90-129 Days	1.4	1.4	1.5	1.1	1.0	1.0	0.7
	130-180 Days	2.2	1.7	1.9	1.2	1.1	0.9	0.9
Extended day programs	No Answer	10.3	9.2	8.8	9.9	9.7	9.2	9.4
	None	61.5	65.1	66.5	70.2	73.2	75.7	78.5
	1-9 Days	16.1	15.7	14.8	12.2	10.6	9.1	7.0
	10-29 Days	5.3	4.6	4.5	3.4	2.8	2.7	2.2
	30-59 Days	2.7	2.1	2.2	1.8	1.5	1.4	1.1
	60-89 Days	1.4	1.3	1.2	1.0	0.8	0.7	0.7
	90-129 Days	1.1	0.8	0.8	0.7	0.5	0.4	0.4
	130-180 Days	1.6	1.1	1.2	0.7	0.8	0.7	0.6

**Number of Days per School Year Spent in Various Afterschool Activities  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Supervised youth activities (4H, scouts)	No Answer	11.1	10.4	9.9	11.2	10.8	10.0	10.1
	None	53.2	56.8	60.9	64.1	66.0	67.9	70.4
	1-9 Days	11.3	9.7	8.9	8.0	7.6	7.4	6.5
	10-29 Days	6.8	6.7	5.9	5.3	5.3	5.2	4.9
	30-59 Days	6.4	6.4	5.5	4.8	4.3	4.1	3.6
	60-89 Days	4.2	4.2	3.8	3.1	2.7	2.6	2.1
	90-129 Days	3.2	2.8	2.4	1.9	1.6	1.4	1.2
	130-180 Days	3.8	3.1	2.7	1.7	1.7	1.4	1.3
Afterschool classes (art, sport, etc)	No Answer	9.9	9.2	8.7	9.8	9.7	9.1	9.2
	None	38.4	41.1	44.2	47.6	50.3	53.2	57.8
	1-9 Days	16.2	14.3	13.9	11.7	10.8	10.4	8.8
	10-29 Days	9.0	9.1	7.9	7.2	6.7	6.5	6.3
	30-59 Days	8.0	7.7	7.3	6.8	6.1	6.2	5.4
	60-89 Days	5.5	5.9	5.4	5.1	5.0	4.7	3.5
	90-129 Days	5.1	5.0	4.8	4.7	4.4	3.7	3.1
	130-180 Days	7.9	7.7	7.8	7.1	6.9	6.1	5.8
Socialization without adult supervision	No Answer	8.1	7.4	6.8	7.9	7.8	7.5	8.0
	None	36.4	24.0	16.6	12.8	9.2	7.3	7.0
	1-9 Days	20.2	19.1	16.1	13.2	10.6	8.5	7.6
	10-29 Days	10.3	13.2	13.8	13.8	12.1	10.9	10.5
	30-59 Days	7.3	10.4	12.3	14.4	14.6	14.8	15.3
	60-89 Days	5.4	8.0	11.1	12.9	14.7	15.8	15.8
	90-129 Days	4.8	7.6	10.1	12.0	14.7	16.8	17.2
	130-180 Days	7.4	10.4	13.2	13.0	16.4	18.4	18.8
At home without adult supervision	No Answer	9.4	8.6	7.9	9.0	8.7	8.4	9.0
	None	30.5	20.9	15.8	12.4	10.2	9.3	8.3
	1-9 Days	27.0	23.6	19.5	16.6	13.8	13.0	11.7
	10-29 Days	11.8	14.8	15.8	15.7	15.6	15.5	14.8
	30-59 Days	7.2	10.7	13.1	15.2	15.6	16.5	16.4
	60-89 Days	4.9	7.5	9.7	11.8	13.4	14.0	14.6
	90-129 Days	3.7	6.2	8.3	9.6	11.4	11.8	12.0
	130-180 Days	5.5	7.7	9.9	9.7	11.3	11.5	13.2
At home with adult supervision	No Answer	9.7	9.0	8.5	9.5	9.1	8.7	9.2
	None	12.1	10.7	10.1	9.0	8.5	8.5	9.6
	1-9 Days	12.8	11.0	10.7	9.6	9.3	8.8	9.3
	10-29 Days	8.7	9.5	10.0	10.0	10.4	10.5	11.5
	30-59 Days	8.2	10.3	11.4	12.5	13.4	14.4	14.9
	60-89 Days	8.4	10.2	12.4	14.1	15.0	15.5	15.8
	90-129 Days	14.9	16.4	17.8	18.4	18.2	18.3	16.5
	130-180 Days	25.3	23.0	19.1	16.8	16.1	15.4	13.0

**Number of Days per School Year Spent in Various Afterschool Activities  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Work for pay outside home	No Answer	9.4	8.3	8.0	9.1	8.7	8.0	8.5
	None	52.1	47.1	44.8	44.9	38.1	28.9	22.4
	1-9 Days	17.5	17.9	16.6	13.0	9.4	6.4	4.9
	10-29 Days	8.6	10.8	11.5	10.7	8.8	7.4	6.6
	30-59 Days	4.9	6.6	8.2	9.1	10.1	10.1	10.2
	60-89 Days	2.6	3.8	4.6	5.8	9.5	12.7	13.3
	90-129 Days	1.8	2.4	2.8	3.8	8.0	13.5	15.9
	130-180 Days	3.2	3.1	3.4	3.6	7.5	13.1	18.2
Work for pay inside a home	No Answer	9.0	8.2	7.9	9.3	9.1	8.8	9.4
	None	40.3	39.2	40.0	43.4	47.3	51.9	56.4
	1-9 Days	20.2	19.3	18.0	16.0	14.8	14.5	13.2
	10-29 Days	11.7	12.4	12.2	11.4	10.5	9.3	8.5
	30-59 Days	6.9	8.1	8.5	8.1	7.0	6.4	5.2
	60-89 Days	3.9	4.4	4.7	4.4	4.1	3.6	2.6
	90-129 Days	2.7	3.0	3.0	3.0	2.8	2.2	1.9
	130-180 Days	5.4	5.5	5.6	4.5	4.4	3.4	3.0
Volunteer outside the home	No Answer	9.6	8.5	8.0	9.3	9.1	8.6	9.0
	None	50.8	49.3	49.5	49.1	48.1	46.3	44.1
	1-9 Days	19.4	20.0	19.1	18.1	17.3	18.0	17.2
	10-29 Days	8.8	9.5	10.0	10.4	11.5	12.6	14.0
	30-59 Days	4.7	5.5	5.8	6.1	6.4	7.2	7.8
	60-89 Days	2.3	3.0	3.3	3.1	3.4	3.6	3.9
	90-129 Days	1.7	1.9	2.1	2.0	2.0	1.8	1.9
	130-180 Days	2.6	2.4	2.3	1.8	2.2	1.8	2.1
Volunteer inside a home	No Answer	8.9	8.1	7.7	9.0	9.0	8.5	9.0
	None	41.2	40.1	41.7	42.3	43.2	43.9	45.4
	1-9 Days	20.8	20.0	18.7	16.9	15.2	15.0	14.2
	10-29 Days	10.2	11.1	10.3	10.2	10.2	10.1	10.3
	30-59 Days	6.4	7.1	7.3	7.7	7.5	8.3	7.6
	60-89 Days	3.8	4.3	4.5	4.6	4.8	4.5	4.5
	90-129 Days	3.1	3.4	3.5	3.5	3.6	3.3	3.1
	130-180 Days	5.6	6.0	6.2	5.9	6.6	6.2	5.8

Note: The number of school days in a year is estimated as 180 days.

**Perceived Safety in Neighborhood and in School  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Out alone during the day	No Answer	5.3	5.6	5.2	6.1	6.2	5.6	5.9
	Very Safe	49.3	56.2	60.5	65.7	68.7	73.7	76.3
	Somewhat Safe	33.2	30.0	27.6	23.1	20.6	17.3	14.7
	Somewhat Unsafe	8.2	5.7	4.6	3.2	2.7	1.9	1.7
	Very Unsafe	4.0	2.6	2.1	1.8	1.8	1.4	1.4
Out alone after dark	No Answer	5.6	5.8	5.4	6.3	6.4	5.7	6.1
	Very Safe	17.4	22.5	27.4	33.9	37.7	42.8	46.3
	Somewhat Safe	29.3	33.4	35.4	34.5	33.4	32.7	31.1
	Somewhat Unsafe	23.3	22.2	19.8	16.7	15.2	13.0	11.2
	Very Unsafe	24.3	16.2	12.1	8.6	7.4	5.8	5.4
Inside home alone during the day	No Answer	5.8	6.1	5.7	6.5	6.6	5.9	6.1
	Very Safe	65.3	71.6	76.0	78.1	79.3	81.7	82.7
	Somewhat Safe	21.9	17.7	14.8	12.5	11.3	10.1	9.0
	Somewhat Unsafe	4.4	2.9	2.0	1.5	1.5	1.1	1.0
	Very Unsafe	2.5	1.8	1.5	1.4	1.4	1.2	1.2
Inside home alone after dark	No Answer	5.9	6.0	5.7	6.5	6.5	6.0	6.1
	Very Safe	39.5	47.0	53.0	57.6	60.0	63.1	66.0
	Somewhat Safe	27.9	27.4	26.7	24.2	23.0	21.7	19.7
	Somewhat Unsafe	15.6	12.7	9.9	8.0	7.0	6.3	5.7
	Very Unsafe	11.0	6.9	4.8	3.8	3.4	2.9	2.5
Traveling to/from school	No Answer	5.8	6.0	5.6	6.6	6.6	6.0	6.2
	Very Safe	62.6	64.8	65.9	68.5	69.4	72.5	74.5
	Somewhat Safe	22.8	21.9	21.8	20.2	19.6	18.0	16.2
	Somewhat Unsafe	5.4	4.6	4.1	2.9	2.5	2.0	1.7
	Very Unsafe	3.5	2.7	2.6	1.8	1.8	1.6	1.5
Staying in class	No Answer	5.4	5.8	5.5	6.4	6.4	5.9	6.1
	Very Safe	81.9	78.7	74.9	72.8	69.9	72.3	72.9
	Somewhat Safe	9.7	12.0	15.1	16.5	18.6	17.1	16.5
	Somewhat Unsafe	1.4	1.6	2.2	2.0	2.4	2.2	2.3
	Very Unsafe	1.5	1.9	2.3	2.3	2.7	2.4	2.3
Alone in school hallways	No Answer	5.6	6.0	5.7	6.5	6.5	6.0	6.2
	Very Safe	70.6	68.3	69.1	65.7	65.5	68.9	70.1
	Somewhat Safe	18.2	18.8	18.6	20.7	20.7	18.9	18.0
	Somewhat Unsafe	3.6	4.4	3.9	4.1	4.2	3.4	3.1
	Very Unsafe	2.1	2.6	2.8	2.9	3.1	2.8	2.6
Staying for afterschool activities	No Answer	6.4	6.5	6.1	6.9	6.9	6.3	6.4
	Very Safe	69.9	70.3	69.6	68.3	67.0	70.0	70.9
	Somewhat Safe	18.0	17.2	18.3	18.9	19.8	18.0	17.3
	Somewhat Unsafe	3.5	3.4	3.2	3.1	3.1	2.9	2.7
	Very Unsafe	2.2	2.6	2.8	2.8	3.2	2.8	2.7

**Annual Prevalence of Weapon Use and Violent Behavior  
by Indiana 6th - 12th Graders, 2007**  
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Physical fight	No Answer	5.8	5.9	5.5	6.6	6.7	6.1	6.3
	Never	59.4	58.9	56.4	60.7	63.2	67.5	71.4
	1-2 times	22.5	22.7	25.0	22.3	20.7	18.8	16.5
	3-9 times	8.0	8.3	8.8	7.2	6.3	5.3	4.0
	10+ times	4.3	4.1	4.3	3.3	3.0	2.3	1.8
Shouting argument	No Answer	6.0	6.3	5.8	6.8	6.9	6.2	6.5
	Never	45.2	39.8	34.1	34.4	33.4	35.6	36.8
	1-2 times	25.4	25.1	25.7	25.8	25.3	25.5	26.3
	3-9 times	12.8	15.1	17.9	18.0	19.2	18.9	18.6
	10+ times	10.5	13.7	16.5	15.0	15.2	13.8	11.8
Carried gun for hunting	No Answer	6.2	6.3	5.9	7.0	7.0	6.2	6.4
	Never	72.8	71.3	70.4	70.2	71.2	73.2	73.8
	1-2 times	8.7	8.6	8.9	8.1	7.3	7.1	6.5
	3-9 times	5.6	6.0	6.2	6.2	5.9	5.5	5.6
	10+ times	6.7	7.8	8.6	8.6	8.5	8.0	7.7
Carried gun for other reasons	No Answer	6.2	6.3	6.0	7.0	7.0	6.3	6.5
	Never	89.4	87.9	86.5	85.7	85.4	87.0	86.9
	1-2 times	2.3	2.9	3.5	3.6	3.5	3.1	2.9
	3-9 times	0.9	1.2	1.6	1.5	1.6	1.4	1.5
	10+ times	1.2	1.7	2.4	2.2	2.6	2.2	2.2
Carried weapon for hunting	No Answer	6.3	6.4	6.0	7.1	7.1	6.4	6.6
	Never	81.2	79.3	78.6	78.5	78.6	80.6	81.1
	1-2 times	6.2	6.3	6.3	5.5	4.9	4.6	3.8
	3-9 times	3.0	3.4	3.7	3.5	3.4	3.0	3.1
	10+ times	3.3	4.6	5.3	5.4	5.9	5.4	5.4
Carried weapon for other reasons	No Answer	6.3	6.4	6.2	7.2	7.1	6.4	6.6
	Never	86.1	84.0	80.8	80.8	79.4	81.5	80.9
	1-2 times	4.0	4.6	5.8	5.2	5.4	4.7	4.3
	3-9 times	1.6	2.2	2.9	2.7	2.7	2.4	2.6
	10+ times	2.0	2.8	4.2	4.2	5.3	4.9	5.6
Carried gun to school	No Answer	6.1	6.3	6.0	7.0	7.0	6.3	6.5
	Never	93.2	92.5	92.3	91.4	91.1	91.8	91.7
	1-2 times	0.4	0.5	0.8	0.7	0.7	0.8	0.7
	3-9 times	0.1	0.2	0.3	0.2	0.3	0.3	0.3
	10+ times	0.2	0.5	0.7	0.6	0.9	0.8	0.8
Carried weapon to school	No Answer	6.1	6.4	6.0	7.1	7.1	6.4	6.5
	Never	91.3	89.9	88.2	87.5	86.4	87.8	87.5
	1-2 times	1.9	2.5	3.7	3.1	3.3	2.7	2.7
	3-9 times	0.3	0.6	0.9	1.0	1.2	1.0	1.0
	10+ times	0.3	0.7	1.2	1.4	2.1	2.0	2.2

**Annual Prevalence of Weapon Use and Violent Behavior  
by Indiana 6th - 12th Graders, 2007**  
**(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Threatened by weapon on school property	No Answer	6.3	6.5	6.2	7.2	7.2	6.4	6.7
	Never	87.6	86.5	85.4	85.0	85.4	87.9	88.7
	1-2 times	4.7	5.0	6.1	5.6	5.0	3.8	3.1
	3-9 times	0.8	1.1	1.3	1.1	1.3	1.0	0.7
	10+ times	0.6	0.8	1.1	1.0	1.1	0.9	0.8
Property stolen while at school	No Answer	6.4	6.7	6.3	7.4	7.3	6.5	6.6
	Never	68.6	63.4	60.7	60.3	61.8	65.7	68.7
	1-2 times	17.9	21.5	23.5	24.2	23.2	21.6	19.4
	3-9 times	5.0	6.0	6.8	6.3	5.8	4.7	4.1
	10+ times	2.1	2.3	2.7	1.9	2.0	1.5	1.2
Missed school: Felt unsafe on school property	No Answer	6.3	6.6	6.1	7.1	7.2	6.4	6.6
	Never	89.9	88.8	88.3	88.2	86.4	89.3	87.9
	1-2 times	2.8	3.4	4.0	3.2	4.5	2.8	4.1
	3-9 times	0.5	0.6	0.9	0.7	1.0	0.7	0.6
	10+ times	0.5	0.6	0.8	0.8	1.0	0.8	0.7
Missed school: Felt unsafe on way to/from school	No Answer	6.2	6.5	6.0	7.1	7.2	6.6	6.7
	Never	91.4	91.0	90.7	90.1	89.5	90.8	90.5
	1-2 times	1.7	1.6	2.0	1.7	1.9	1.4	1.6
	3-9 times	0.3	0.4	0.5	0.4	0.5	0.4	0.5
	10+ times	0.4	0.5	0.8	0.7	0.9	0.8	0.7

**Gambling during the Last Twelve Months  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Played cards	No Answer	5.3	5.4	5.4	6.8	7.0	6.6	6.8
	Not at all	79.1	74.5	69.3	67.1	64.7	65.5	63.2
	Less than once a month	9.1	11.6	14.4	14.8	15.7	15.5	16.8
	About once a month	3.9	5.1	6.6	7.1	7.9	7.6	8.5
	About once a week	1.8	2.3	2.9	3.1	3.5	3.5	3.4
Bet on games	Daily	0.8	1.0	1.4	1.1	1.3	1.4	1.2
	No Answer	5.5	5.5	5.5	7.0	7.2	6.7	7.0
	Not at all	77.1	72.5	68.1	66.6	66.2	68.0	68.0
	Less than once a month	10.0	12.2	14.0	14.1	14.1	13.1	13.5
	About once a month	4.0	5.7	7.0	7.3	7.3	7.2	7.1
Bet on sports	About once a week	2.2	2.8	3.4	3.3	3.7	3.4	3.2
	Daily	1.1	1.4	1.9	1.5	1.5	1.6	1.4
	No Answer	5.9	5.8	6.0	7.4	7.5	7.1	7.4
	Not at all	76.1	72.1	68.1	67.0	67.8	69.6	69.4
	Less than once a month	11.0	13.0	14.7	14.4	13.7	13.4	13.8
Bought lottery tickets	About once a month	3.8	5.1	6.3	6.8	6.4	5.6	5.7
	About once a week	1.9	2.6	3.1	3.0	3.2	2.9	2.5
	Daily	1.3	1.4	1.8	1.4	1.5	1.4	1.2
	No Answer	6.0	5.8	5.9	7.4	7.5	7.1	7.2
	Not at all	80.7	79.3	78.0	78.2	78.5	78.2	61.0
Gambled in a casino	Less than once a month	8.0	9.1	9.4	8.6	8.2	8.6	19.0
	About once a month	3.1	3.3	3.6	3.4	3.1	3.2	7.6
	About once a week	1.5	1.7	1.9	1.6	1.7	1.9	3.7
	Daily	0.7	0.8	1.1	0.8	0.9	1.0	1.5
	No Answer	6.0	5.9	5.9	7.3	7.5	7.1	7.2
Played online	Not at all	93.0	92.6	92.2	90.7	90.2	90.3	89.5
	Less than once a month	0.6	0.7	0.9	0.9	1.0	1.3	1.9
	About once a month	0.1	0.3	0.3	0.4	0.4	0.4	0.5
	About once a week	0.1	0.1	0.2	0.2	0.3	0.3	0.3
	Daily	0.2	0.4	0.5	0.5	0.6	0.6	0.6
	No Answer	5.9	5.9	5.9	7.5	7.5	7.1	7.3
	Not at all	90.8	89.8	89.4	88.3	88.1	89.0	88.8
	Less than once a month	1.6	2.1	2.1	1.9	1.9	1.4	1.5
	About once a month	0.5	0.9	1.0	0.9	0.9	0.8	0.8
	About once a week	0.5	0.6	0.6	0.6	0.6	0.7	0.6
	Daily	0.7	0.7	1.0	0.8	0.9	0.9	0.9

**Gambling during the Last Twelve Months  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Felt bad about the amount bet	No Answer	6.6	6.4	6.1	7.8	7.7	7.3	7.5
	Yes	5.6	5.1	5.0	4.1	3.9	3.4	3.7
	No	19.1	24.3	29.3	30.0	30.9	30.4	33.7
	I do not bet for money	68.7	64.2	59.6	58.2	57.5	58.9	55.1
Would like to stop betting but could not	No Answer	6.6	6.5	6.2	7.8	7.7	7.4	7.5
	Yes	4.9	4.0	3.5	2.6	2.1	1.7	1.5
	No	19.5	24.6	30.0	30.7	32.1	31.6	35.2
	I do not bet for money	69.1	64.9	60.3	58.9	58.2	59.3	55.8

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	3.0	3.1	2.8	3.1	2.9	2.4	2.4
	Never	86.0	79.6	71.1	64.3	57.3	53.2	47.8
	7 years or younger	2.1	2.3	2.9	2.6	2.4	2.3	1.6
	8 years	1.2	1.5	1.7	1.7	1.8	1.3	1.2
	9 years	1.2	1.4	1.7	1.5	1.6	1.3	1.2
	10 years	2.2	2.4	2.7	2.4	2.3	1.9	1.6
	11 years	2.2	3.1	3.2	3.0	2.6	2.0	1.8
	12 years	1.7	4.1	4.8	4.6	4.6	4.1	3.7
	13 years	0.4	2.2	5.9	5.9	5.5	4.7	4.4
	14 years		0.3	2.7	6.9	5.9	4.9	4.5
	15 years			0.5	3.6	8.8	8.3	6.7
	16 years				0.4	3.8	9.6	10.2
	17 years					0.3	3.6	8.3
	18 years or older						0.5	4.7
Cigars	No Answer	2.1	2.4	2.5	3.1	3.1	3.1	3.1
	Never	92.4	87.7	81.3	73.4	63.8	56.8	47.9
	7 years or younger	0.8	1.1	1.1	1.1	1.0	1.0	0.7
	8 years	0.6	0.6	1.0	0.6	0.8	0.6	0.4
	9 years	0.6	0.7	0.9	0.7	0.9	0.7	0.5
	10 years	1.1	1.4	1.5	1.4	1.4	1.0	0.9
	11 years	1.3	1.6	1.9	1.8	1.4	1.1	0.9
	12 years	0.8	2.5	2.8	3.0	3.0	2.4	2.1
	13 years	0.2	1.8	4.2	4.1	4.4	3.7	2.8
	14 years		0.3	2.4	6.5	6.0	4.6	4.3
	15 years			0.4	3.9	9.1	9.1	6.9
	16 years				0.4	4.7	11.2	12.6
	17 years					0.3	4.0	9.7
	18 years or older						0.7	7.3
Pipe	No Answer	2.5	3.1	3.5	4.3	4.6	5.2	5.0
	Never	96.2	93.6	90.1	85.0	79.7	75.1	69.9
	7 years or younger	0.1	0.4	0.3	0.4	0.6	0.5	0.5
	8 years	0.1	0.2	0.3	0.2	0.3	0.3	0.2
	9 years	0.2	0.1	0.4	0.3	0.3	0.3	0.2
	10 years	0.3	0.4	0.5	0.6	0.6	0.5	0.3
	11 years	0.2	0.5	0.7	0.9	0.6	0.6	0.4
	12 years	0.3	1.0	1.2	1.3	1.5	1.1	1.1
	13 years	0.1	0.6	1.7	2.2	2.2	1.8	1.4
	14 years		0.1	1.1	2.9	3.0	2.3	1.9
	15 years			0.1	1.7	4.3	4.2	3.5
	16 years				0.2	2.2	5.9	5.9
	17 years					0.2	1.8	6.0
	18 years or older						0.2	3.9

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smokeless Tobacco	No Answer	2.0	2.3	2.3	2.7	3.2	3.2	3.1
	Never	94.8	91.7	87.6	81.1	73.4	69.6	65.1
	7 years or younger	0.5	0.7	0.8	0.9	1.1	0.9	0.9
	8 years	0.2	0.4	0.5	0.6	0.6	0.5	0.4
	9 years	0.3	0.4	0.4	0.5	0.6	0.5	0.4
	10 years	0.7	0.7	1.0	0.8	1.0	0.9	0.8
	11 years	0.8	0.9	0.9	1.0	0.7	0.7	0.7
	12 years	0.5	1.5	1.7	1.9	2.0	1.8	1.6
	13 years	0.2	1.2	2.8	3.0	2.7	2.4	1.9
	14 years		0.1	1.8	4.1	4.0	3.0	2.7
	15 years			0.2	3.0	7.0	6.1	5.2
	16 years				0.4	3.4	7.8	7.9
	17 years					0.2	2.4	6.2
	18 years or older						0.3	3.2
Alcohol	No Answer	3.6	4.0	4.1	4.3	4.3	4.4	4.1
	Never	75.3	66.2	56.6	49.2	39.6	35.9	29.9
	7 years or younger	4.3	4.0	3.7	2.8	3.1	2.4	2.4
	8 years	1.7	1.8	1.7	1.3	1.1	1.0	0.7
	9 years	2.2	1.8	1.7	1.3	1.1	0.6	0.5
	10 years	3.6	3.7	3.6	2.8	2.5	1.8	1.4
	11 years	4.7	4.4	3.7	2.7	2.2	1.6	1.2
	12 years	3.9	7.4	6.7	5.4	4.7	3.8	3.6
	13 years	0.7	5.9	10.4	9.5	7.5	6.1	5.4
	14 years		0.8	6.7	11.5	10.6	8.6	7.6
	15 years			0.9	8.2	15.4	13.7	12.5
	16 years				1.0	7.4	14.3	15.4
	17 years					0.5	5.3	10.3
	18 years or older						0.5	5.0
Marijuana	No Answer	2.4	2.7	2.8	3.3	3.8	4.1	3.8
	Never	94.6	89.6	82.4	76.2	66.9	63.7	58.9
	7 years or younger	0.2	0.4	0.6	0.6	0.7	0.5	0.5
	8 years	0.1	0.3	0.5	0.3	0.6	0.4	0.4
	9 years	0.3	0.3	0.5	0.6	0.6	0.5	0.3
	10 years	0.4	0.9	1.0	0.9	1.1	0.8	0.7
	11 years	0.9	1.3	1.6	1.4	1.4	1.0	0.9
	12 years	0.8	2.2	2.5	2.9	2.6	2.3	2.1
	13 years	0.2	2.1	4.5	4.0	4.4	3.7	3.3
	14 years		0.3	3.0	5.6	5.5	4.2	4.6
	15 years			0.5	3.9	7.9	7.3	7.2
	16 years				0.4	4.2	8.5	8.6
	17 years					0.2	2.7	6.2
	18 years or older						0.2	2.5

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	2.3	2.5	2.7	3.3	3.9	3.9	4.0
	Never	97.3	96.4	95.4	93.1	90.3	89.0	87.0
	7 years or younger	0.1	0.2	0.2	0.2	0.3	0.2	0.3
	8 years	0.0	0.0	0.1	0.0	0.1	0.1	0.1
	9 years	0.0	0.1	0.1	0.0	0.1	0.1	0.1
	10 years	0.1	0.1	0.1	0.1	0.2	0.1	0.0
	11 years	0.0	0.1	0.2	0.1	0.1	0.1	0.2
	12 years	0.1	0.3	0.3	0.4	0.3	0.2	0.2
	13 years	0.0	0.3	0.5	0.5	0.5	0.4	0.2
	14 years		0.1	0.4	1.0	0.9	0.6	0.5
	15 years			0.1	1.1	1.7	1.4	1.0
	16 years				0.1	1.5	2.5	2.3
	17 years					0.1	1.2	2.8
	18 years or older						0.1	1.4
Crack	No Answer	2.5	2.8	3.0	3.5	4.0	4.3	4.1
	Never	97.0	96.2	95.5	94.4	92.9	92.5	92.1
	7 years or younger	0.1	0.1	0.2	0.2	0.3	0.1	0.3
	8 years	0.0	0.0	0.1	0.1	0.0	0.1	0.0
	9 years	0.0	0.0	0.1	0.0	0.1	0.1	0.0
	10 years	0.1	0.1	0.1	0.0	0.1	0.1	0.1
	11 years	0.1	0.2	0.2	0.1	0.2	0.1	0.1
	12 years	0.1	0.2	0.2	0.2	0.2	0.1	0.1
	13 years	0.0	0.2	0.4	0.3	0.3	0.3	0.2
	14 years		0.0	0.3	0.5	0.5	0.3	0.2
	15 years			0.0	0.4	0.7	0.7	0.3
	16 years				0.1	0.6	0.8	0.9
	17 years					0.1	0.4	0.9
	18 years or older						0.1	0.6
Inhalants	No Answer	2.9	2.9	3.1	3.5	4.0	4.2	4.4
	Never	91.5	90.1	89.0	88.7	87.1	87.3	87.1
	7 years or younger	1.4	1.1	1.0	0.7	0.7	0.4	0.4
	8 years	0.5	0.5	0.5	0.3	0.2	0.2	0.2
	9 years	0.5	0.6	0.5	0.2	0.3	0.2	0.1
	10 years	0.9	0.9	0.9	0.4	0.5	0.3	0.4
	11 years	1.2	1.1	1.0	0.8	0.5	0.4	0.3
	12 years	1.0	1.6	1.3	1.1	0.8	0.8	0.6
	13 years	0.1	1.0	1.7	1.5	1.3	0.9	1.0
	14 years		0.2	0.9	1.6	1.6	1.2	0.9
	15 years			0.1	1.0	2.0	1.8	1.2
	16 years				0.1	0.9	1.5	1.5
	17 years					0.0	0.6	1.3
	18 years or older						0.1	0.5

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Amphetamines	No Answer	3.5	3.0	3.0	3.6	4.1	4.3	4.4
	Never	96.3	96.1	95.0	92.4	89.3	88.3	87.2
	7 years or younger	0.0	0.1	0.2	0.2	0.2	0.1	0.2
	8 years		0.0	0.1	0.1	0.1	0.1	0.0
	9 years	0.0	0.1	0.1	0.1	0.0	0.1	0.0
	10 years	0.0	0.1	0.1	0.2	0.2	0.1	0.1
	11 years	0.1	0.1	0.2	0.2	0.2	0.1	0.1
	12 years	0.1	0.3	0.3	0.5	0.4	0.3	0.2
	13 years		0.2	0.5	0.7	0.9	0.6	0.5
	14 years		0.0	0.5	1.1	1.2	1.0	0.8
	15 years			0.1	0.8	2.0	1.7	1.7
	16 years				0.1	1.2	2.2	2.3
	17 years					0.1	0.8	1.8
	18 years or older						0.1	0.5
Methamphetamines	No Answer	3.1	3.0	2.9	3.5	4.1	4.3	4.3
	Never	96.6	96.2	96.0	94.7	93.3	92.7	92.2
	7 years or younger	0.0	0.1	0.1	0.1	0.3	0.1	0.2
	8 years	0.0	0.1	0.1	0.0	0.0	0.1	0.0
	9 years	0.0	0.0	0.1	0.0	0.1	0.1	0.0
	10 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	11 years	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	12 years	0.1	0.2	0.2	0.2	0.2	0.1	0.1
	13 years	0.0	0.2	0.3	0.4	0.2	0.3	0.2
	14 years		0.0	0.2	0.5	0.5	0.3	0.3
	15 years			0.0	0.4	0.7	0.6	0.5
	16 years				0.0	0.4	0.7	0.9
	17 years					0.0	0.4	0.7
	18 years or older						0.1	0.5
Ritalin/Adderall	No Answer	4.0	3.1	3.2	3.7	4.2	4.4	4.5
	Never	95.3	95.0	93.7	90.7	87.0	86.2	84.5
	7 years or younger	0.1	0.2	0.2	0.2	0.3	0.1	0.3
	8 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	9 years	0.1	0.1	0.2	0.1	0.1	0.1	0.0
	10 years	0.1	0.3	0.2	0.1	0.2	0.2	0.1
	11 years	0.1	0.2	0.3	0.2	0.3	0.1	0.1
	12 years	0.2	0.5	0.4	0.5	0.5	0.4	0.4
	13 years	0.0	0.4	0.8	1.1	1.0	0.7	0.6
	14 years		0.0	0.7	1.5	1.3	1.3	0.8
	15 years			0.1	1.5	3.2	2.2	1.7
	16 years				0.2	1.6	3.0	2.9
	17 years					0.2	1.1	2.8
	18 years or older						0.1	1.1

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Tranquilizers	No Answer	3.2	3.0	2.8	3.5	4.0	4.3	4.3
	Never	93.9	93.4	91.8	89.3	85.8	85.6	84.2
	7 years or younger	0.3	0.3	0.3	0.2	0.3	0.2	0.3
	8 years	0.2	0.3	0.1	0.1	0.1	0.1	0.0
	9 years	0.3	0.1	0.3	0.2	0.1	0.1	0.0
	10 years	0.4	0.4	0.3	0.3	0.3	0.1	0.1
	11 years	0.8	0.5	0.5	0.4	0.3	0.2	0.2
	12 years	0.7	0.9	0.9	0.9	0.6	0.5	0.4
	13 years	0.2	1.1	1.6	1.3	1.3	0.9	0.7
	14 years		0.1	1.2	2.1	1.8	1.4	0.9
	15 years			0.2	1.7	3.3	2.5	1.9
	16 years				0.2	2.0	2.8	3.5
	17 years					0.1	1.2	2.6
	18 years or older						0.2	1.0
Narcotics	No Answer	3.5	3.0	2.9	3.5	4.0	4.1	4.5
	Never	95.9	95.4	93.5	90.5	86.5	84.9	82.6
	7 years or younger	0.0	0.1	0.1	0.1	0.2	0.1	0.3
	8 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	9 years	0.0	0.0	0.1	0.1	0.1	0.1	0.0
	10 years	0.1	0.2	0.2	0.2	0.2	0.2	0.1
	11 years	0.2	0.2	0.3	0.2	0.3	0.1	0.1
	12 years	0.1	0.4	0.6	0.6	0.5	0.3	0.3
	13 years	0.0	0.5	1.0	1.2	1.1	0.7	0.6
	14 years		0.1	1.0	1.6	1.7	1.5	1.0
	15 years			0.2	1.6	3.3	2.6	2.1
	16 years				0.3	1.9	3.6	3.6
	17 years					0.1	1.5	3.4
	18 years or older						0.2	1.4
LSD Only	No Answer	3.7	3.2	3.0	3.6	4.1	4.4	4.5
	Never	96.1	95.8	95.0	93.3	91.7	90.6	90.0
	7 years or younger	0.1	0.1	0.1	0.2	0.3	0.2	0.2
	8 years	0.0	0.0	0.1	0.0	0.1	0.1	0.1
	9 years		0.0	0.1	0.1	0.0	0.0	0.1
	10 years	0.0	0.1	0.2	0.1	0.2	0.1	0.1
	11 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	12 years	0.1	0.2	0.3	0.3	0.2	0.3	0.1
	13 years	0.0	0.3	0.5	0.4	0.4	0.4	0.3
	14 years		0.1	0.5	0.9	0.7	0.6	0.3
	15 years			0.2	0.9	1.2	1.1	0.8
	16 years				0.1	0.9	1.4	1.4
	17 years					0.1	0.7	1.3
	18 years or older						0.1	0.8

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Psychedelics	No Answer	4.4	3.2	3.2	3.8	4.3	4.3	4.7
	Never	95.4	96.3	95.8	94.5	92.8	92.1	90.3
	7 years or younger	0.0	0.1	0.1	0.1	0.2	0.1	0.2
	8 years	0.0	0.0	0.0	0.0	0.1	0.1	0.0
	9 years	0.0	0.0	0.0	0.0	0.1	0.1	0.0
	10 years	0.0	0.1	0.1	0.1	0.1	0.1	0.0
	11 years	0.0	0.1	0.1	0.1	0.1	0.1	0.0
	12 years	0.1	0.1	0.1	0.2	0.2	0.1	0.1
	13 years		0.1	0.2	0.3	0.3	0.2	0.2
	14 years		0.0	0.2	0.5	0.4	0.5	0.2
	15 years			0.0	0.3	0.8	0.7	0.6
	16 years				0.1	0.6	1.1	1.3
	17 years					0.0	0.4	1.5
	18 years or older						0.1	0.7
Heroin	No Answer	3.3	3.1	2.9	3.7	4.2	4.4	4.4
	Never	96.4	96.2	96.0	94.8	93.9	93.3	93.3
	7 years or younger	0.1	0.1	0.1	0.2	0.2	0.2	0.3
	8 years	0.0	0.0	0.1		0.1	0.0	0.1
	9 years	0.0	0.0	0.0	0.1	0.0	0.0	0.0
	10 years	0.0	0.1	0.1	0.1	0.1	0.1	0.0
	11 years	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	12 years	0.1	0.1	0.1	0.2	0.2	0.2	0.1
	13 years	0.0	0.2	0.3	0.2	0.2	0.2	0.2
	14 years		0.0	0.2	0.3	0.3	0.2	0.2
	15 years			0.0	0.3	0.4	0.4	0.3
	16 years				0.0	0.3	0.4	0.4
	17 years					0.0	0.3	0.5
	18 years or older						0.1	0.3
Steroids	No Answer	3.1	3.1	3.1	3.7	4.3	4.3	4.6
	Never	96.1	95.7	95.2	94.2	93.2	93.1	92.6
	7 years or younger	0.1	0.2	0.2	0.1	0.3	0.2	0.2
	8 years	0.1	0.0	0.0	0.0	0.1	0.1	0.0
	9 years	0.1	0.1	0.1	0.1	0.1	0.0	0.0
	10 years	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	11 years	0.2	0.2	0.1	0.1	0.1	0.1	0.0
	12 years	0.1	0.2	0.3	0.2	0.1	0.1	0.1
	13 years	0.0	0.3	0.4	0.3	0.2	0.2	0.1
	14 years		0.0	0.4	0.4	0.3	0.2	0.2
	15 years			0.0	0.6	0.5	0.2	0.3
	16 years				0.1	0.7	0.7	0.6
	17 years					0.1	0.5	0.7
	18 years or older						0.1	0.5

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	4.2	3.1	3.2	3.7	4.2	4.3	4.7
	Never	95.6	96.0	95.1	93.5	91.7	91.1	89.7
	7 years or younger	0.0	0.1	0.1	0.1	0.2	0.1	0.2
	8 years	0.0	0.0	0.1	0.0	0.1	0.1	0.0
	9 years	0.0	0.0	0.1	0.0	0.1	0.0	0.0
	10 years	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	11 years	0.0	0.2	0.1	0.2	0.1	0.1	0.0
	12 years	0.1	0.3	0.3	0.3	0.3	0.1	0.1
	13 years	0.0	0.2	0.4	0.4	0.4	0.3	0.3
	14 years		0.0	0.4	0.8	0.7	0.3	0.3
	15 years			0.1	0.8	1.2	0.9	0.6
	16 years				0.1	0.9	1.6	1.2
	17 years					0.1	0.8	1.6
	18 years or older						0.1	1.1
Rohypnol	No Answer	4.3	3.3	3.1	3.6	4.2	4.5	4.7
	Never	95.5	96.2	96.0	95.2	94.2	93.8	93.6
	7 years or younger	0.0	0.1	0.1	0.1	0.2	0.1	0.2
	8 years	0.0	0.0	0.0	0.0	0.1	0.0	0.0
	9 years	0.0	0.0	0.1	0.0	0.0	0.0	0.0
	10 years	0.0	0.1	0.1	0.1	0.1	0.1	0.0
	11 years	0.0	0.2	0.1	0.1	0.1	0.1	0.1
	12 years	0.0	0.1	0.1	0.2	0.1	0.1	0.1
	13 years	0.0	0.1	0.2	0.3	0.1	0.2	0.1
	14 years			0.1	0.2	0.3	0.1	0.1
	15 years			0.1	0.2	0.4	0.2	0.3
	16 years				0.0	0.2	0.4	0.3
	17 years					0.0	0.2	0.3
	18 years or older						0.1	0.2
GHB	No Answer	4.6	3.3	3.2	3.8	4.3	4.3	4.9
	Never	95.3	96.3	96.2	95.2	94.5	94.4	93.8
	7 years or younger	0.0	0.1	0.1	0.1	0.2	0.0	0.2
	8 years	0.0	0.0	0.0	0.0	0.0	0.1	0.0
	9 years	0.0	0.0	0.0	0.0	0.1	0.0	0.0
	10 years	0.0	0.0	0.0	0.1	0.1	0.1	0.0
	11 years	0.0	0.1	0.1	0.0	0.1	0.1	0.0
	12 years	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	13 years	0.0	0.0	0.1	0.2	0.1	0.2	0.1
	14 years			0.1	0.2	0.2	0.1	0.2
	15 years			0.0	0.1	0.2	0.2	0.1
	16 years				0.0	0.1	0.2	0.2
	17 years					0.2	0.2	0.2
	18 years or older						0.1	0.2

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Over the counter drugs	No Answer	4.1	3.6	3.6	4.0	4.6	4.8	4.9
	Never	92.8	92.2	90.4	88.1	85.1	84.5	82.8
	7 years or younger	1.0	0.9	1.0	0.8	0.9	0.4	0.7
	8 years	0.3	0.3	0.3	0.2	0.2	0.2	0.1
	9 years	0.2	0.3	0.3	0.1	0.2	0.1	0.1
	10 years	0.4	0.5	0.5	0.4	0.4	0.3	0.2
	11 years	0.5	0.6	0.5	0.4	0.4	0.3	0.2
	12 years	0.6	0.7	0.8	0.9	0.9	0.6	0.5
	13 years	0.1	0.8	1.4	1.1	1.2	0.9	0.8
	14 years		0.2	1.1	2.3	1.9	1.3	1.3
	15 years			0.3	1.4	2.6	2.4	2.0
	16 years				0.2	1.5	3.0	2.9
	17 years					0.1	1.0	2.6
18 years or older							0.2	1.0

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	0.6	0.5	0.6	0.6	0.4	0.5	0.4
	Never	87.5	81.1	71.9	65.5	58.6	54.2	48.8
	Once or Twice	8.9	11.3	14.6	14.8	16.0	15.7	17.4
	Occasionally	1.3	3.0	5.5	8.3	10.6	12.8	15.0
	Past Regularly	1.1	2.1	3.2	4.3	5.0	5.2	5.2
	Current Regularly	0.6	2.0	4.2	6.4	9.4	11.5	13.1
Cigars	No Answer	0.3	0.3	0.4	0.6	0.6	0.6	0.5
	Never	93.7	89.0	81.8	74.1	64.9	57.6	48.7
	1-5 times	4.8	7.5	11.6	14.7	16.9	19.0	20.6
	6-19 times	0.6	1.2	2.5	4.0	6.9	9.0	11.7
	20-40 times	0.2	0.8	1.2	2.5	4.2	5.9	7.6
	40+ times	0.4	1.3	2.4	4.0	6.5	8.0	10.9
Pipe	No Answer	0.6	0.6	0.8	1.2	1.3	1.2	1.3
	Never	97.8	95.4	91.8	87.1	81.7	77.6	72.6
	1-5 times	1.3	2.5	4.1	5.9	7.4	9.1	10.8
	6-19 times	0.2	0.7	1.4	2.1	3.1	4.3	5.9
	20-40 times	0.0	0.3	0.7	1.3	2.0	2.7	3.4
	40+ times	0.1	0.5	1.2	2.4	4.5	5.0	6.1
Smokeless Tobacco	No Answer	0.4	0.4	0.4	0.5	0.5	0.5	0.6
	Never	96.2	93.4	88.9	82.4	75.5	71.5	67.2
	1-5 times	2.7	4.3	6.5	8.8	11.2	12.5	13.8
	6-19 times	0.4	0.8	1.6	2.5	3.4	3.8	4.2
	20-40 times	0.2	0.4	0.8	1.6	2.4	2.6	2.7
	40+ times	0.2	0.9	1.9	4.2	7.1	9.1	11.6
Alcohol	No Answer	0.8	0.7	0.8	0.9	1.0	1.1	1.1
	Never	75.5	66.2	56.0	48.7	39.3	35.7	29.9
	1-5 times	19.4	23.2	25.7	24.6	23.4	21.2	18.9
	6-19 times	3.0	5.6	9.0	11.6	13.2	13.3	13.8
	20-40 times	0.7	2.1	3.8	6.0	8.2	9.7	10.6
	40+ times	0.7	2.2	4.7	8.3	14.8	19.0	25.7
Marijuana	No Answer	0.6	0.6	0.8	0.7	1.0	1.0	1.1
	Never	95.9	90.9	82.7	77.1	67.8	64.6	59.6
	1-5 times	2.4	4.5	7.5	8.8	11.1	12.1	12.9
	6-19 times	0.5	1.4	2.9	3.9	4.8	5.3	6.2
	20-40 times	0.2	0.8	1.7	2.5	3.2	3.7	4.5
	40+ times	0.4	1.8	4.4	7.0	12.1	13.3	15.8
Cocaine	No Answer	0.4	0.4	0.6	0.7	0.8	0.8	1.0
	Never	99.1	98.1	96.8	95.1	92.6	91.1	89.1
	1-5 times	0.4	0.8	1.5	2.6	3.7	4.2	5.2
	6-19 times	0.1	0.2	0.4	0.5	1.1	1.6	1.7
	20-40 times	0.0	0.2	0.2	0.4	0.7	1.0	1.2
	40+ times	0.1	0.2	0.5	0.6	1.1	1.3	1.7

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Crack	No Answer	0.6	0.5	0.7	0.7	0.7	0.7	0.8
	Never	98.8	98.1	97.2	96.7	95.6	95.3	94.8
	1-5 times	0.5	0.8	1.2	1.4	1.8	2.2	2.5
	6-19 times	0.1	0.2	0.3	0.4	0.6	0.6	0.5
	20-40 times	0.0	0.2	0.2	0.3	0.3	0.5	0.4
	40+ times	0.1	0.2	0.4	0.6	0.9	0.8	0.9
Inhalants	No Answer	0.9	0.7	0.6	0.8	0.7	0.8	0.9
	Never	92.5	91.1	89.7	90.0	88.6	88.9	89.4
	1-5 times	5.3	5.9	6.4	6.0	6.6	6.5	5.7
	6-19 times	0.8	1.3	1.7	1.7	1.9	1.8	1.9
	20-40 times	0.3	0.4	0.7	0.6	0.9	0.9	1.0
	40+ times	0.3	0.6	0.9	0.9	1.3	1.2	1.1
Amphetamines	No Answer	1.8	0.9	0.8	0.9	0.9	0.8	1.1
	Never	97.8	97.8	96.4	94.2	91.2	90.5	89.3
	1-5 times	0.2	0.7	1.4	2.3	3.5	3.7	4.0
	6-19 times	0.1	0.2	0.7	1.0	1.8	2.2	2.2
	20-40 times	0.0	0.2	0.3	0.6	1.0	1.1	1.4
	40+ times	0.0	0.2	0.4	0.9	1.6	1.7	2.0
Methamphetamines	No Answer	1.3	0.7	0.7	0.7	0.8	0.8	1.1
	Never	98.3	98.3	97.7	97.1	96.0	95.7	94.8
	1-5 times	0.2	0.5	0.8	1.2	1.5	1.7	2.0
	6-19 times	0.0	0.2	0.2	0.3	0.5	0.5	0.8
	20-40 times	0.0	0.1	0.2	0.2	0.4	0.4	0.4
	40+ times	0.1	0.2	0.4	0.4	0.7	0.9	1.0
Ritalin/Adderall	No Answer	2.1	1.0	0.9	1.0	1.0	0.9	1.1
	Never	96.9	96.7	95.2	92.2	88.7	87.9	86.4
	1-5 times	0.7	1.1	2.3	3.5	5.0	5.3	5.7
	6-19 times	0.1	0.5	0.7	1.5	2.2	2.6	3.1
	20-40 times	0.1	0.2	0.4	0.7	1.1	1.4	1.5
	40+ times	0.1	0.4	0.6	1.1	2.0	2.0	2.2
Tranquilizers	No Answer	1.2	0.8	0.7	1.0	1.0	0.9	1.1
	Never	95.2	94.8	92.7	90.5	87.1	87.0	85.6
	1-5 times	2.9	3.0	4.1	4.9	6.3	5.9	6.0
	6-19 times	0.4	0.6	1.3	1.7	2.5	3.1	3.4
	20-40 times	0.2	0.3	0.5	0.9	1.3	1.3	1.8
	40+ times	0.2	0.4	0.7	1.1	1.9	1.8	2.2
Narcotics	No Answer	1.8	0.8	0.8	0.9	1.0	0.8	1.3
	Never	97.4	97.0	94.5	91.9	87.8	86.2	84.0
	1-5 times	0.5	1.2	2.5	3.4	4.9	5.3	5.8
	6-19 times	0.1	0.5	1.0	1.6	2.6	3.3	3.8
	20-40 times	0.0	0.2	0.5	1.0	1.5	1.9	2.2
	40+ times	0.1	0.3	0.7	1.2	2.2	2.4	2.9

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Psychedelics (All)	No Answer	1.4	0.7	0.6	0.7	0.7	0.5	0.7
	Never	98.0	97.9	96.4	95.1	93.2	92.3	90.4
	1-5 times	0.4	0.8	1.7	2.3	3.3	4.3	5.3
	6-19 times	0.1	0.2	0.6	0.8	1.1	1.3	1.7
	20-40 times	0.0	0.2	0.3	0.4	0.7	0.5	0.6
	40+ times	0.1	0.2	0.5	0.7	1.0	1.2	1.2
LSD Only	No Answer	1.7	1.0	0.9	1.0	1.1	0.9	1.1
	Never	97.9	97.8	96.5	95.5	94.0	93.2	92.4
	1-5 times	0.2	0.7	1.5	2.1	2.8	3.7	4.0
	6-19 times	0.0	0.2	0.5	0.7	0.8	0.9	1.2
	20-40 times	0.0	0.1	0.3	0.3	0.5	0.4	0.5
	40+ times	0.1	0.2	0.4	0.5	0.8	1.0	0.8
Other Psychedelics	No Answer	2.5	1.2	0.9	1.0	1.1	0.9	1.2
	Never	97.1	98.1	97.7	96.8	95.5	95.0	93.0
	1-5 times	0.2	0.3	0.8	1.1	1.8	2.2	3.4
	6-19 times	0.1	0.1	0.3	0.4	0.7	0.9	1.1
	20-40 times	0.0	0.1	0.1	0.2	0.3	0.3	0.4
	40+ times	0.0	0.1	0.3	0.5	0.6	0.7	0.8
Heroin	No Answer	1.4	0.8	0.7	1.0	1.1	0.8	1.1
	Never	98.3	98.2	97.7	97.2	96.5	96.4	96.1
	1-5 times	0.2	0.5	0.8	0.9	1.0	1.2	1.2
	6-19 times	0.0	0.2	0.3	0.3	0.5	0.4	0.5
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.4	0.4
	40+ times	0.0	0.2	0.4	0.4	0.7	0.7	0.8
Steroids	No Answer	1.2	1.1	1.0	0.9	1.1	1.0	1.1
	Never	97.8	97.4	97.0	96.5	95.7	96.0	95.7
	1-5 times	0.7	0.9	1.1	1.2	1.3	1.2	1.2
	6-19 times	0.1	0.2	0.3	0.4	0.5	0.5	0.6
	20-40 times	0.0	0.1	0.1	0.3	0.3	0.4	0.5
	40+ times	0.1	0.3	0.6	0.7	1.1	0.9	1.0
Injected Drugs	No Answer	3.3	4.2	4.4	5.7	5.8	6.0	5.7
	Never	95.9	94.2	93.8	92.4	91.8	91.6	91.7
	1-5 times	0.5	0.8	0.8	0.9	1.0	0.9	1.0
	6-19 times	0.1	0.3	0.3	0.3	0.4	0.4	0.4
	20-40 times	0.0	0.2	0.1	0.2	0.2	0.3	0.2
	40+ times	0.1	0.2	0.5	0.6	0.8	0.8	0.9
MDMA	No Answer	2.3	1.1	0.8	0.9	1.0	0.9	1.1
	Never	97.4	97.8	96.9	95.6	94.1	93.6	92.6
	1-5 times	0.2	0.7	1.3	2.0	2.8	3.6	4.0
	6-19 times	0.0	0.2	0.4	0.6	0.8	0.8	1.0
	20-40 times	0.0	0.1	0.2	0.3	0.4	0.3	0.5
	40+ times	0.0	0.2	0.4	0.5	0.9	0.8	0.9

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Rohypnol	No Answer	2.6	1.2	1.0	1.0	1.2	0.9	1.2
	Never	97.2	98.1	97.9	97.5	96.8	97.0	96.6
	1-5 times	0.1	0.4	0.6	0.8	0.9	0.9	1.0
	6-19 times	0.1	0.1	0.2	0.2	0.4	0.4	0.3
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.2	0.3
	40+ times	0.0	0.1	0.2	0.3	0.5	0.5	0.5
GHB	No Answer	2.8	1.4	1.1	1.0	1.3	1.1	1.3
	Never	97.0	98.1	98.1	97.7	97.2	97.3	97.1
	1-5 times	0.1	0.2	0.4	0.6	0.6	0.7	0.7
	6-19 times	0.0	0.1	0.1	0.3	0.3	0.3	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.1	0.2
	40+ times	0.0	0.1	0.2	0.3	0.5	0.5	0.5
Over the counter drugs	No Answer	2.1	1.2	1.2	1.2	1.2	1.2	1.4
	Never	94.0	93.6	91.0	89.1	86.0	85.8	84.1
	1-5 times	2.5	3.0	4.3	5.1	6.4	6.4	7.1
	6-19 times	0.8	1.0	1.5	1.9	2.7	3.1	3.6
	20-40 times	0.2	0.5	0.6	1.0	1.2	1.0	1.4
	40+ times	0.5	0.8	1.3	1.7	2.4	2.4	2.4

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	1.4	1.2	1.1	1.2	1.1	1.3	1.0
	None	93.0	88.3	82.3	76.3	70.1	65.5	60.7
	Few times	3.8	6.3	9.3	12.1	15.0	16.5	20.0
	1-5 cigarettes/day	1.1	2.3	3.7	4.7	5.9	6.5	7.4
	1/2 pack/day	0.4	0.8	1.8	2.6	3.5	4.8	5.0
	1 pack/day	0.2	0.5	1.1	1.9	3.0	4.0	4.1
	1 & 1/2 pack/day	0.1	0.2	0.3	0.5	0.6	0.5	0.8
	2+ pack/day	0.1	0.3	0.4	0.7	0.8	1.0	1.0
Cigars	No Answer	2.6	3.4	3.5	4.5	4.8	5.0	4.9
	Never	93.5	89.1	83.7	76.8	68.4	62.2	53.1
	1-5 times	2.8	4.7	7.4	10.1	12.8	15.6	19.7
	6-19 times	0.6	1.2	2.0	3.3	6.0	7.5	9.5
	20-40 times	0.3	0.6	1.1	2.0	3.1	3.5	5.0
Pipe	40+ times	0.3	1.1	2.2	3.3	4.9	6.2	7.8
	No Answer	2.9	3.5	4.0	5.1	5.6	6.0	5.9
	Never	95.8	93.2	90.0	85.5	80.3	77.3	72.6
	1-5 times	0.8	1.7	3.1	4.4	6.0	7.2	9.9
	6-19 times	0.2	0.7	1.2	1.9	2.6	3.5	4.6
Smokeless Tobacco	20-40 times	0.1	0.3	0.7	1.1	1.9	2.2	2.8
	40+ times	0.1	0.5	1.0	1.9	3.5	3.8	4.3
	No Answer	2.5	3.2	3.4	4.4	4.8	4.9	4.9
	Never	95.1	92.0	88.0	82.0	75.9	72.9	70.0
	1-5 times	1.7	2.7	4.8	6.4	7.9	8.8	9.7
Alcohol	6-19 times	0.4	0.8	1.5	2.3	3.2	3.1	3.4
	20-40 times	0.2	0.3	0.7	1.4	2.1	2.1	2.3
	40+ times	0.2	0.8	1.6	3.5	6.1	8.2	9.7
	No Answer	2.9	3.4	3.8	4.4	4.9	5.0	5.0
	Never	81.5	72.5	62.8	54.8	45.7	41.2	35.5
Marijuana	1-5 times	12.5	16.9	20.8	21.5	22.1	21.9	20.5
	6-19 times	2.1	4.4	7.1	10.0	11.6	13.2	13.9
	20-40 times	0.5	1.4	2.6	4.3	6.9	7.7	9.4
	40+ times	0.5	1.4	2.9	4.9	8.8	11.0	15.7
	No Answer	2.9	3.5	3.6	4.8	5.2	5.3	5.6
	Never	94.6	89.9	83.4	77.9	70.7	68.7	65.9
	1-5 times	1.5	3.3	5.6	6.9	8.8	9.7	10.5
	6-19 times	0.5	1.2	2.7	3.5	4.2	4.4	4.5
	20-40 times	0.3	0.7	1.6	1.9	2.9	3.1	3.2
	40+ times	0.3	1.3	3.1	4.8	8.2	8.9	10.5

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	2.9	3.5	3.9	4.9	5.4	5.5	5.7
	Never	96.8	95.5	94.3	92.3	90.0	88.7	87.5
	1-5 times	0.2	0.6	1.0	1.6	2.5	3.3	3.8
	6-19 times	0.1	0.2	0.4	0.5	0.8	1.0	1.2
	20-40 times	0.1	0.1	0.2	0.2	0.7	0.7	0.9
	40+ times	0.0	0.1	0.3	0.4	0.5	0.8	0.9
Crack	No Answer	2.9	3.5	3.8	5.0	5.5	5.5	5.7
	Never	96.5	95.6	94.7	93.3	92.2	91.8	91.6
	1-5 times	0.3	0.5	0.7	0.8	1.1	1.4	1.4
	6-19 times	0.1	0.1	0.4	0.3	0.5	0.6	0.4
	20-40 times	0.1	0.1	0.1	0.2	0.4	0.2	0.4
	40+ times	0.0	0.1	0.3	0.3	0.3	0.5	0.4
Inhalants	No Answer	3.2	3.5	3.9	5.1	5.5	5.6	5.9
	Never	92.8	91.3	90.4	89.7	89.0	89.7	89.6
	1-5 times	2.7	3.4	3.6	3.3	3.4	2.6	2.8
	6-19 times	0.9	1.0	1.2	1.1	1.1	1.0	0.9
	20-40 times	0.3	0.4	0.5	0.4	0.5	0.5	0.4
	40+ times	0.2	0.3	0.4	0.4	0.4	0.6	0.4
Amphetamines	No Answer	4.0	3.7	4.0	5.1	5.6	5.6	5.9
	Never	95.7	95.4	94.0	91.6	89.2	89.1	88.5
	1-5 times	0.2	0.5	1.0	1.7	2.6	2.5	2.7
	6-19 times	0.0	0.2	0.5	0.8	1.3	1.2	1.6
	20-40 times	0.0	0.1	0.1	0.4	0.8	0.7	0.6
	40+ times	0.1	0.3	0.4	0.5	0.5	0.8	0.7
Methamphetamines	No Answer	3.5	3.7	4.0	5.0	5.6	5.8	5.9
	Never	96.3	95.7	94.9	93.5	92.5	91.9	91.5
	1-5 times	0.1	0.3	0.5	0.7	0.9	1.0	1.2
	6-19 times	0.0	0.1	0.3	0.4	0.4	0.5	0.5
	20-40 times	0.0	0.1	0.1	0.1	0.3	0.2	0.4
	40+ times	0.1	0.2	0.2	0.2	0.2	0.5	0.4
Ritalin/Adderall	No Answer	4.1	3.8	4.1	5.1	5.7	5.6	5.9
	Never	95.3	94.7	93.3	90.4	87.5	87.5	86.4
	1-5 times	0.4	0.7	1.4	2.3	3.5	3.5	4.2
	6-19 times	0.1	0.4	0.6	1.1	1.8	1.8	1.9
	20-40 times	0.1	0.2	0.3	0.6	0.7	0.8	0.8
	40+ times	0.1	0.3	0.3	0.4	0.9	0.9	0.8
Tranquilizers	No Answer	3.5	3.7	4.0	5.1	5.6	5.7	5.9
	Never	94.1	93.3	91.5	89.1	86.7	86.7	85.6
	1-5 times	1.8	1.9	2.5	3.4	4.1	3.9	4.5
	6-19 times	0.4	0.6	1.0	1.2	1.9	1.8	2.2
	20-40 times	0.1	0.2	0.5	0.6	0.9	0.9	0.8
	40+ times	0.1	0.3	0.4	0.5	0.8	0.9	0.9

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	4.0	3.7	4.1	5.1	5.5	5.7	5.9
	Never	95.5	94.8	92.8	89.8	87.0	85.6	84.4
	1-5 times	0.3	0.7	1.7	2.6	3.6	4.2	4.5
	6-19 times	0.1	0.4	0.8	1.1	1.9	2.1	2.5
	20-40 times	0.0	0.2	0.3	0.7	0.9	1.1	1.5
	40+ times	0.0	0.1	0.4	0.6	1.1	1.3	1.2
Psychedelics (All)	No Answer	3.7	3.5	3.7	4.8	5.0	5.1	5.4
	Never	95.9	95.4	94.0	92.2	91.1	90.1	88.8
	1-5 times	0.2	0.5	1.2	1.6	2.1	2.7	3.7
	6-19 times	0.1	0.2	0.5	0.7	0.8	0.7	0.9
	20-40 times	0.0	0.2	0.2	0.3	0.4	0.5	0.4
	40+ times	0.2	0.3	0.4	0.5	0.6	0.9	0.8
LSD Only	No Answer	4.0	3.8	4.1	5.2	5.5	5.7	5.9
	Never	95.8	95.3	94.1	92.4	91.5	90.7	90.3
	1-5 times	0.1	0.5	1.0	1.3	1.8	2.1	2.4
	6-19 times	0.0	0.1	0.5	0.6	0.6	0.6	0.7
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.3	0.3
	40+ times	0.0	0.1	0.2	0.3	0.4	0.6	0.4
Other Psychedelics	No Answer	4.6	4.1	4.1	5.3	5.7	5.4	5.9
	Never	95.2	95.2	94.8	93.1	92.0	91.5	90.0
	1-5 times	0.1	0.2	0.4	0.7	1.2	1.6	2.6
	6-19 times	0.0	0.1	0.2	0.4	0.5	0.5	0.7
	20-40 times	0.0	0.1	0.1	0.2	0.3	0.4	0.3
	40+ times	0.1	0.2	0.3	0.3	0.4	0.6	0.6
Heroin	No Answer	3.6	3.8	4.1	5.1	5.5	5.7	6.1
	Never	96.2	95.5	94.8	93.6	93.0	92.4	92.2
	1-5 times	0.1	0.3	0.5	0.6	0.7	0.7	0.7
	6-19 times	0.0	0.1	0.2	0.3	0.2	0.4	0.3
	20-40 times	0.0	0.1	0.2	0.1	0.3	0.3	0.3
	40+ times	0.0	0.2	0.2	0.2	0.3	0.4	0.4
Steroids	No Answer	3.4	3.7	4.2	5.3	5.6	5.7	6.0
	Never	95.9	95.1	94.3	93.0	92.4	92.1	91.8
	1-5 times	0.4	0.7	0.7	0.8	0.8	0.7	0.9
	6-19 times	0.1	0.2	0.3	0.2	0.4	0.5	0.5
	20-40 times	0.1	0.1	0.1	0.2	0.3	0.4	0.3
	40+ times	0.1	0.2	0.4	0.4	0.5	0.6	0.6
Injected Drugs	No Answer	4.9	5.8	5.8	7.0	7.3	7.5	7.4
	Never	94.6	93.2	92.8	91.4	90.8	90.5	90.6
	1-5 times	0.4	0.6	0.6	0.8	0.7	0.6	0.8
	6-19 times	0.1	0.2	0.2	0.3	0.3	0.6	0.4
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.3	0.3
	40+ times	0.0	0.1	0.3	0.3	0.5	0.5	0.6

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	4.4	3.9	4.1	5.2	5.5	5.5	5.9
	Never	95.4	95.2	94.2	92.5	91.4	91.0	90.0
	1-5 times	0.1	0.5	0.9	1.3	1.9	2.2	2.7
	6-19 times	0.0	0.2	0.4	0.5	0.5	0.6	0.7
	20-40 times	0.0	0.1	0.1	0.3	0.4	0.3	0.2
	40+ times	0.1	0.2	0.3	0.4	0.4	0.4	0.5
Rohypnol	No Answer	4.6	4.0	4.2	5.2	5.7	5.5	5.9
	Never	95.3	95.4	95.0	93.7	93.1	93.1	92.8
	1-5 times	0.1	0.3	0.4	0.5	0.6	0.6	0.6
	6-19 times	0.0	0.1	0.2	0.3	0.2	0.2	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.2	0.2
	40+ times	0.0	0.1	0.2	0.1	0.2	0.4	0.3
GHB	No Answer	4.7	4.1	4.3	5.4	5.7	5.7	6.0
	Never	95.1	95.5	95.1	93.8	93.3	93.2	92.9
	1-5 times	0.1	0.2	0.3	0.3	0.4	0.4	0.5
	6-19 times	0.0	0.1	0.2	0.2	0.2	0.2	0.2
	20-40 times	0.0	0.0	0.1	0.1	0.2	0.2	0.2
	40+ times	0.0	0.1	0.1	0.1	0.1	0.3	0.2
Over the counter drugs	No Answer	4.2	4.2	4.5	5.7	6.0	6.4	6.5
	Never	92.8	92.0	90.1	87.7	85.8	85.3	84.9
	1-5 times	1.8	2.1	2.9	3.4	4.2	4.6	4.7
	6-19 times	0.7	0.9	1.3	1.6	2.0	1.9	2.0
	20-40 times	0.3	0.4	0.5	0.7	0.8	0.7	1.0
	40+ times	0.2	0.4	0.7	0.8	1.2	1.0	1.0

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	1.2	1.1	1.0	1.1	1.1	1.2	0.9
	None	95.9	92.8	88.8	84.0	79.0	75.1	72.3
	Few times	1.8	3.1	4.7	6.6	8.3	9.3	10.8
	1-5 cigarettes/day	0.6	1.5	2.6	3.4	4.6	5.3	6.3
	1/2 pack/day	0.2	0.5	1.2	1.9	2.8	3.8	4.0
	1 pack/day	0.2	0.4	1.0	1.8	2.7	3.8	4.0
	1 & 1/2 pack/day	0.1	0.1	0.2	0.3	0.6	0.5	0.6
	2+ pack/day	0.1	0.4	0.5	0.8	0.9	1.0	1.1
Cigars	No Answer	3.5	4.1	4.5	5.8	6.1	6.0	6.0
	Never	94.7	91.6	88.2	83.2	77.3	74.4	68.4
	1-5 times	1.4	2.7	4.4	6.6	9.7	11.7	16.0
	6-19 times	0.2	0.7	1.0	1.8	3.0	3.1	4.3
	20-40 times	0.1	0.4	0.8	1.1	1.3	1.6	1.6
Pipe	40+ times	0.1	0.5	1.1	1.6	2.7	3.2	3.6
	No Answer	3.6	4.5	4.7	6.0	6.6	6.6	6.7
	Never	95.7	93.6	91.6	88.4	84.8	83.9	80.8
	1-5 times	0.6	1.1	2.0	2.8	4.1	5.0	7.1
	6-19 times	0.1	0.4	0.8	1.3	1.8	1.8	2.4
Smokeless Tobacco	20-40 times	0.0	0.1	0.4	0.6	1.2	1.0	1.1
	40+ times	0.0	0.2	0.5	0.9	1.5	1.6	1.8
	No Answer	3.4	4.2	4.4	5.3	5.9	6.0	6.0
	Never	95.3	92.9	90.6	86.0	81.5	79.6	77.3
	1-5 times	0.9	1.7	2.9	4.3	5.3	5.5	6.0
Alcohol	6-19 times	0.2	0.5	0.9	1.5	2.2	2.2	2.7
	20-40 times	0.1	0.3	0.6	1.0	1.6	1.8	2.1
	40+ times	0.1	0.4	0.7	1.9	3.6	5.0	5.9
	No Answer	3.7	4.3	4.5	5.5	5.8	6.0	6.0
	Never	89.8	83.9	77.6	70.6	62.5	59.2	52.5
Marijuana	1-5 times	5.6	9.2	13.0	16.6	20.1	22.1	24.1
	6-19 times	0.6	1.6	2.8	4.6	6.9	7.9	11.0
	20-40 times	0.2	0.5	1.1	1.5	2.3	2.6	3.2
	40+ times	0.2	0.5	0.9	1.2	2.4	2.3	3.2
	No Answer	3.7	4.2	4.7	6.0	6.3	6.4	6.5
	Never	94.7	91.8	87.0	83.1	78.4	77.9	75.6
	1-5 times	1.0	2.2	4.3	5.3	6.4	6.6	7.2
	6-19 times	0.3	0.8	1.6	2.3	3.3	3.2	3.6
	20-40 times	0.1	0.5	1.1	1.5	2.0	1.9	2.1
	40+ times	0.1	0.6	1.3	1.9	3.7	4.0	4.9

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	3.8	4.4	4.9	6.2	6.7	6.7	6.8
	Never	96.0	94.9	94.0	92.1	90.9	90.5	90.0
	1-5 times	0.1	0.4	0.6	1.1	1.4	1.7	1.9
	6-19 times	0.0	0.2	0.3	0.3	0.5	0.7	0.7
	20-40 times	0.0	0.0	0.1	0.2	0.2	0.2	0.2
	40+ times	0.0	0.0	0.2	0.1	0.2	0.2	0.2
Crack	No Answer	3.9	4.6	4.9	6.3	6.6	6.7	6.9
	Never	95.8	94.9	94.2	92.7	92.0	91.9	91.7
	1-5 times	0.2	0.4	0.4	0.6	0.8	0.9	0.8
	6-19 times	0.0	0.1	0.3	0.3	0.4	0.3	0.4
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.1	0.1	0.2	0.1	0.1
Inhalants	No Answer	4.1	4.5	5.0	6.2	6.7	6.7	6.9
	Never	93.6	92.7	92.2	91.1	90.9	91.0	91.1
	1-5 times	1.7	1.9	1.9	1.9	1.6	1.6	1.3
	6-19 times	0.4	0.6	0.5	0.5	0.5	0.4	0.5
	20-40 times	0.1	0.1	0.2	0.2	0.1	0.2	0.1
	40+ times	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Amphetamines	No Answer	4.9	4.7	5.1	6.2	6.6	6.7	6.9
	Never	94.8	94.8	93.7	92.0	90.6	90.7	90.5
	1-5 times	0.2	0.3	0.7	1.1	1.7	1.5	1.6
	6-19 times	0.0	0.2	0.3	0.5	0.7	0.7	0.7
	20-40 times	0.0	0.0	0.1	0.1	0.2	0.2	0.2
	40+ times	0.1	0.1	0.1	0.1	0.1	0.2	0.1
Methamphetamines	No Answer	4.6	4.6	5.0	6.3	6.6	6.5	7.1
	Never	95.3	95.0	94.4	92.9	92.3	92.1	91.6
	1-5 times	0.1	0.2	0.3	0.5	0.6	0.8	0.8
	6-19 times	0.0	0.1	0.2	0.2	0.3	0.3	0.4
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.2	0.1
	40+ times	0.0	0.0	0.1	0.0	0.1	0.1	0.1
Ritalin/Adderall	No Answer	5.0	4.8	5.0	6.3	6.8	6.7	6.9
	Never	94.6	94.3	93.3	91.3	89.6	90.1	89.8
	1-5 times	0.2	0.5	1.0	1.5	2.3	2.0	2.2
	6-19 times	0.1	0.2	0.4	0.6	0.8	0.7	0.7
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.3	0.2
	40+ times	0.0	0.1	0.1	0.1	0.2	0.2	0.2
Tranquilizers	No Answer	4.3	4.7	5.0	6.3	6.6	6.7	7.1
	Never	94.5	93.7	92.7	90.5	89.4	89.7	89.0
	1-5 times	0.9	1.1	1.5	2.1	2.5	2.2	2.5
	6-19 times	0.2	0.2	0.5	0.7	0.9	0.7	1.0
	20-40 times	0.1	0.1	0.2	0.2	0.3	0.4	0.3
	40+ times	0.0	0.1	0.2	0.2	0.2	0.3	0.2

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	4.8	4.8	5.0	6.2	6.6	6.8	7.1
	Never	94.9	94.3	93.3	90.9	89.7	88.9	88.0
	1-5 times	0.2	0.6	1.0	1.7	2.1	2.8	2.9
	6-19 times	0.1	0.3	0.4	0.8	1.0	1.0	1.3
	20-40 times	0.0	0.0	0.1	0.2	0.4	0.3	0.4
	40+ times	0.0	0.0	0.1	0.2	0.2	0.3	0.3
Psychedelics (All)	No Answer	4.6	4.6	4.7	6.0	6.4	6.5	6.7
	Never	95.2	94.8	94.0	92.3	91.6	91.2	90.7
	1-5 times	0.2	0.3	0.7	1.0	1.2	1.5	1.6
	6-19 times	0.0	0.2	0.3	0.4	0.4	0.5	0.6
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.1	0.2
	40+ times	0.0	0.0	0.1	0.1	0.2	0.2	0.2
LSD Only	No Answer	4.9	4.9	4.9	6.3	6.7	6.9	7.0
	Never	95.0	94.6	94.0	92.2	91.7	91.3	91.3
	1-5 times	0.1	0.3	0.7	0.9	1.0	1.1	0.9
	6-19 times		0.1	0.3	0.3	0.3	0.4	0.4
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.2
	40+ times	0.0	0.0	0.1	0.1	0.1	0.2	0.1
Other Psychedelics	No Answer	5.5	4.9	5.1	6.4	6.8	6.8	7.2
	Never	94.4	94.7	94.2	92.8	92.0	91.9	91.0
	1-5 times	0.1	0.2	0.3	0.5	0.7	1.0	1.2
	6-19 times	0.0	0.1	0.2	0.2	0.3	0.3	0.4
	20-40 times		0.0	0.1	0.0	0.1	0.1	0.1
	40+ times	0.0	0.0	0.1	0.1	0.1	0.0	0.1
Heroin	No Answer	4.5	4.8	5.0	6.3	6.8	6.9	7.1
	Never	95.3	94.8	94.3	92.9	92.3	91.9	91.7
	1-5 times	0.1	0.2	0.3	0.4	0.4	0.5	0.4
	6-19 times	0.0	0.2	0.2	0.2	0.2	0.4	0.4
	20-40 times		0.0	0.1	0.1	0.2	0.1	0.2
	40+ times	0.0	0.0	0.0	0.1	0.1	0.1	0.2
Steroids	No Answer	4.4	4.7	5.1	6.5	6.7	6.9	7.1
	Never	95.2	94.5	93.9	92.3	91.8	91.8	91.6
	1-5 times	0.2	0.5	0.5	0.6	0.8	0.5	0.5
	6-19 times	0.1	0.2	0.2	0.3	0.4	0.4	0.3
	20-40 times	0.0	0.1	0.2	0.2	0.2	0.2	0.2
	40+ times	0.0	0.1	0.2	0.1	0.2	0.2	0.1
Injected Drugs	No Answer	5.0	5.9	6.0	7.1	7.5	7.7	7.5
	Never	94.7	93.5	93.1	91.7	91.2	90.7	91.0
	1-5 times	0.2	0.3	0.4	0.6	0.5	0.6	0.6
	6-19 times	0.0	0.1	0.2	0.2	0.3	0.3	0.3
	20-40 times		0.1	0.1	0.1	0.2	0.2	0.1
	40+ times	0.1	0.1	0.2	0.2	0.4	0.5	0.4

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	5.3	5.0	5.1	6.4	6.8	6.8	7.1
	Never	94.6	94.5	93.9	92.2	91.5	91.6	90.9
	1-5 times	0.1	0.4	0.6	1.0	1.0	1.0	1.2
	6-19 times	0.0	0.1	0.2	0.3	0.4	0.3	0.4
	20-40 times		0.0	0.1	0.1	0.1	0.1	0.2
	40+ times		0.0	0.1	0.1	0.1	0.1	0.1
Rohypnol	No Answer	5.6	4.8	5.1	6.5	6.7	6.9	7.3
	Never	94.4	94.8	94.3	92.9	92.5	92.2	91.9
	1-5 times	0.0	0.2	0.3	0.4	0.4	0.5	0.4
	6-19 times	0.0	0.1	0.2	0.1	0.2	0.2	0.2
	20-40 times		0.0	0.1	0.1	0.1	0.1	0.1
	40+ times		0.0	0.0	0.0	0.1	0.1	0.1
GHB	No Answer	5.7	5.0	5.1	6.4	6.8	7.0	7.4
	Never	94.2	94.8	94.4	93.0	92.6	92.3	92.0
	1-5 times	0.1	0.1	0.2	0.3	0.3	0.4	0.3
	6-19 times		0.1	0.2	0.2	0.2	0.2	0.3
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.1	0.0
	40+ times		0.0	0.0	0.0	0.0	0.0	0.0
Over the counter drugs	No Answer	5.1	5.1	5.5	6.8	7.2	7.4	7.5
	Never	92.9	92.4	91.1	89.3	88.2	88.3	88.3
	1-5 times	1.5	1.8	2.2	2.5	3.1	3.1	2.9
	6-19 times	0.3	0.5	0.7	0.8	0.9	0.7	0.9
	20-40 times	0.1	0.1	0.3	0.3	0.3	0.3	0.2
	40+ times	0.1	0.1	0.3	0.2	0.3	0.3	0.3

**Binge Drinking in the Past Two Weeks  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Binge drinking	No Answer	6.4	6.0	5.3	5.7	5.9	5.8	5.3
	None	88.6	85.7	82.2	76.9	70.0	67.2	61.5
	Once	2.8	3.8	5.4	7.0	8.9	9.8	10.4
	Twice	0.8	2.0	2.7	4.1	5.8	6.8	8.6
	3-5 times	0.9	1.3	2.2	3.4	5.3	5.7	8.2
	6-9 times	0.2	0.3	0.9	1.1	1.7	2.1	2.5
	10+ times	0.3	0.8	1.3	1.7	2.5	2.6	3.5

**Main Sources of Alcoholic Beverages in the Past Month  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Main sources of alcoholic beverages	No Answer	11.9	11.3	10.7	12.3	13.0	13.5	13.7
	No drink	81.1	76.8	71.6	65.4	57.9	54.7	48.8
	Liquor stores/ convenience stores/ supermarkets	0.1	0.2	0.3	0.4	1.2	1.6	2.6
	Restaurants/bars/ clubs	0.1	0.0	0.1	0.1	0.2	0.3	0.5
	Public events	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	Had someone else buy it	0.5	1.1	2.6	4.8	8.9	11.5	14.8
	Received from person 21 or older	0.9	1.4	2.6	3.5	4.9	5.5	7.2
	Took it from a store	0.1	0.3	0.5	0.6	0.7	0.7	0.5
	Family members	2.8	3.9	4.5	4.3	3.8	3.1	3.1
	Other ways	2.5	4.8	7.1	8.5	9.3	9.1	8.7

**Most Important Reasons for Drinking  
by Indiana Male 6th - 12th Graders, 2007**  
(Values expressed as percentages, multiple responses allowed)

	Grade						
	6th	7th	8th	9th	10th	11th	12th
Did not drink alcoholic beverages	79.7	71.0	61.4	53.8	44.3	41.0	35.1
To experiment	10.4	11.9	11.7	9.5	8.2	6.5	5.0
To relax or relieve tension	0.4	0.7	1.3	1.6	2.2	2.3	2.5
To feel good or get high	0.4	0.7	1.2	1.4	1.6	1.7	1.6
To seek deeper insights or understanding	0.2	0.2	0.3	0.4	0.4	0.6	0.6
To have a good time with friends	1.0	2.3	4.8	8.6	12.6	15.2	17.5
To fit in with a group I like	0.4	0.6	0.7	0.7	0.9	0.9	0.9
To get away from my problems	0.7	1.0	1.4	1.8	2.3	2.2	2.7
Because of boredom	1.1	1.7	3.1	3.7	4.8	4.9	5.4
Because of anger	0.8	1.2	1.8	2.2	2.7	2.9	2.6
To get through the day	0.1	0.3	0.4	0.6	0.7	0.6	0.7
To increase the effects of other drugs	0.1	0.2	0.3	0.8	1.2	1.1	1.4
To decrease the effects of other drugs	0.1	0.0	0.1	0.1	0.1	0.1	0.2
To get to sleep	0.3	0.6	0.7	0.7	1.2	1.3	1.6
Because it tastes good	3.8	6.4	9.7	12.7	15.1	16.7	20.1
Because I am hooked	0.6	1.2	1.2	1.5	1.9	2.0	2.2

**Consequences of ATOD Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had a hangover	No Answer	5.5	5.5	5.0	5.5	5.6	5.5	5.3
	Never	90.5	85.3	78.6	72.9	66.3	62.3	57.3
	Once	2.0	4.1	6.4	7.7	9.6	10.8	11.4
	Twice	0.9	1.8	3.4	4.4	5.9	6.7	7.2
	3-5 times	0.6	1.6	3.1	4.3	5.6	6.8	8.0
	6-10 times	0.2	0.5	1.2	1.6	2.3	2.6	3.8
	11 or more times	0.3	1.2	2.2	3.5	4.7	5.3	7.0
Had nausea or vomited	No Answer	5.9	5.9	5.5	6.1	6.1	5.9	5.7
	Never	90.6	86.9	82.5	77.7	69.9	64.5	58.9
	Once	2.0	3.8	6.0	7.4	10.3	12.0	13.5
	Twice	0.7	1.5	2.6	3.4	5.2	6.9	7.5
	3-5 times	0.4	1.0	1.8	2.7	4.6	5.7	8.2
	6-10 times	0.2	0.4	0.7	1.2	1.7	2.2	3.0
	11 or more times	0.2	0.6	0.9	1.5	2.1	2.8	3.1
Had a memory loss	No Answer	6.2	6.2	5.8	6.6	6.6	6.1	6.2
	Never	91.6	89.0	85.8	81.3	75.0	73.2	69.5
	Once	1.3	2.4	3.6	4.5	6.3	6.8	7.2
	Twice	0.4	0.9	1.7	2.2	3.5	3.7	4.5
	3-5 times	0.2	0.7	1.1	2.3	3.6	4.2	5.5
	6-10 times	0.1	0.3	0.7	1.1	1.9	2.2	2.9
	11 or more times	0.2	0.5	1.2	2.0	3.1	3.8	4.0
Poor on school test	No Answer	6.3	6.3	5.8	6.6	6.5	6.2	6.1
	Never	91.4	89.5	88.0	85.7	84.6	85.9	85.6
	Once	1.0	1.6	2.2	2.2	2.5	2.1	2.2
	Twice	0.5	0.8	1.0	1.4	1.6	1.3	1.4
	3-5 times	0.3	0.7	1.0	1.4	1.7	1.7	1.9
	6-10 times	0.1	0.3	0.6	0.7	0.9	0.7	0.9
	11 or more times	0.3	0.9	1.3	1.9	2.2	2.1	1.8
Missed school	No Answer	6.4	6.4	5.9	6.5	6.6	6.3	6.2
	Never	91.9	90.0	89.2	87.5	85.9	86.5	84.9
	Once	0.7	1.3	1.7	1.9	2.5	2.1	2.6
	Twice	0.3	0.7	1.0	1.1	1.2	1.4	1.7
	3-5 times	0.3	0.6	0.9	1.2	1.7	1.4	1.9
	6-10 times	0.1	0.4	0.5	0.8	0.8	0.9	1.1
	11 or more times	0.2	0.6	0.9	1.1	1.3	1.4	1.5
Got into a fight	No Answer	6.4	6.2	6.0	6.6	6.6	6.2	6.3
	Never	89.7	87.1	84.2	82.0	78.8	78.5	77.2
	Once	1.7	2.7	3.9	4.5	5.8	5.8	6.3
	Twice	0.7	1.2	1.8	2.4	3.1	3.5	3.8
	3-5 times	0.6	1.2	1.8	1.8	2.8	3.1	3.4
	6-10 times	0.3	0.6	0.8	0.9	1.0	1.0	1.2
	11 or more times	0.5	1.1	1.5	1.8	1.9	1.9	1.8

**Consequences of ATOD Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Damaged property	No Answer	6.5	6.4	5.9	6.8	6.7	6.4	6.3
	Never	92.1	90.7	89.0	87.3	85.6	86.6	86.2
	Once	0.7	1.1	2.0	2.0	2.7	2.6	2.9
	Twice	0.3	0.5	0.8	1.1	1.6	1.2	1.6
	3-5 times	0.2	0.5	0.9	1.3	1.5	1.6	1.4
	6-10 times	0.1	0.2	0.4	0.5	0.6	0.4	0.7
	11 or more times	0.2	0.6	0.9	1.1	1.3	1.1	0.9
Had trouble with parents	No Answer	6.6	6.3	6.0	6.7	6.6	6.4	6.4
	Never	90.0	88.0	85.8	83.1	79.7	79.1	78.6
	Once	1.8	2.5	3.6	4.5	6.7	7.0	7.4
	Twice	0.5	1.0	1.4	1.8	2.5	2.6	3.2
	3-5 times	0.3	0.8	1.0	1.5	2.0	2.3	2.3
	6-10 times	0.2	0.3	0.6	0.8	0.7	0.7	0.8
	11 or more times	0.7	1.1	1.5	1.6	1.7	1.9	1.4
Had trouble with police	No Answer	6.6	6.4	6.1	6.7	6.7	6.8	6.3
	Never	91.5	90.0	88.4	87.0	85.7	85.3	85.6
	Once	0.8	1.5	2.1	2.8	3.9	4.5	5.1
	Twice	0.4	0.7	1.1	1.3	1.4	1.5	1.3
	3-5 times	0.3	0.7	1.0	1.1	1.2	0.8	0.9
	6-10 times	0.1	0.2	0.5	0.4	0.4	0.3	0.3
	11 or more times	0.3	0.6	0.9	0.7	0.7	0.8	0.5
Driven under the influence	No Answer	6.8	6.5	6.3	7.0	7.2	6.7	6.7
	Never	92.3	91.5	90.1	88.2	83.0	77.5	72.4
	Once	0.4	0.8	1.4	1.9	3.8	5.1	5.7
	Twice	0.2	0.3	0.6	0.9	1.8	2.9	3.3
	3-5 times	0.1	0.3	0.6	0.8	1.6	2.8	4.3
	6-10 times	0.1	0.1	0.3	0.4	0.8	1.5	2.3
	11 or more times	0.2	0.4	0.8	0.8	1.9	3.5	5.3
Rode with a drunk driver	No Answer	6.8	6.2	6.0	6.5	6.7	6.5	6.4
	Never	89.6	87.8	82.9	79.2	73.0	69.9	66.8
	Once	1.3	1.9	3.0	4.0	5.3	5.9	6.0
	Twice	0.5	1.0	1.8	2.3	3.4	3.6	3.6
	3-5 times	0.7	1.0	2.1	2.9	3.9	4.4	5.5
	6-10 times	0.3	0.6	1.1	1.3	1.8	2.2	3.0
	11 or more times	0.8	1.6	2.9	3.9	5.8	7.4	8.7
Forgot things you did while high	No Answer	6.8	6.5	6.2	6.9	6.9	6.5	6.5
	Never	91.2	88.1	84.3	79.3	73.1	70.7	67.6
	Once	1.2	2.5	3.7	5.3	6.5	7.3	7.2
	Twice	0.4	1.1	1.8	2.5	3.7	4.1	5.0
	3-5 times	0.2	0.8	1.8	2.5	4.2	4.8	5.6
	6-10 times	0.1	0.3	0.7	1.2	1.9	2.1	3.0
	11 or more times	0.2	0.6	1.4	2.3	3.7	4.5	5.2

**Consequences of ATOD Use  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had been told to cut down	No Answer	6.8	6.4	6.2	6.8	6.7	6.3	6.4
	Never	91.9	90.9	89.1	86.5	83.7	83.4	82.3
	Once	0.6	1.0	1.6	2.3	3.0	3.0	3.4
	Twice	0.3	0.5	0.7	1.1	1.8	1.9	2.4
	3-5 times	0.1	0.4	0.9	1.0	1.8	2.0	2.2
	6-10 times	0.1	0.1	0.4	0.6	0.8	0.9	1.1
	11 or more times	0.3	0.6	1.1	1.6	2.2	2.5	2.2
Used alcohol or drugs to relax or fit in	No Answer	6.6	6.4	6.0	6.8	6.8	6.4	6.5
	Never	90.7	88.3	84.8	80.7	77.2	75.2	72.8
	Once	1.3	2.1	3.2	4.0	4.8	5.1	4.6
	Twice	0.6	0.9	1.6	2.0	2.5	2.6	3.4
	3-5 times	0.4	1.0	1.6	2.4	3.3	3.7	4.0
	6-10 times	0.2	0.3	0.8	1.1	1.5	1.9	2.5
	11 or more times	0.3	0.9	2.0	2.9	4.0	5.1	6.1
Used alcohol or drugs alone	No Answer	6.6	6.5	6.0	6.8	6.7	6.5	6.5
	Never	89.6	86.2	81.2	77.2	72.5	72.2	70.3
	Once	1.7	2.4	4.1	4.7	5.9	5.6	5.3
	Twice	0.8	1.6	2.5	2.9	3.6	3.4	4.2
	3-5 times	0.6	1.4	2.6	3.3	4.2	4.2	5.2
	6-10 times	0.3	0.6	1.3	1.8	2.0	2.3	2.4
	11 or more times	0.4	1.2	2.3	3.3	5.0	5.9	6.1

**Perceived Risk of Selected Substance Use Behavior  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	4.6	5.2	5.3	6.5	6.6	6.7	6.7
	No Risk	10.1	9.3	9.1	8.5	7.7	6.9	6.7
	Slight Risk	13.9	12.5	12.8	11.2	10.1	9.5	8.5
	Moderate Risk	28.7	29.3	28.7	27.7	27.2	27.4	25.8
	Great Risk	42.6	43.6	44.0	46.0	48.4	49.5	52.4
Smoke marijuana occasionally	No Answer	4.9	5.5	5.6	6.6	6.7	6.7	6.7
	No Risk	10.2	11.0	14.0	16.6	19.0	20.2	22.5
	Slight Risk	10.8	12.7	17.0	20.4	24.7	26.5	28.6
	Moderate Risk	29.5	30.3	28.2	25.6	23.9	22.1	21.2
	Great Risk	44.5	40.5	35.3	30.8	25.7	24.5	20.9
Smoke marijuana regularly	No Answer	5.1	5.7	5.7	6.8	6.9	6.8	6.9
	No Risk	9.9	9.8	11.0	11.4	12.1	12.0	12.0
	Slight Risk	4.0	4.3	7.3	10.4	12.6	14.2	16.1
	Moderate Risk	14.2	14.7	17.6	19.3	22.3	23.2	25.9
	Great Risk	66.7	65.4	58.5	52.1	46.1	43.7	39.1
Use cocaine occasionally	No Answer	5.3	5.8	5.7	6.9	7.0	6.8	6.9
	No Risk	9.6	8.7	8.5	7.4	7.1	6.3	6.5
	Slight Risk	8.2	7.5	6.7	6.7	6.7	6.6	6.8
	Moderate Risk	29.2	29.5	27.8	26.4	24.4	23.7	22.3
	Great Risk	47.8	48.5	51.4	52.6	54.8	56.6	57.5
Use cocaine regularly	No Answer	6.1	6.7	6.7	7.8	7.9	7.7	7.8
	No Risk	9.7	8.5	8.3	7.0	6.7	5.8	5.7
	Slight Risk	3.2	2.7	2.3	2.4	2.0	2.1	2.0
	Moderate Risk	11.6	9.4	9.2	9.2	8.5	8.0	7.4
	Great Risk	69.4	72.7	73.5	73.6	74.9	76.4	77.0
Occasionally consume 1-2 alcoholic drinks	No Answer	5.7	6.1	6.2	7.1	7.3	7.1	7.1
	No Risk	22.4	24.0	28.5	31.3	34.1	37.8	41.4
	Slight Risk	35.6	36.2	35.1	32.8	33.1	32.6	31.1
	Moderate Risk	17.8	16.5	13.8	13.4	11.6	10.1	9.0
	Great Risk	18.6	17.3	16.3	15.4	13.9	12.4	11.3
Binge drink weekly	No Answer	5.1	5.7	6.0	7.0	7.1	7.0	7.1
	No Risk	11.1	10.6	11.6	11.6	11.4	11.3	12.1
	Slight Risk	11.4	12.6	14.9	16.9	18.5	20.7	22.2
	Moderate Risk	28.8	28.5	29.2	28.7	29.0	29.1	29.6
	Great Risk	43.6	42.5	38.3	35.8	33.9	31.9	29.0

**Perceived Peer Approval of Using Selected Substances  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade					
		6th	7th	8th	9th	10th	11th
Smoke 1+ pack cigarettes per day	No Answer	7.6	7.5	7.2	8.1	8.2	8.3
	Strongly Approve	4.3	4.3	4.5	4.4	4.8	4.1
	Approve	1.5	2.3	4.0	5.6	7.0	8.5
	Do Not Know	14.8	16.4	19.6	20.4	21.0	21.8
	Disapprove	14.7	15.6	16.6	18.2	18.9	20.0
Smoke marijuana occasionally	Strongly Disapprove	57.0	53.8	48.2	43.3	40.1	38.2
	No Answer	7.8	7.7	7.2	8.2	8.3	8.2
	Strongly Approve	4.3	4.7	5.7	6.1	7.6	6.9
	Approve	1.5	3.4	6.3	9.1	12.0	14.0
	Do Not Know	12.7	14.0	17.6	19.2	20.2	20.0
Smoke marijuana regularly	Disapprove	10.7	11.5	12.8	13.9	14.4	15.4
	Strongly Disapprove	63.0	58.8	50.3	43.5	37.4	35.6
	No Answer	8.0	7.9	7.3	8.2	8.5	8.4
	Strongly Approve	4.3	4.7	5.9	6.4	7.6	6.8
	Approve	1.0	2.1	4.0	6.1	7.8	9.0
Use cocaine occasionally	Do Not Know	11.9	13.4	16.8	18.1	18.9	18.6
	Disapprove	8.1	8.8	10.2	12.2	12.6	13.5
	Strongly Disapprove	66.7	63.1	55.8	49.1	44.6	43.6
	No Answer	8.1	7.8	7.5	8.2	8.6	8.4
	Strongly Approve	4.1	3.8	4.0	3.6	3.6	3.3
Use cocaine regularly	Approve	0.8	1.1	1.4	1.8	2.3	2.4
	Do Not Know	11.7	13.0	15.1	15.8	15.5	15.0
	Disapprove	10.5	10.4	11.0	12.2	11.8	12.0
	Strongly Disapprove	64.8	63.8	61.1	58.4	58.2	58.9
	No Answer	8.3	8.1	7.9	8.7	8.8	8.8
Occasionally consume 1-2 alcoholic drinks	Strongly Approve	4.2	3.8	4.0	3.7	3.7	3.1
	Approve	0.6	1.0	1.2	1.4	1.7	1.5
	Do Not Know	11.5	12.5	14.7	15.6	14.9	14.5
	Disapprove	7.4	7.5	8.3	9.5	9.0	8.9
	Strongly Disapprove	68.0	67.0	63.9	61.1	61.8	63.2
Binge drink weekly	No Answer	8.2	7.9	7.5	8.2	8.3	8.3
	Strongly Approve	4.6	5.5	7.0	8.7	11.2	12.1
	Approve	5.2	8.8	13.4	18.4	24.2	27.0
	Do Not Know	18.5	20.5	24.2	24.6	24.0	23.1
	Disapprove	14.3	13.7	12.9	12.2	10.8	9.7
	Strongly Disapprove	49.1	43.6	35.1	27.9	21.4	19.8
	No Answer	8.4	8.2	7.6	8.4	8.6	8.7
	Strongly Approve	4.6	4.9	6.2	7.8	10.2	10.2
	Approve	1.9	4.0	7.1	10.6	15.1	16.9
	Do Not Know	13.7	16.4	20.6	22.6	22.7	23.6
	Disapprove	11.0	11.7	13.1	13.3	12.8	12.5
	Strongly Disapprove	60.4	54.8	45.4	37.2	30.5	28.1
							24.5

**Perceived Parental Approval of Using Selected Substances  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	7.6	7.6	7.3	8.1	8.3	8.3	8.3
	Strongly Approve	4.1	3.6	3.6	3.1	3.0	2.4	2.7
	Approve	0.5	0.7	0.9	1.1	1.8	2.5	2.4
	Do Not Know	4.8	5.0	5.5	6.4	6.9	6.8	7.1
	Disapprove	3.5	4.1	4.6	5.8	8.2	9.1	12.3
Smoke marijuana occasionally	Strongly Disapprove	79.5	79.1	78.2	75.5	71.7	70.9	67.1
	No Answer	7.8	7.8	7.4	8.4	8.6	8.3	8.4
	Strongly Approve	4.0	3.5	3.6	3.0	3.1	2.5	2.7
	Approve	0.3	0.6	1.0	1.1	2.1	2.2	2.6
	Do Not Know	4.5	4.9	5.6	6.5	7.1	7.1	7.2
Smoke marijuana regularly	Disapprove	2.6	2.9	3.8	4.9	6.8	7.4	9.5
	Strongly Disapprove	80.8	80.4	78.7	76.1	72.3	72.5	69.6
	No Answer	7.9	7.9	7.3	8.4	8.6	8.4	8.5
	Strongly Approve	4.1	3.5	3.6	3.1	3.1	2.5	2.7
	Approve	0.2	0.4	0.7	0.8	1.3	1.3	1.6
Use cocaine occasionally	Do Not Know	4.4	4.7	5.6	6.1	6.9	6.5	6.8
	Disapprove	2.3	2.5	3.3	4.3	5.7	6.1	7.1
	Strongly Disapprove	81.1	81.0	79.4	77.3	74.4	75.1	73.4
	No Answer	8.0	7.9	7.5	8.5	8.8	8.6	8.5
	Strongly Approve	3.9	3.4	3.4	2.9	2.8	2.3	2.4
Use cocaine regularly	Approve	0.1	0.2	0.3	0.3	0.3	0.4	0.5
	Do Not Know	4.3	4.6	4.9	5.3	5.7	5.5	5.4
	Disapprove	2.4	2.4	2.7	3.0	4.1	3.9	4.6
	Strongly Disapprove	81.2	81.4	81.2	80.0	78.3	79.3	78.7
	No Answer	8.1	8.0	7.6	8.6	8.8	8.5	8.6
Occasionally consume 1-2 alcoholic drinks	Strongly Approve	3.9	3.4	3.4	2.9	2.9	2.3	2.4
	Approve	0.1	0.2	0.3	0.3	0.3	0.5	0.3
	Do Not Know	4.3	4.6	4.8	5.3	5.7	5.5	5.4
	Disapprove	2.1	2.1	2.5	2.8	3.9	3.6	4.1
	Strongly Disapprove	81.4	81.7	81.3	80.1	78.4	79.7	79.1
Binge drink weekly	No Answer	7.9	8.0	7.5	8.3	8.6	8.3	8.4
	Strongly Approve	4.1	3.6	3.7	3.3	3.3	3.0	3.2
	Approve	3.2	3.4	4.4	4.9	7.3	8.5	12.2
	Do Not Know	7.4	8.5	9.3	10.6	12.1	12.0	13.6
	Disapprove	6.2	6.9	8.1	9.9	11.9	13.1	15.2
	Strongly Disapprove	71.3	69.7	66.9	63.0	56.9	55.0	47.4
	No Answer	8.2	8.3	7.8	8.8	8.9	8.9	8.6
	Strongly Approve	4.0	3.5	3.6	3.1	3.1	2.5	2.8
	Approve	0.7	0.8	1.3	1.4	2.7	2.9	4.1
	Do Not Know	5.3	5.8	6.9	8.1	9.3	9.4	10.9
	Disapprove	3.5	4.5	5.2	6.7	8.8	10.3	13.2
	Strongly Disapprove	78.3	77.1	75.2	71.9	67.3	65.9	60.3

**Days being Absent from School in the Past Year  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

	<b>Grade</b>						
	<b>6th</b>	<b>7th</b>	<b>8th</b>	<b>9th</b>	<b>10th</b>	<b>11th</b>	<b>12th</b>
No Answer	2.0	1.4	1.2	0.9	0.6	0.5	0.5
Never	12.9	13.5	12.0	15.0	13.7	12.5	10.5
Once	10.7	11.4	11.2	11.5	11.3	11.0	9.1
Twice	17.2	17.6	16.3	16.0	16.2	15.5	14.2
3-5 days	33.6	32.6	32.6	32.2	31.9	32.8	32.6
6-10 days	15.9	15.5	17.3	16.1	16.5	17.2	19.8
11-19 days	5.1	5.3	6.2	5.0	5.8	6.6	8.4
20 or more days	2.4	2.7	3.4	3.4	3.9	3.9	4.9

**Participation in Some Form of Camp  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

	<b>Grade</b>						
	<b>6th</b>	<b>7th</b>	<b>8th</b>	<b>9th</b>	<b>10th</b>	<b>11th</b>	<b>12th</b>
No participation in camps	75.1	79.7	81.7	84.6	86.7	85.3	87.2
Afternoons R.O.C.K. in Indiana	7.7	5.7	4.8	2.2	1.3	0.9	0.8
Youth Leadership Programs	7.1	8.2	8.4	8.5	7.5	8.3	7.1
S.A.D.D., S.T.A.N.D. or other prevention programs	8.7	5.3	4.3	3.8	3.5	4.1	3.8
Participated in 2+ types of camps	1.4	1.1	0.8	0.9	1.0	1.4	1.0

**Participation in Organized Family Events  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

	<b>Grade</b>						
	<b>6th</b>	<b>7th</b>	<b>8th</b>	<b>9th</b>	<b>10th</b>	<b>11th</b>	<b>12th</b>
No Answer	4.9	4.1	3.8	3.4	3.3	2.8	3.0
Never	16.8	17.7	20.2	21.4	21.8	22.8	24.6
Once per Week	38.8	41.0	41.5	42.7	45.7	46.2	46.6
Twice per Week	18.9	18.9	18.6	17.8	16.5	16.4	14.6
3+ Times per Week	20.7	18.3	15.9	14.8	12.7	11.8	11.2

**Number of Days per School Year Spent in Various Afterschool Activities  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
School sports team	No Answer	7.2	7.5	7.2	9.0	9.5	9.2	9.5
	None	36.1	32.1	34.2	35.9	40.2	42.9	46.6
	1-9 Days	13.1	9.9	8.5	6.0	4.7	4.3	3.5
	10-29 Days	8.4	6.2	5.4	3.1	2.5	2.2	2.1
	30-59 Days	11.2	11.7	10.7	8.6	6.6	6.7	6.2
	60-89 Days	9.1	11.5	11.3	10.9	10.1	10.1	10.5
	90-129 Days	7.0	11.4	11.1	12.5	11.8	11.2	10.7
	130-180 Days	8.0	9.8	11.6	14.1	14.6	13.4	10.9
School arts programs	No Answer	8.7	9.1	8.6	10.3	10.6	10.1	10.7
	None	65.2	64.6	64.8	65.6	66.1	65.9	65.4
	1-9 Days	13.6	13.3	12.9	9.2	8.4	7.8	7.2
	10-29 Days	4.6	4.7	4.9	4.0	3.6	4.0	4.1
	30-59 Days	2.4	2.9	2.9	3.3	3.4	3.6	3.8
	60-89 Days	1.6	1.6	2.0	2.7	2.8	3.2	2.9
	90-129 Days	1.1	1.1	1.4	2.0	2.3	2.4	2.7
	130-180 Days	2.7	2.7	2.6	3.0	3.0	3.1	3.0
School club/intramurals	No Answer	9.5	10.3	9.8	11.1	11.4	10.7	11.3
	None	61.0	61.3	63.0	60.9	58.2	54.3	52.3
	1-9 Days	13.4	12.6	12.0	11.7	11.3	11.9	11.6
	10-29 Days	7.2	7.0	6.5	7.0	8.4	11.2	11.5
	30-59 Days	4.0	3.9	3.8	4.3	5.1	5.8	6.8
	60-89 Days	1.9	2.0	2.1	2.2	2.3	2.9	2.9
	90-129 Days	1.3	1.4	1.3	1.3	1.5	1.6	1.6
	130-180 Days	1.8	1.4	1.6	1.5	1.9	1.6	1.9
Youth centers (YMCA, etc)	No Answer	10.0	9.8	9.5	10.9	11.2	10.5	10.8
	None	68.8	68.7	69.6	71.4	71.7	71.9	73.3
	1-9 Days	9.8	9.5	9.2	8.0	7.6	7.6	6.7
	10-29 Days	4.0	4.3	4.0	3.9	3.6	3.9	3.5
	30-59 Days	2.5	3.1	2.8	2.5	2.7	2.7	2.6
	60-89 Days	1.7	1.9	1.8	1.2	1.4	1.5	1.4
	90-129 Days	1.2	1.2	1.3	0.9	0.8	0.9	0.6
	130-180 Days	1.9	1.5	1.8	1.2	1.0	0.9	1.1
Extended day programs	No Answer	11.3	10.6	10.2	11.7	11.8	11.4	11.4
	None	66.1	68.3	69.6	72.2	74.4	76.0	78.0
	1-9 Days	13.2	13.1	12.3	9.7	8.0	7.0	6.1
	10-29 Days	4.2	3.8	3.6	2.7	2.4	2.6	1.8
	30-59 Days	2.0	1.6	1.7	1.6	1.3	1.3	0.9
	60-89 Days	1.2	1.1	1.0	0.8	0.7	0.5	0.5
	90-129 Days	0.8	0.7	0.6	0.6	0.4	0.4	0.4
	130-180 Days	1.3	0.9	1.0	0.8	0.8	0.7	0.7

**Number of Days per School Year Spent in Various Afterschool Activities  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Supervised youth activities (4H, scouts)	No Answer	11.5	11.2	10.8	12.3	12.6	11.6	11.8
	None	55.0	59.6	63.0	66.1	67.6	68.7	70.7
	1-9 Days	9.3	7.8	7.2	6.8	6.2	6.6	5.5
	10-29 Days	6.2	5.9	5.2	4.7	4.5	4.9	4.4
	30-59 Days	6.6	6.1	5.4	4.5	4.1	3.7	3.4
	60-89 Days	4.4	4.0	3.6	2.6	2.3	2.1	1.8
	90-129 Days	3.0	2.5	2.2	1.5	1.2	1.0	1.0
	130-180 Days	3.9	2.9	2.7	1.4	1.5	1.4	1.4
Afterschool classes (art, sport, etc)	No Answer	11.1	10.9	10.4	11.5	12.1	11.3	11.5
	None	47.6	50.2	53.4	56.8	58.4	60.0	63.5
	1-9 Days	13.7	12.3	11.8	9.5	8.6	8.3	7.6
	10-29 Days	8.2	8.5	7.0	6.2	5.8	5.9	5.3
	30-59 Days	6.7	6.5	5.9	5.7	4.8	5.1	4.0
	60-89 Days	4.2	4.2	3.8	3.3	3.3	3.0	2.2
	90-129 Days	3.2	2.9	2.7	2.6	2.4	2.2	1.8
	130-180 Days	5.4	4.6	4.9	4.4	4.6	4.2	4.1
Socialization without adult supervision	No Answer	9.5	9.2	8.7	9.9	10.2	9.7	9.9
	None	39.0	27.6	20.9	16.1	11.8	9.8	9.0
	1-9 Days	18.5	17.5	15.0	12.8	10.2	7.8	7.0
	10-29 Days	10.0	13.1	13.7	13.9	12.5	10.9	10.2
	30-59 Days	6.8	9.9	11.8	13.6	15.0	14.8	15.2
	60-89 Days	5.2	6.9	9.8	11.6	13.3	15.3	14.8
	90-129 Days	4.4	6.5	8.6	10.3	12.4	14.7	15.9
	130-180 Days	6.5	9.3	11.5	11.7	14.6	17.0	17.9
At home without adult supervision	No Answer	10.6	10.5	9.5	10.8	11.0	10.5	11.1
	None	32.8	23.6	19.0	15.3	12.6	11.8	10.4
	1-9 Days	25.1	22.3	18.6	15.8	13.1	12.4	11.0
	10-29 Days	11.4	14.3	15.5	15.8	15.5	15.5	14.9
	30-59 Days	7.0	10.2	12.7	14.5	15.2	16.2	15.5
	60-89 Days	4.4	6.9	9.1	10.7	12.2	12.3	13.9
	90-129 Days	3.4	5.2	6.8	7.7	9.7	10.1	10.6
	130-180 Days	5.3	7.0	8.8	9.3	10.7	11.2	12.6
At home with adult supervision	No Answer	10.5	10.6	10.0	11.3	11.3	10.7	11.1
	None	15.6	14.3	13.7	12.3	11.1	11.5	12.3
	1-9 Days	12.0	10.6	10.6	9.3	9.3	8.5	9.1
	10-29 Days	8.6	9.4	9.5	9.5	10.6	11.0	11.7
	30-59 Days	8.3	9.6	10.9	11.9	13.1	14.0	13.9
	60-89 Days	8.0	9.6	11.3	13.2	13.7	14.4	15.1
	90-129 Days	12.9	14.8	15.9	16.2	15.9	15.5	14.3
	130-180 Days	24.2	21.1	18.0	16.2	15.0	14.4	12.3

**Number of Days per School Year Spent in Various Afterschool Activities  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Work for pay outside home	No Answer	10.4	9.9	9.4	10.9	10.9	10.1	10.5
	None	49.4	43.9	41.7	41.3	36.0	28.7	23.5
	1-9 Days	16.8	16.7	15.5	11.7	8.4	6.3	4.5
	10-29 Days	9.2	11.6	11.6	10.8	9.1	8.1	6.8
	30-59 Days	5.5	7.1	8.8	9.9	10.5	10.2	10.6
	60-89 Days	2.8	4.3	5.4	6.6	9.2	12.5	12.6
	90-129 Days	2.1	2.7	3.3	4.1	7.6	11.0	13.8
	130-180 Days	3.8	3.8	4.3	4.7	8.3	13.2	17.8
Work for pay inside a home	No Answer	10.5	10.1	9.7	11.5	11.6	10.9	11.8
	None	46.0	44.9	46.6	49.7	53.7	57.4	61.0
	1-9 Days	18.1	17.4	16.0	14.5	12.9	12.7	10.6
	10-29 Days	9.9	10.4	10.0	9.1	8.2	7.3	6.7
	30-59 Days	5.7	7.0	6.6	6.2	5.4	4.9	3.8
	60-89 Days	3.1	3.2	3.9	3.2	2.8	2.5	1.8
	90-129 Days	2.0	2.1	2.2	1.9	1.9	1.4	1.5
	130-180 Days	4.8	4.8	5.1	3.9	3.7	2.9	2.8
Volunteer outside the home	No Answer	10.7	10.3	9.8	11.5	11.5	10.6	11.3
	None	53.7	51.7	52.0	51.5	51.2	49.9	48.0
	1-9 Days	17.8	18.1	17.4	16.1	15.4	15.8	15.4
	10-29 Days	7.8	8.7	9.0	9.2	10.1	11.2	12.2
	30-59 Days	4.1	4.8	5.1	5.2	5.3	6.1	6.3
	60-89 Days	2.0	2.5	2.8	2.8	2.7	2.9	3.0
	90-129 Days	1.3	1.5	1.7	1.6	1.4	1.4	1.3
	130-180 Days	2.5	2.4	2.3	2.1	2.4	2.1	2.4
Volunteer inside a home	No Answer	10.3	10.1	9.5	11.2	11.6	10.7	11.5
	None	48.0	46.6	48.8	48.7	49.5	49.6	50.2
	1-9 Days	18.2	17.7	16.5	14.7	13.2	12.2	11.7
	10-29 Days	8.7	9.2	8.4	8.3	8.6	9.4	8.9
	30-59 Days	5.1	5.7	5.6	6.0	5.7	6.7	6.3
	60-89 Days	2.9	3.3	3.5	3.6	3.4	3.9	3.7
	90-129 Days	2.2	2.5	2.6	2.5	2.5	2.4	2.4
	130-180 Days	4.7	4.9	5.1	5.1	5.4	5.1	5.3

Note: The number of school days in a year is estimated as 180 days.

**Perceived Safety in Neighborhood and in School  
by Indiana Male 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Out alone during the day	No Answer	5.9	6.6	6.2	7.1	7.4	7.0	7.0
	Very Safe	53.3	60.9	65.7	71.3	74.5	78.7	81.2
	Somewhat Safe	31.1	25.8	22.9	17.6	14.4	11.4	9.0
	Somewhat Unsafe	6.6	4.3	3.3	2.2	1.9	1.2	1.2
	Very Unsafe	3.2	2.5	2.0	1.8	1.9	1.7	1.6
Out alone after dark	No Answer	6.2	6.8	6.3	7.3	7.6	7.1	7.0
	Very Safe	22.4	28.6	36.4	45.8	51.9	58.6	63.8
	Somewhat Safe	31.3	34.6	34.5	31.9	28.7	25.8	21.8
	Somewhat Unsafe	21.4	18.5	15.0	10.2	7.9	5.4	4.5
	Very Unsafe	18.7	11.5	7.8	4.8	3.8	3.0	2.9
Inside home alone during the day	No Answer	6.4	6.9	6.7	7.4	7.8	7.3	7.2
	Very Safe	69.4	75.0	79.1	81.5	82.2	84.0	85.5
	Somewhat Safe	19.0	14.2	11.3	8.7	7.6	6.4	5.2
	Somewhat Unsafe	3.2	2.1	1.4	1.0	0.9	0.7	0.7
	Very Unsafe	2.1	1.8	1.5	1.5	1.6	1.6	1.4
Inside home alone after dark	No Answer	6.4	6.8	6.7	7.3	7.8	7.4	7.2
	Very Safe	46.8	55.3	62.9	69.4	72.6	76.0	79.1
	Somewhat Safe	26.3	24.3	21.7	17.2	15.0	12.6	10.1
	Somewhat Unsafe	12.7	9.0	5.7	3.8	2.6	2.0	1.8
	Very Unsafe	7.8	4.7	3.0	2.3	2.0	2.0	1.7
Traveling to/from school	No Answer	6.3	7.0	6.7	7.4	7.9	7.5	7.4
	Very Safe	65.0	66.7	68.2	71.1	72.6	75.7	77.9
	Somewhat Safe	21.4	19.5	19.3	17.0	15.4	13.2	11.5
	Somewhat Unsafe	4.3	3.9	3.3	2.3	1.9	1.6	1.4
	Very Unsafe	2.9	2.8	2.6	2.1	2.1	2.0	1.9
Staying in class	No Answer	5.9	6.8	6.5	7.3	7.7	7.3	7.3
	Very Safe	80.9	76.7	73.8	71.9	69.5	72.4	73.9
	Somewhat Safe	9.9	12.1	14.4	15.8	16.9	14.7	13.1
	Somewhat Unsafe	1.4	1.8	2.3	1.9	2.1	2.1	2.5
	Very Unsafe	1.9	2.6	3.0	3.1	3.7	3.5	3.2
Alone in school hallways	No Answer	6.1	7.0	6.7	7.4	7.8	7.4	7.3
	Very Safe	71.1	68.4	69.5	66.9	67.0	71.0	72.8
	Somewhat Safe	17.4	17.1	16.8	18.5	17.8	15.1	13.7
	Somewhat Unsafe	3.2	4.3	3.6	3.6	3.5	2.7	2.6
	Very Unsafe	2.2	3.2	3.4	3.7	3.9	3.9	3.5
Staying for afterschool activities	No Answer	7.3	7.7	7.1	7.9	8.3	7.7	7.5
	Very Safe	70.0	70.4	70.2	69.7	68.7	71.7	73.6
	Somewhat Safe	17.2	15.7	16.3	16.3	16.5	14.4	13.0
	Somewhat Unsafe	3.1	3.0	2.9	2.7	2.6	2.4	2.3
	Very Unsafe	2.4	3.1	3.5	3.4	4.0	3.8	3.5

**Annual Prevalence of Weapon Use and Violent Behavior  
by Indiana Male 6th - 12th Graders, 2007**  
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Physical fight	No Answer	6.3	6.8	6.5	7.4	8.0	7.5	7.5
	Never	46.2	45.8	45.9	49.9	53.7	57.4	62.0
	1-2 times	30.1	29.8	30.5	28.0	25.2	24.1	21.6
	3-9 times	11.5	11.6	11.2	9.7	8.4	7.3	5.9
	10+ times	5.9	6.0	6.0	4.9	4.6	3.7	2.9
Shouting argument	No Answer	6.6	7.3	6.8	7.7	8.4	7.5	7.6
	Never	41.5	37.0	32.9	33.3	31.8	34.3	33.6
	1-2 times	26.4	25.5	26.4	26.0	25.2	25.0	27.0
	3-9 times	14.0	16.1	17.5	17.8	19.5	18.9	19.0
	10+ times	11.5	14.1	16.3	15.2	15.1	14.3	12.8
Carried gun for hunting	No Answer	7.0	7.4	7.0	7.9	8.5	7.6	7.7
	Never	59.2	55.9	55.1	55.0	55.8	57.7	59.4
	1-2 times	13.0	12.5	12.3	11.6	10.2	10.2	9.3
	3-9 times	8.8	10.0	10.0	9.8	9.8	9.2	9.1
	10+ times	12.0	14.2	15.6	15.7	15.8	15.3	14.5
Carried gun for other reasons	No Answer	7.1	7.4	7.2	8.1	8.5	7.8	7.8
	Never	85.8	83.1	80.5	79.7	78.3	80.4	80.4
	1-2 times	3.6	4.6	5.2	5.9	5.8	5.3	5.0
	3-9 times	1.6	1.9	2.8	2.5	2.7	2.5	2.7
	10+ times	1.9	3.0	4.2	3.9	4.7	4.1	4.1
Carried weapon for hunting	No Answer	7.2	7.5	7.2	8.0	8.7	8.0	7.9
	Never	72.6	68.8	67.7	67.6	66.9	69.3	70.7
	1-2 times	9.2	9.6	9.3	8.5	7.4	7.1	6.0
	3-9 times	5.0	5.7	6.2	5.7	5.9	5.2	5.2
	10+ times	5.9	8.3	9.7	10.2	11.1	10.4	10.2
Carried weapon for other reasons	No Answer	7.1	7.6	7.4	8.2	8.8	8.0	8.0
	Never	81.3	77.4	73.3	73.1	69.9	72.7	71.6
	1-2 times	5.8	6.6	7.7	7.4	7.9	6.8	6.3
	3-9 times	2.4	3.5	4.4	4.1	4.2	3.8	4.4
	10+ times	3.4	4.9	7.2	7.2	9.2	8.8	9.8
Carried gun to school	No Answer	6.9	7.3	7.1	7.9	8.5	7.8	7.8
	Never	92.1	90.9	90.4	89.5	88.4	88.9	89.1
	1-2 times	0.6	0.8	1.1	1.2	1.1	1.3	1.3
	3-9 times	0.1	0.3	0.3	0.3	0.5	0.6	0.6
	10+ times	0.4	0.7	1.1	1.1	1.5	1.4	1.3
Carried weapon to school	No Answer	6.8	7.4	7.0	8.1	8.5	8.0	7.8
	Never	89.5	87.1	84.6	83.6	81.1	82.7	82.4
	1-2 times	2.8	3.6	5.4	4.6	5.2	4.3	4.4
	3-9 times	0.5	0.8	1.2	1.4	1.9	1.6	1.6
	10+ times	0.4	1.1	1.8	2.3	3.3	3.4	3.8

**Annual Prevalence of Weapon Use and Violent Behavior  
by Indiana Male 6th - 12th Graders, 2007**  
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Threatened by weapon on school property	No Answer	7.0	7.6	7.2	8.2	8.6	7.8	8.0
	Never	85.6	83.4	82.3	81.3	81.1	83.6	85.1
	1-2 times	5.5	6.3	7.1	7.3	6.6	5.5	4.4
	3-9 times	1.0	1.5	1.8	1.6	1.9	1.6	1.2
	10+ times	0.8	1.1	1.7	1.5	1.7	1.6	1.3
Property stolen while at school	No Answer	7.1	7.7	7.3	8.3	8.8	7.9	7.9
	Never	66.9	60.6	59.2	57.1	58.3	61.1	64.4
	1-2 times	17.7	21.6	22.4	24.3	23.2	22.7	20.8
	3-9 times	5.6	7.2	7.4	7.6	6.9	5.9	5.0
	10+ times	2.6	3.0	3.6	2.7	2.9	2.4	1.8
Missed school: Felt unsafe on school property	No Answer	7.0	7.6	7.0	8.0	8.7	7.9	7.9
	Never	89.8	88.3	87.8	87.7	85.4	88.1	86.7
	1-2 times	2.3	2.7	3.2	2.4	3.5	2.0	3.6
	3-9 times	0.4	0.6	0.8	0.8	1.0	0.7	0.7
	10+ times	0.6	0.8	1.1	1.1	1.4	1.3	1.1
Missed school: Felt unsafe on way to/from school	No Answer	6.9	7.6	7.1	8.0	8.9	8.2	8.0
	Never	91.0	90.0	89.8	89.0	87.7	88.8	88.9
	1-2 times	1.4	1.4	1.7	1.6	1.6	1.3	1.5
	3-9 times	0.2	0.4	0.4	0.4	0.6	0.5	0.6
	10+ times	0.5	0.7	1.0	1.1	1.3	1.2	1.0

**Gambling during the Last Twelve Months  
by Indiana Male 6th - 12th Graders, 2007**  
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Played cards	No Answer	6.2	6.7	6.5	8.1	8.7	8.2	8.5
	Not at all	72.2	64.9	58.5	54.1	49.2	48.7	46.0
	Less than once a month	12.0	15.2	18.4	19.5	21.0	21.7	23.0
	About once a month	5.7	7.9	9.7	11.2	12.8	12.6	14.4
	About once a week	2.7	3.7	4.6	5.2	6.0	6.3	6.0
	Daily	1.2	1.5	2.2	1.9	2.2	2.5	2.2
Bet on games	No Answer	6.5	6.9	6.8	8.4	8.9	8.3	8.7
	Not at all	69.4	62.1	56.7	53.7	50.8	51.7	51.5
	Less than once a month	13.1	15.8	17.4	18.1	18.9	18.4	19.3
	About once a month	5.9	8.8	10.5	11.5	12.1	12.6	12.1
	About once a week	3.3	4.4	5.6	5.6	6.6	6.2	5.9
	Daily	1.7	2.1	3.0	2.6	2.6	2.9	2.5
Bet on sports	No Answer	6.9	7.3	7.2	8.9	9.3	8.7	9.1
	Not at all	67.7	61.4	56.3	54.2	53.3	54.7	54.6
	Less than once a month	14.6	16.9	19.1	18.6	18.5	18.8	19.7
	About once a month	5.7	7.9	9.5	10.7	10.8	10.0	9.7
	About once a week	3.0	4.4	5.0	5.2	5.5	5.3	4.6
	Daily	2.1	2.2	3.0	2.4	2.6	2.6	2.3
Bought lottery tickets	No Answer	7.0	7.3	7.3	8.7	9.3	8.8	8.9
	Not at all	78.8	76.7	75.1	74.6	74.6	73.4	56.2
	Less than once a month	8.0	9.2	9.5	9.1	8.5	9.2	17.9
	About once a month	3.5	3.8	4.2	4.0	3.8	4.3	9.6
	About once a week	1.7	2.1	2.3	2.3	2.3	2.8	5.0
	Daily	0.9	1.0	1.5	1.3	1.4	1.5	2.4
Gambled in a casino	No Answer	7.0	7.3	7.3	8.8	9.3	8.8	9.0
	Not at all	91.6	90.5	90.0	88.1	87.1	87.2	86.0
	Less than once a month	0.7	0.9	1.1	1.4	1.5	1.7	2.7
	About once a month	0.2	0.4	0.5	0.6	0.6	0.7	0.8
	About once a week	0.1	0.2	0.3	0.3	0.5	0.5	0.5
	Daily	0.4	0.6	0.8	0.8	1.0	1.0	1.1
Played online	No Answer	7.0	7.5	7.3	9.0	9.3	8.9	9.3
	Not at all	88.4	86.1	85.9	84.5	83.7	84.5	84.2
	Less than once a month	2.3	2.9	2.8	2.8	2.9	2.2	2.5
	About once a month	0.7	1.4	1.5	1.4	1.5	1.5	1.4
	About once a week	0.6	0.8	1.0	1.0	1.0	1.2	1.1
	Daily	1.0	1.1	1.5	1.3	1.5	1.7	1.6

**Gambling during the Last Twelve Months  
by Indiana Male 6th - 12th Graders, 2007**  
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Felt bad about the amount bet	No Answer	7.6	8.1	7.4	9.2	9.6	9.0	9.3
	Yes	7.7	7.1	7.2	6.1	6.4	5.8	6.1
	No	25.9	32.7	39.3	41.5	43.6	44.8	48.0
	I do not bet for money	58.9	52.1	46.1	43.2	40.4	40.4	36.6
Would like to stop betting but could not	No Answer	7.6	8.3	7.6	9.4	9.4	9.1	9.3
	Yes	6.8	5.3	4.6	3.8	3.1	2.8	2.3
	No	26.5	33.7	40.9	43.0	46.4	47.1	51.2
	I do not bet for money	59.1	52.7	46.9	43.8	41.1	41.0	37.3

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	2.8	2.8	2.4	2.4	2.3	2.2	2.3
	Never	87.7	80.7	70.1	65.6	58.7	55.0	52.5
	7 years or younger	1.6	1.6	2.0	1.6	1.6	1.2	0.9
	8 years	0.8	1.1	1.6	1.3	1.3	1.1	0.7
	9 years	1.1	1.1	1.4	1.5	1.4	1.2	0.9
	10 years	2.0	2.2	2.4	2.2	2.2	1.4	1.1
	11 years	2.3	3.0	3.7	2.8	2.6	2.1	1.7
	12 years	1.4	4.6	5.6	4.7	4.5	3.9	3.1
	13 years	0.2	2.5	7.5	6.6	6.2	5.3	4.1
	14 years		0.4	3.0	7.6	6.7	6.0	4.7
	15 years			0.3	3.3	8.8	8.2	6.7
	16 years				0.2	3.6	9.4	9.6
	17 years					0.1	3.0	8.2
	18 years or older						0.2	3.5
Cigars	No Answer	1.4	1.3	1.4	1.9	2.0	2.2	2.0
	Never	95.5	92.5	87.7	85.4	79.5	77.6	73.8
	7 years or younger	0.5	0.6	0.6	0.4	0.4	0.3	0.2
	8 years	0.3	0.3	0.5	0.4	0.4	0.3	0.2
	9 years	0.4	0.4	0.5	0.4	0.4	0.2	0.1
	10 years	0.6	0.7	0.9	0.8	0.7	0.5	0.2
	11 years	0.8	1.2	1.3	1.0	0.8	0.5	0.3
	12 years	0.5	1.7	2.2	1.4	1.5	1.2	0.8
	13 years	0.1	1.1	3.2	2.8	2.4	1.6	1.3
	14 years		0.1	1.5	3.4	3.4	2.4	1.8
	15 years			0.1	1.9	5.9	4.7	3.2
	16 years				0.1	2.5	6.2	6.6
	17 years					0.1	2.1	6.3
	18 years or older						0.1	3.0
Pipe	No Answer	1.6	2.0	2.6	3.1	3.5	4.2	4.1
	Never	97.9	96.4	93.8	91.0	87.7	85.0	82.7
	7 years or younger	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	8 years	0.0	0.1	0.1	0.0	0.1	0.2	0.1
	9 years	0.1	0.1	0.1	0.1	0.1	0.0	0.1
	10 years	0.1	0.2	0.2	0.3	0.3	0.1	0.1
	11 years	0.1	0.2	0.3	0.4	0.3	0.3	0.1
	12 years	0.1	0.4	0.8	0.7	0.7	0.7	0.3
	13 years	0.1	0.5	1.3	1.4	1.1	0.8	0.7
	14 years		0.0	0.6	1.7	1.7	1.2	1.2
	15 years			0.1	1.1	3.1	2.4	1.8
	16 years				0.1	1.3	3.7	3.2
	17 years					0.1	1.3	3.4
	18 years or older						0.0	2.2

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smokeless Tobacco	No Answer	1.2	1.2	1.3	1.6	1.8	2.0	1.8
	Never	97.6	96.7	95.0	93.8	91.9	91.2	91.3
	7 years or younger	0.2	0.2	0.3	0.2	0.3	0.2	0.1
	8 years	0.1	0.2	0.2	0.2	0.1	0.1	0.1
	9 years	0.1	0.2	0.2	0.2	0.1	0.2	0.1
	10 years	0.2	0.2	0.3	0.2	0.2	0.2	0.1
	11 years	0.3	0.3	0.4	0.4	0.3	0.2	0.1
	12 years	0.2	0.5	0.6	0.4	0.5	0.3	0.3
	13 years	0.0	0.5	1.1	1.1	0.9	0.6	0.4
	14 years			0.0	0.6	1.1	0.9	0.6
	15 years				0.1	0.8	1.9	1.6
	16 years					0.0	0.9	1.8
	17 years						0.1	0.8
	18 years or older							0.0
Alcohol	No Answer	2.6	2.8	3.5	3.3	3.3	3.9	3.5
	Never	78.2	67.5	52.6	46.8	37.4	34.0	29.8
	7 years or younger	3.1	3.2	2.9	1.9	1.8	1.5	1.2
	8 years	1.4	1.6	1.5	0.9	0.8	0.5	0.4
	9 years	1.8	1.4	1.4	1.2	0.9	0.6	0.3
	10 years	3.2	3.0	3.0	2.0	1.6	1.3	0.9
	11 years	5.2	4.6	4.0	2.9	2.0	1.4	1.1
	12 years	3.9	9.1	8.5	5.6	4.6	3.6	2.6
	13 years	0.5	6.3	14.6	11.7	8.6	6.9	5.5
	14 years			0.6	7.4	15.5	13.2	9.9
	15 years				0.7	7.9	18.6	16.5
	16 years					0.5	6.8	15.3
	17 years						0.3	4.4
	18 years or older							0.3
Marijuana	No Answer	1.6	1.7	1.8	2.3	2.6	2.8	2.4
	Never	96.3	91.9	83.6	78.9	70.4	68.1	65.3
	7 years or younger	0.1	0.2	0.3	0.2	0.2	0.1	0.1
	8 years	0.1	0.2	0.3	0.2	0.2	0.1	0.1
	9 years	0.2	0.2	0.4	0.3	0.2	0.2	0.1
	10 years	0.3	0.5	0.7	0.5	0.4	0.4	0.3
	11 years	0.6	1.0	1.2	1.0	1.0	0.7	0.5
	12 years	0.6	2.1	2.7	2.1	2.2	1.5	1.2
	13 years	0.2	1.9	5.6	4.2	4.3	3.4	2.7
	14 years			0.3	3.0	6.5	5.9	4.5
	15 years				0.4	3.5	8.9	7.7
	16 years					0.3	3.6	8.1
	17 years						0.2	2.3
	18 years or older							0.1

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	1.4	1.5	1.5	1.9	2.2	2.4	2.1
	Never	98.2	97.5	96.5	95.3	93.5	92.5	92.7
	7 years or younger	0.1	0.1	0.1	0.0	0.1	0.0	0.1
	8 years	0.0	0.0	0.0		0.0	0.0	0.0
	9 years	0.0		0.0	0.0	0.0	0.0	0.0
	10 years	0.1	0.1	0.1	0.1	0.1	0.0	0.0
	11 years	0.1	0.2	0.1	0.1	0.1	0.1	0.0
	12 years	0.1	0.2	0.2	0.2	0.2	0.1	0.1
	13 years	0.0	0.4	0.7	0.5	0.4	0.4	0.2
	14 years		0.0	0.6	0.9	0.7	0.6	0.5
	15 years			0.1	0.8	1.8	1.3	0.5
	16 years				0.1	0.9	1.8	1.4
	17 years					0.1	0.6	1.5
	18 years or older						0.0	0.7
Crack	No Answer	1.6	1.5	1.7	2.1	2.3	2.7	2.0
	Never	97.8	97.5	96.7	96.2	95.7	95.1	96.1
	7 years or younger	0.1	0.1	0.1	0.1	0.0	0.1	0.0
	8 years	0.0	0.0	0.0	0.0	0.0	0.0	
	9 years	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	10 years	0.1	0.1	0.1	0.0	0.1	0.1	0.0
	11 years	0.1	0.1	0.1	0.0	0.1	0.0	0.0
	12 years	0.1	0.2	0.3	0.1	0.1	0.1	0.1
	13 years	0.1	0.3	0.5	0.3	0.2	0.2	0.1
	14 years		0.1	0.4	0.5	0.4	0.3	0.2
	15 years			0.1	0.5	0.7	0.6	0.3
	16 years				0.1	0.3	0.5	0.4
	17 years					0.0	0.3	0.4
	18 years or older						0.0	0.3
Inhalants	No Answer	2.4	1.7	2.0	2.2	2.3	2.7	2.3
	Never	92.8	91.5	87.8	88.6	88.2	88.9	91.2
	7 years or younger	1.0	0.8	0.7	0.4	0.3	0.2	0.2
	8 years	0.5	0.3	0.4	0.2	0.2	0.1	0.1
	9 years	0.3	0.4	0.3	0.2	0.2	0.2	0.1
	10 years	0.7	0.8	0.8	0.4	0.4	0.3	0.1
	11 years	1.2	1.1	1.2	0.9	0.4	0.2	0.2
	12 years	0.8	1.8	2.3	1.2	1.1	0.7	0.4
	13 years	0.1	1.4	3.0	2.3	1.6	1.4	0.9
	14 years		0.2	1.4	2.4	1.9	1.4	0.9
	15 years			0.1	1.1	2.4	1.9	1.1
	16 years				0.2	0.8	1.5	1.3
	17 years					0.1	0.5	0.9
	18 years or older						0.0	0.3

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Amphetamines	No Answer	2.7	1.8	1.8	2.1	2.3	2.8	2.2
	Never	97.0	97.2	95.3	93.2	90.1	88.9	90.4
	7 years or younger	0.1	0.0	0.1	0.0	0.0	0.1	0.0
	8 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	9 years	0.0	0.0	0.1	0.0	0.1	0.0	0.0
	10 years	0.0	0.1	0.1	0.1	0.1	0.1	0.0
	11 years	0.1	0.1	0.2	0.2	0.2	0.1	0.1
	12 years	0.1	0.3	0.6	0.6	0.4	0.4	0.2
	13 years	0.0	0.3	1.0	1.0	1.3	0.8	0.5
	14 years		0.1	0.7	1.8	1.9	1.3	0.9
	15 years			0.1	0.9	2.5	2.4	1.6
	16 years				0.1	1.0	2.3	2.2
	17 years					0.1	0.7	1.4
	18 years or older						0.0	0.4
Methamphetamines	No Answer	2.4	1.7	1.8	2.0	2.4	2.7	2.1
	Never	97.3	97.4	96.9	96.3	95.4	94.7	95.6
	7 years or younger	0.0	0.1	0.0	0.0	0.0	0.1	0.0
	8 years	0.0	0.0	0.0		0.0	0.0	0.0
	9 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	10 years	0.0	0.1	0.0	0.0	0.1	0.1	0.0
	11 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	12 years	0.1	0.2	0.2	0.2	0.1	0.1	0.1
	13 years	0.0	0.3	0.5	0.3	0.3	0.3	0.2
	14 years		0.1	0.4	0.6	0.5	0.3	0.3
	15 years			0.0	0.4	0.7	0.8	0.4
	16 years				0.1	0.4	0.6	0.6
	17 years					0.0	0.2	0.4
	18 years or older						0.0	0.2
Ritalin/Adderall	No Answer	3.1	2.1	2.0	2.2	2.5	2.7	2.3
	Never	96.1	96.1	93.7	90.8	87.9	88.2	88.4
	7 years or younger	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	8 years	0.1	0.1	0.0	0.1	0.1	0.0	0.0
	9 years	0.1	0.1	0.1	0.0	0.0	0.0	0.0
	10 years	0.1	0.2	0.2	0.1	0.1	0.1	0.0
	11 years	0.2	0.2	0.4	0.3	0.2	0.2	0.1
	12 years	0.1	0.5	0.8	0.8	0.5	0.3	0.2
	13 years	0.1	0.5	1.5	1.3	1.1	0.7	0.4
	14 years		0.1	1.2	2.5	2.3	1.2	1.0
	15 years			0.1	1.6	3.5	2.5	1.9
	16 years				0.1	1.7	2.9	2.7
	17 years					0.1	1.0	2.2
	18 years or older						0.0	0.8

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Tranquilizers	No Answer	2.4	1.9	1.7	2.3	2.2	2.6	2.1
	Never	93.8	92.1	87.8	86.2	83.9	84.3	86.5
	7 years or younger	0.3	0.3	0.2	0.2	0.1	0.1	0.1
	8 years	0.2	0.1	0.1	0.1	0.1	0.1	
	9 years	0.4	0.2	0.3	0.1	0.1		0.1
	10 years	0.7	0.5	0.5	0.3	0.3	0.1	0.1
	11 years	1.1	0.9	0.8	0.4	0.3	0.2	0.1
	12 years	1.0	2.0	2.1	1.2	0.9	0.6	0.4
	13 years	0.2	1.9	3.7	2.5	2.0	1.1	0.7
	14 years		0.2	2.5	3.8	3.2	2.0	1.2
	15 years			0.3	2.6	4.7	3.8	2.5
	16 years				0.3	2.1	4.0	3.1
	17 years					0.1	1.1	2.4
	18 years or older						0.1	0.8
Narcotics	No Answer	2.4	1.7	1.6	2.0	2.1	2.5	2.0
	Never	97.0	96.3	93.7	91.3	88.5	88.2	89.3
	7 years or younger	0.1	0.1	0.1	0.0	0.1	0.1	0.0
	8 years	0.1	0.0	0.0	0.1	0.0	0.0	0.0
	9 years	0.0	0.0	0.0	0.0	0.1	0.0	
	10 years	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	11 years	0.2	0.2	0.2	0.2	0.2	0.1	0.1
	12 years	0.1	0.6	0.8	0.6	0.4	0.3	0.2
	13 years	0.1	0.8	1.8	1.3	1.2	0.7	0.3
	14 years		0.1	1.4	2.5	2.1	1.5	0.8
	15 years			0.1	1.7	3.4	2.6	1.9
	16 years				0.1	1.6	3.1	2.8
	17 years					0.1	0.7	1.7
	18 years or older						0.1	0.8
LSD Only	No Answer	2.9	1.9	1.8	2.1	2.3	2.6	2.2
	Never	97.0	97.4	96.7	95.9	95.2	94.6	95.5
	7 years or younger	0.0	0.0	0.1	0.0	0.0	0.1	0.0
	8 years	0.0		0.0	0.0	0.0	0.0	
	9 years	0.0	0.0	0.1	0.0	0.0	0.0	0.0
	10 years	0.0	0.1	0.0	0.0	0.1	0.1	0.0
	11 years	0.0	0.1	0.1	0.1	0.1	0.1	0.0
	12 years	0.1	0.2	0.3	0.2	0.2	0.1	0.1
	13 years	0.0	0.3	0.5	0.4	0.3	0.2	0.1
	14 years		0.0	0.4	0.6	0.4	0.3	0.3
	15 years			0.0	0.6	0.7	0.7	0.3
	16 years				0.1	0.6	0.8	0.8
	17 years					0.1	0.3	0.5
	18 years or older						0.0	0.2

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Psychedelics	No Answer	3.1	2.1	1.9	2.2	2.3	2.7	2.2
	Never	96.7	97.5	97.2	96.7	96.1	95.7	95.9
	7 years or younger	0.0	0.0	0.1	0.0	0.0	0.0	0.0
	8 years	0.0	0.0	0.0	0.0	0.0		
	9 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	10 years	0.0	0.1	0.1		0.0	0.1	0.0
	11 years	0.0	0.0	0.1	0.0	0.1	0.0	0.0
	12 years	0.1	0.1	0.2	0.2	0.1	0.0	0.0
	13 years	0.0	0.2	0.3	0.3	0.2	0.1	0.1
	14 years		0.0	0.2	0.4	0.3	0.2	0.1
	15 years			0.0	0.2	0.5	0.4	0.2
	16 years				0.0	0.3	0.6	0.6
	17 years					0.0	0.1	0.4
	18 years or older							0.2
Heroin	No Answer	2.4	1.6	1.7	2.1	2.3	2.5	2.1
	Never	97.3	97.7	97.4	96.9	96.5	96.3	96.9
	7 years or younger	0.0	0.1	0.1	0.0	0.0	0.0	0.0
	8 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	9 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	10 years	0.0	0.1	0.0	0.0	0.0	0.1	0.0
	11 years	0.1	0.1	0.1	0.0	0.0	0.0	0.1
	12 years	0.1	0.2	0.1	0.1	0.1	0.1	0.0
	13 years	0.0	0.2	0.3	0.2	0.2	0.1	0.0
	14 years		0.1	0.3	0.3	0.2	0.1	0.1
	15 years			0.0	0.2	0.4	0.2	0.3
	16 years				0.0	0.2	0.3	0.2
	17 years					0.0	0.1	0.2
	18 years or older							0.1
Steroids	No Answer	2.0	1.8	1.7	2.2	2.3	2.6	2.2
	Never	97.6	97.4	97.5	97.1	96.8	96.7	97.3
	7 years or younger	0.0	0.1	0.1	0.0	0.1	0.0	0.0
	8 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	9 years	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	10 years	0.0	0.1	0.0	0.0	0.1	0.1	0.0
	11 years	0.1	0.1	0.0	0.0	0.1	0.0	0.0
	12 years	0.1	0.2	0.2	0.1	0.1	0.0	0.0
	13 years	0.0	0.2	0.2	0.1	0.1	0.0	0.0
	14 years		0.0	0.2	0.2	0.1	0.1	0.0
	15 years			0.0	0.2	0.2	0.2	0.1
	16 years				0.0	0.1	0.1	0.1
	17 years					0.0	0.0	0.1
	18 years or older						0.0	0.0

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	3.3	2.1	1.9	2.2	2.3	2.6	2.3
	Never	96.6	97.1	96.2	95.3	94.4	93.8	94.3
	7 years or younger	0.0	0.0	0.1	0.0	0.0	0.0	0.0
	8 years	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	9 years	0.0	0.0	0.0	0.0	0.1		
	10 years	0.0	0.0	0.0	0.0	0.1	0.1	0.0
	11 years	0.0	0.1	0.2	0.1	0.0	0.1	0.1
	12 years	0.0	0.2	0.3	0.2	0.1	0.1	0.0
	13 years	0.0	0.3	0.7	0.5	0.4	0.2	0.1
	14 years		0.1	0.6	0.8	0.6	0.4	0.3
	15 years			0.0	0.6	1.1	0.9	0.5
	16 years				0.1	0.8	1.3	0.9
	17 years					0.0	0.5	0.8
	18 years or older						0.0	0.6
Rohypnol	No Answer	3.4	2.0	1.8	2.2	2.3	2.6	2.2
	Never	96.5	97.5	97.4	96.7	96.4	96.2	97.1
	7 years or younger	0.0	0.0	0.1	0.0	0.0	0.0	0.0
	8 years			0.0	0.0	0.0	0.0	
	9 years	0.0	0.0	0.0	0.0	0.0		
	10 years	0.0	0.0	0.1	0.0	0.0	0.0	0.0
	11 years	0.0	0.1	0.1	0.0	0.1	0.1	0.0
	12 years	0.0	0.1	0.1	0.2	0.1	0.1	0.0
	13 years	0.0	0.1	0.3	0.2	0.2	0.1	0.1
	14 years		0.0	0.2	0.4	0.3	0.2	0.1
	15 years			0.0	0.2	0.4	0.2	0.1
	16 years				0.0	0.2	0.3	0.1
	17 years					0.0	0.1	0.1
	18 years or older							0.1
GHB	No Answer	3.6	2.3	1.9	2.2	2.4	2.7	2.2
	Never	96.4	97.5	97.7	97.3	97.0	96.7	97.4
	7 years or younger	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	8 years	0.0		0.0	0.0			0.0
	9 years	0.0	0.0	0.0	0.0			
	10 years		0.0	0.0		0.1	0.0	0.0
	11 years	0.0	0.1	0.0	0.0	0.0	0.0	0.0
	12 years	0.0	0.0	0.1	0.0	0.1	0.0	0.0
	13 years	0.0	0.0	0.1	0.1	0.1	0.0	0.0
	14 years		0.0	0.1	0.1	0.1	0.0	0.1
	15 years				0.1	0.1	0.3	0.1
	16 years				0.0	0.1	0.1	0.0
	17 years					0.0	0.0	0.0
	18 years or older							0.0

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Over the counter drugs	No Answer	3.3	2.4	2.1	2.6	2.7	3.0	2.4
	Never	92.1	90.9	86.6	84.5	83.5	84.6	86.4
	7 years or younger	1.1	0.9	1.2	0.7	0.5	0.4	0.5
	8 years	0.5	0.3	0.4	0.3	0.2	0.1	0.1
	9 years	0.4	0.4	0.3	0.2	0.2	0.1	0.1
	10 years	0.8	0.6	0.7	0.5	0.4	0.3	0.2
	11 years	0.8	0.8	1.0	0.7	0.5	0.2	0.2
	12 years	0.9	1.8	1.9	1.5	1.2	0.7	0.6
	13 years	0.2	1.6	3.3	2.9	2.0	1.4	0.7
	14 years		0.2	2.4	3.8	3.0	2.0	1.2
	15 years			0.2	2.1	4.0	2.9	2.1
	16 years				0.2	1.7	3.3	2.8
	17 years					0.1	0.9	1.9
	18 years or older						0.1	0.7

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	0.4	0.4	0.3	0.3	0.2	0.2	0.2
	Never	89.3	82.3	71.1	66.7	59.7	55.9	53.7
	Once or Twice	7.6	10.0	14.5	14.3	15.5	16.3	17.1
	Occasionally	1.4	3.7	6.9	8.3	9.9	11.2	12.3
	Past Regularly	0.8	1.8	3.0	4.1	5.0	5.0	5.1
	Current Regularly	0.5	1.8	4.1	6.3	9.7	11.3	11.5
Cigars	No Answer	0.2	0.3	0.2	0.2	0.4	0.4	0.2
	Never	96.3	93.1	88.2	86.1	80.2	78.3	74.4
	1-5 times	2.8	4.7	8.1	9.1	12.8	13.9	17.4
	6-19 times	0.3	0.7	1.4	1.8	2.6	3.4	3.5
	20-40 times	0.2	0.4	0.7	0.8	1.3	1.1	1.7
Pipe	40+ times	0.2	0.8	1.4	1.9	2.7	2.8	2.9
	No Answer	0.3	0.6	0.8	0.9	1.0	1.2	1.1
	Never	98.8	97.4	95.0	92.5	89.3	87.0	84.8
	1-5 times	0.6	1.4	2.7	3.8	5.2	6.3	8.3
	6-19 times	0.1	0.4	0.7	1.3	1.7	2.3	2.5
Smokeless Tobacco	20-40 times	0.0	0.1	0.4	0.7	1.1	1.1	1.3
	40+ times	0.1	0.2	0.4	0.8	1.8	2.1	2.0
	No Answer	0.2	0.2	0.2	0.2	0.3	0.3	0.2
	Never	98.5	97.4	96.1	95.0	93.1	92.8	92.6
	1-5 times	1.0	1.9	2.9	3.8	5.0	5.1	5.5
Alcohol	6-19 times	0.1	0.2	0.4	0.4	0.6	0.7	0.7
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.2	0.2
	40+ times	0.1	0.1	0.3	0.2	0.7	0.8	0.8
	No Answer	0.7	0.7	0.7	0.9	0.7	0.9	0.9
	Never	78.1	67.1	51.9	46.3	37.1	33.7	29.7
Marijuana	1-5 times	17.7	23.6	29.4	28.5	28.8	27.2	24.3
	6-19 times	2.5	5.5	9.9	12.2	15.2	16.7	17.2
	20-40 times	0.6	1.7	4.0	5.8	8.9	9.7	11.9
	40+ times	0.4	1.5	4.0	6.3	9.3	11.8	16.0
	No Answer	0.4	0.5	0.6	0.6	0.6	0.8	0.6
Cocaine	Never	97.1	92.6	83.9	79.3	70.7	68.6	65.5
	1-5 times	1.8	4.5	8.2	9.8	12.2	13.0	14.1
	6-19 times	0.4	1.2	2.8	3.3	5.1	5.6	6.4
	20-40 times	0.1	0.5	1.6	2.4	3.4	3.5	4.4
	40+ times	0.2	0.8	2.9	4.6	7.9	8.4	8.9
	No Answer	0.4	0.4	0.3	0.4	0.4	0.6	0.5
	Never	99.2	98.5	97.5	96.3	94.6	93.6	93.7
	1-5 times	0.3	0.8	1.5	2.3	3.3	3.5	3.3
	6-19 times	0.1	0.1	0.3	0.4	0.8	1.0	1.1
	20-40 times	0.0	0.1	0.2	0.3	0.4	0.6	0.7
	40+ times	0.1	0.1	0.2	0.3	0.5	0.6	0.7

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Crack	No Answer	0.4	0.3	0.4	0.3	0.4	0.6	0.3
	Never	98.9	98.4	97.7	97.6	97.1	96.9	97.5
	1-5 times	0.6	0.9	1.4	1.5	1.7	1.7	1.3
	6-19 times	0.1	0.2	0.2	0.2	0.4	0.3	0.4
	20-40 times	0.1	0.1	0.1	0.2	0.2	0.2	0.3
	40+ times	0.0	0.1	0.2	0.2	0.2	0.3	0.2
Inhalants	No Answer	1.0	0.5	0.4	0.6	0.5	0.5	0.4
	Never	93.4	92.0	88.2	89.1	89.0	90.3	92.4
	1-5 times	4.4	5.4	8.1	7.0	7.1	6.3	5.1
	6-19 times	0.7	1.1	1.9	1.9	2.0	1.6	1.3
	20-40 times	0.2	0.5	0.7	0.8	0.7	0.7	0.5
	40+ times	0.4	0.5	0.7	0.7	0.7	0.6	0.3
Amphetamines	No Answer	1.7	0.8	0.6	0.5	0.6	0.6	0.5
	Never	97.9	98.1	96.2	93.9	90.9	90.0	91.2
	1-5 times	0.2	0.7	1.9	3.0	4.4	4.7	3.9
	6-19 times	0.1	0.2	0.6	1.3	2.0	2.3	2.1
	20-40 times	0.0	0.1	0.3	0.7	1.2	1.2	1.1
	40+ times	0.0	0.1	0.3	0.7	0.9	1.2	1.1
Methamphetamine	No Answer	1.2	0.5	0.4	0.5	0.6	0.6	0.4
	Never	98.4	98.4	97.9	97.4	96.7	96.3	97.0
	1-5 times	0.3	0.7	1.1	1.3	1.5	1.6	1.4
	6-19 times	0.0	0.2	0.3	0.3	0.4	0.4	0.4
	20-40 times	0.0	0.0	0.1	0.2	0.4	0.4	0.3
	40+ times	0.1	0.1	0.2	0.2	0.3	0.6	0.5
Ritalin/Adderall	No Answer	1.8	1.0	0.7	0.6	0.6	0.8	0.6
	Never	97.2	96.9	94.4	91.6	88.7	89.1	89.2
	1-5 times	0.8	1.5	3.1	4.6	5.9	5.4	5.5
	6-19 times	0.1	0.3	0.9	1.5	2.3	2.5	2.5
	20-40 times	0.1	0.1	0.4	0.8	1.3	1.0	1.1
	40+ times	0.1	0.2	0.5	0.9	1.2	1.2	1.0
Tranquilizers	No Answer	0.9	0.7	0.6	0.5	0.6	0.7	0.5
	Never	94.8	92.5	88.0	86.5	84.3	84.9	87.0
	1-5 times	3.5	5.0	7.8	7.9	8.6	8.2	6.7
	6-19 times	0.6	1.3	2.0	2.7	3.2	3.0	2.9
	20-40 times	0.2	0.2	0.8	1.1	1.8	1.4	1.5
	40+ times	0.2	0.3	0.7	1.3	1.5	1.8	1.4
Narcotics	No Answer	1.5	0.8	0.5	0.4	0.5	0.6	0.5
	Never	97.8	96.9	94.2	91.7	89.1	88.9	90.0
	1-5 times	0.5	1.5	3.4	4.5	5.3	5.3	5.0
	6-19 times	0.1	0.4	0.9	1.5	2.3	2.5	2.2
	20-40 times	0.1	0.1	0.5	0.9	1.4	1.2	1.2
	40+ times	0.1	0.2	0.5	0.9	1.4	1.5	1.1

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Psychedelics (All)	No Answer	1.3	0.6	0.3	0.2	0.3	0.4	0.2
	Never	98.2	98.3	97.5	96.8	96.1	95.8	96.3
	1-5 times	0.3	0.7	1.4	2.0	2.5	2.4	2.5
	6-19 times	0.1	0.2	0.4	0.5	0.5	0.6	0.5
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.3	0.2
	40+ times	0.1	0.1	0.2	0.3	0.3	0.4	0.3
LSD Only	No Answer	1.6	0.8	0.5	0.4	0.5	0.6	0.3
	Never	98.1	98.4	97.8	97.3	96.7	96.3	97.1
	1-5 times	0.1	0.5	1.1	1.6	2.0	2.1	1.9
	6-19 times	0.1	0.2	0.3	0.3	0.4	0.4	0.4
	20-40 times	0.0	0.1	0.1	0.2	0.2	0.3	0.2
	40+ times	0.1	0.1	0.2	0.2	0.2	0.3	0.2
Other Psychedelics	No Answer	2.1	1.0	0.6	0.6	0.6	0.8	0.5
	Never	97.6	98.5	98.3	98.1	97.6	97.3	97.6
	1-5 times	0.2	0.3	0.7	0.9	1.3	1.1	1.3
	6-19 times	0.0	0.1	0.2	0.3	0.3	0.4	0.3
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
	40+ times	0.1	0.1	0.1	0.1	0.2	0.3	0.2
Heroin	No Answer	1.3	0.7	0.4	0.4	0.5	0.7	0.4
	Never	98.4	98.5	98.6	98.4	98.0	97.9	98.5
	1-5 times	0.2	0.5	0.7	0.7	0.8	0.7	0.5
	6-19 times	0.1	0.1	0.1	0.2	0.3	0.3	0.2
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.1	0.2
	40+ times	0.1	0.1	0.2	0.2	0.2	0.3	0.2
Steroids	No Answer	0.9	0.7	0.6	0.5	0.6	0.7	0.6
	Never	98.6	98.4	98.5	98.7	98.4	98.5	98.8
	1-5 times	0.3	0.7	0.6	0.5	0.5	0.4	0.3
	6-19 times	0.0	0.1	0.1	0.0	0.2	0.1	0.1
	20-40 times	0.0	0.0	0.1	0.1	0.1	0.0	0.1
	40+ times	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Injected Drugs	No Answer	2.1	2.0	2.4	2.9	2.9	2.8	3.0
	Never	97.1	96.6	96.0	95.5	95.6	95.7	96.1
	1-5 times	0.5	0.8	0.9	1.0	0.9	0.8	0.4
	6-19 times	0.1	0.2	0.3	0.2	0.2	0.2	0.2
	20-40 times	0.1	0.1	0.1	0.1	0.2	0.2	0.1
	40+ times	0.1	0.2	0.2	0.2	0.3	0.3	0.2
MDMA	No Answer	2.1	1.0	0.5	0.5	0.5	0.6	0.4
	Never	97.6	98.1	97.2	96.6	95.7	95.3	95.7
	1-5 times	0.1	0.6	1.6	2.1	2.6	2.9	2.7
	6-19 times	0.0	0.2	0.3	0.4	0.6	0.6	0.6
	20-40 times	0.0	0.0	0.1	0.2	0.3	0.3	0.3
	40+ times	0.1	0.1	0.2	0.2	0.3	0.3	0.3

**Lifetime Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Rohypnol	No Answer	2.1	1.0	0.6	0.5	0.5	0.7	0.5
	Never	97.7	98.4	98.4	98.2	98.0	97.9	98.6
	1-5 times	0.1	0.4	0.7	0.8	1.0	0.9	0.6
	6-19 times	0.0	0.1	0.2	0.3	0.2	0.2	0.2
	20-40 times		0.1	0.0	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.1	0.1	0.2	0.2	0.1
GHB	No Answer	2.5	1.1	0.7	0.6	0.6	0.8	0.5
	Never	97.4	98.6	98.8	98.8	98.7	98.5	99.0
	1-5 times	0.0	0.1	0.3	0.3	0.4	0.4	0.2
	6-19 times	0.0	0.1	0.1	0.1	0.2	0.1	0.1
	20-40 times	0.0	0.0	0.0	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.1	0.1	0.1	0.2	0.1
Over the counter drugs	No Answer	1.7	1.1	0.9	0.9	0.8	0.8	0.7
	Never	92.8	91.1	86.5	84.5	83.4	85.3	86.8
	1-5 times	3.6	4.8	7.4	8.5	8.5	7.8	7.0
	6-19 times	1.0	1.5	2.8	3.0	3.6	3.0	2.7
	20-40 times	0.5	0.7	1.1	1.3	1.6	1.2	1.0
	40+ times	0.4	0.8	1.4	1.8	2.1	1.9	1.7

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	0.9	0.9	0.7	0.8	0.7	0.7	0.7
	None	94.0	88.1	80.2	76.5	71.2	67.8	67.0
	Few times	3.7	7.4	11.8	12.4	14.7	15.9	16.7
	1-5 cigarettes/day	0.9	2.2	4.1	5.7	6.6	7.6	7.7
	1/2 pack/day	0.3	0.8	1.8	2.5	3.9	4.4	4.6
	1 pack/day	0.1	0.4	0.9	1.5	2.1	2.7	2.8
	1 & 1/2 pack/day	0.1	0.1	0.2	0.4	0.5	0.6	0.3
	2+ pack/day	0.1	0.1	0.2	0.2	0.4	0.3	0.2
Cigars	No Answer	1.6	1.7	2.1	2.6	2.9	3.0	2.7
	Never	96.1	93.3	88.8	86.7	82.1	80.9	77.8
	1-5 times	1.6	2.9	5.4	6.3	8.9	10.1	13.0
	6-19 times	0.4	0.9	1.6	1.5	2.3	2.4	2.8
	20-40 times	0.1	0.4	0.7	0.9	1.1	0.9	1.0
Pipe	40+ times	0.2	0.7	1.3	2.0	2.7	2.7	2.7
	No Answer	1.8	2.0	2.6	3.1	3.4	3.5	3.5
	Never	97.5	96.1	93.4	91.0	88.2	86.3	84.6
	1-5 times	0.5	1.1	2.3	3.2	4.0	5.5	6.9
	6-19 times	0.1	0.4	0.8	1.2	1.7	1.9	2.2
Smokeless Tobacco	20-40 times	0.1	0.2	0.4	0.6	1.1	1.0	1.2
	40+ times	0.1	0.2	0.5	0.8	1.5	1.9	1.6
	No Answer	1.6	1.6	2.1	2.5	2.8	2.9	2.8
	Never	97.4	96.4	94.8	93.5	92.1	92.3	92.6
	1-5 times	0.7	1.3	2.1	2.8	3.5	3.3	3.1
Alcohol	6-19 times	0.1	0.4	0.5	0.5	0.6	0.5	0.4
	20-40 times	0.1	0.1	0.2	0.2	0.3	0.2	0.3
	40+ times	0.1	0.1	0.4	0.4	0.7	0.9	0.8
	No Answer	1.9	2.1	2.4	2.7	3.0	3.0	2.8
	Never	83.1	72.4	58.0	51.4	43.0	40.3	36.4
Marijuana	1-5 times	12.2	18.4	25.9	27.2	29.4	30.0	28.6
	6-19 times	2.1	4.5	8.4	10.9	13.3	14.0	16.0
	20-40 times	0.4	1.5	3.0	4.3	6.2	7.0	8.4
	40+ times	0.3	1.1	2.3	3.6	5.1	5.8	7.8
	No Answer	1.8	1.9	2.4	2.9	3.1	3.4	3.1
	Never	96.2	92.1	84.5	80.5	73.9	73.1	72.2
	1-5 times	1.3	3.5	7.0	8.4	10.5	11.1	11.4
	6-19 times	0.4	1.2	2.5	3.1	4.4	4.4	4.9
	20-40 times	0.1	0.4	1.5	2.1	2.9	2.8	3.1
	40+ times	0.1	0.8	2.1	3.2	5.2	5.3	5.3

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	1.8	1.9	2.3	2.7	3.3	3.3	3.0
	Never	97.9	97.1	95.9	94.9	93.3	93.1	93.2
	1-5 times	0.2	0.6	1.2	1.6	2.2	2.3	2.2
	6-19 times	0.1	0.2	0.3	0.4	0.6	0.8	0.8
	20-40 times	0.0	0.1	0.2	0.1	0.4	0.4	0.5
	40+ times	0.0	0.1	0.1	0.2	0.2	0.2	0.4
Crack	No Answer	1.8	1.9	2.3	2.7	3.1	3.2	2.9
	Never	97.7	97.1	96.2	95.8	95.2	95.2	95.6
	1-5 times	0.3	0.6	1.0	1.0	1.1	1.0	0.8
	6-19 times	0.1	0.2	0.2	0.3	0.3	0.3	0.3
	20-40 times	0.1	0.1	0.1	0.0	0.2	0.2	0.2
	40+ times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
Inhalants	No Answer	2.3	2.0	2.3	2.8	3.0	3.2	3.0
	Never	93.6	92.3	89.2	90.3	90.8	92.2	93.5
	1-5 times	2.8	3.5	5.7	4.5	4.1	3.2	2.4
	6-19 times	0.8	1.4	1.8	1.4	1.2	0.9	0.7
	20-40 times	0.3	0.5	0.6	0.5	0.5	0.4	0.3
	40+ times	0.2	0.4	0.4	0.4	0.3	0.2	0.1
Amphetamines	No Answer	2.9	2.2	2.4	2.8	3.2	3.3	3.1
	Never	96.8	96.8	95.1	93.1	90.6	90.5	91.7
	1-5 times	0.2	0.6	1.5	2.2	3.4	3.6	2.9
	6-19 times	0.1	0.2	0.6	1.0	1.6	1.4	1.2
	20-40 times	0.0	0.1	0.3	0.4	0.7	0.7	0.6
	40+ times	0.0	0.1	0.2	0.5	0.5	0.6	0.5
Methamphetamines	No Answer	2.5	2.0	2.4	2.8	3.1	3.2	3.1
	Never	97.3	97.2	96.4	95.7	95.0	95.1	95.5
	1-5 times	0.1	0.5	0.8	0.9	1.2	1.0	0.7
	6-19 times	0.0	0.1	0.2	0.3	0.3	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.3	0.2
	40+ times	0.0	0.1	0.1	0.1	0.1	0.1	0.2
Ritalin/Adderall	No Answer	3.0	2.3	2.5	2.9	3.2	3.3	3.1
	Never	96.3	96.1	93.7	91.0	88.9	90.0	90.7
	1-5 times	0.5	1.0	2.4	3.7	4.7	4.1	4.0
	6-19 times	0.1	0.2	0.7	1.3	1.8	1.4	1.3
	20-40 times	0.0	0.2	0.4	0.5	0.8	0.8	0.5
	40+ times	0.1	0.2	0.3	0.6	0.5	0.5	0.4
Tranquilizers	No Answer	2.3	2.1	2.4	2.9	3.2	3.3	3.1
	Never	94.4	92.5	88.6	87.2	85.5	86.8	88.8
	1-5 times	2.3	3.5	5.6	6.1	6.5	6.0	4.9
	6-19 times	0.6	1.2	2.0	2.0	2.7	2.2	1.8
	20-40 times	0.2	0.4	0.8	0.9	1.3	1.0	0.8
	40+ times	0.1	0.3	0.5	0.8	0.8	0.8	0.6

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	2.9	2.2	2.5	2.9	3.2	3.2	3.2
	Never	96.6	96.0	93.5	91.5	89.3	89.9	91.2
	1-5 times	0.3	1.0	2.4	3.4	4.2	3.8	3.3
	6-19 times	0.1	0.4	0.9	1.2	1.7	1.5	1.1
	20-40 times	0.0	0.2	0.5	0.5	0.8	0.9	0.8
	40+ times	0.1	0.2	0.2	0.4	0.7	0.6	0.4
Psychedelics (All)	No Answer	2.6	2.0	2.2	2.7	2.8	3.0	2.7
	Never	96.9	97.1	96.1	95.3	94.7	94.7	95.1
	1-5 times	0.1	0.4	1.0	1.4	1.7	1.4	1.6
	6-19 times	0.1	0.2	0.4	0.3	0.4	0.5	0.2
	20-40 times	0.0	0.1	0.2	0.1	0.3	0.2	0.2
	40+ times	0.2	0.2	0.2	0.3	0.2	0.3	0.3
LSD Only	No Answer	3.1	2.3	2.4	2.9	3.1	3.2	3.0
	Never	96.8	97.1	96.4	95.6	95.0	95.0	95.5
	1-5 times	0.1	0.4	0.8	1.1	1.4	1.1	1.1
	6-19 times	0.1	0.1	0.2	0.2	0.2	0.4	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.1	0.1
	40+ times	0.0	0.1	0.1	0.1	0.0	0.1	0.1
Other Psychedelics	No Answer	3.4	2.5	2.7	3.2	3.3	3.4	3.2
	Never	96.3	97.0	96.4	95.9	95.3	95.2	95.4
	1-5 times	0.1	0.2	0.5	0.6	0.8	0.8	1.0
	6-19 times	0.0	0.1	0.2	0.1	0.2	0.3	0.1
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.1	0.1
	40+ times	0.0	0.1	0.1	0.1	0.2	0.2	0.2
Heroin	No Answer	2.6	2.1	2.4	3.0	3.2	3.3	3.0
	Never	97.1	97.3	96.9	96.2	95.8	95.8	96.3
	1-5 times	0.2	0.3	0.4	0.5	0.6	0.5	0.3
	6-19 times	0.0	0.2	0.2	0.1	0.2	0.2	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.2	0.1	0.2
	40+ times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
Steroids	No Answer	2.3	2.1	2.6	3.1	3.3	3.4	3.1
	Never	97.4	97.2	96.7	96.3	96.0	96.1	96.4
	1-5 times	0.2	0.4	0.5	0.4	0.4	0.3	0.3
	6-19 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	20-40 times	0.0	0.0	0.0	0.1	0.1	0.1	0.0
	40+ times	0.1	0.0	0.1	0.1	0.1	0.1	0.1
Injected Drugs	No Answer	2.9	2.9	3.1	3.5	3.8	3.6	3.7
	Never	96.5	96.1	95.8	95.3	95.1	95.5	95.7
	1-5 times	0.3	0.6	0.7	0.8	0.6	0.4	0.3
	6-19 times	0.1	0.2	0.2	0.3	0.2	0.2	0.2
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.1	0.1	0.1	0.1	0.1	0.2	0.0

**Annual Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	3.4	2.5	2.5	3.0	3.2	3.3	3.0
	Never	96.5	96.8	95.8	95.1	94.2	94.3	94.7
	1-5 times	0.1	0.4	1.2	1.4	1.9	1.7	1.7
	6-19 times	0.0	0.2	0.3	0.3	0.4	0.5	0.3
	20-40 times	0.0	0.1	0.2	0.1	0.2	0.1	0.2
	40+ times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
Rohypnol	No Answer	3.3	2.5	2.5	3.0	3.2	3.3	3.0
	Never	96.6	97.1	96.8	96.1	95.8	95.8	96.4
	1-5 times	0.1	0.2	0.4	0.6	0.7	0.5	0.3
	6-19 times	0.0	0.1	0.3	0.2	0.2	0.2	0.1
	20-40 times	0.0	0.1	0.0	0.0	0.1	0.1	0.0
	40+ times	0.0	0.0	0.1	0.1	0.1	0.1	0.1
GHB	No Answer	3.6	2.5	2.6	3.0	3.3	3.3	3.1
	Never	96.4	97.3	97.1	96.6	96.2	96.2	96.6
	1-5 times	0.0	0.1	0.2	0.2	0.3	0.2	0.2
	6-19 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	20-40 times	0.0	0.1	0.0	0.0	0.1	0.1	0.0
	40+ times	0.0	0.0	0.0	0.0	0.0	0.1	0.1
Over the counter drugs	No Answer	2.9	2.6	2.9	3.2	3.6	3.5	3.4
	Never	92.4	90.9	86.5	85.1	84.3	86.8	88.3
	1-5 times	3.0	3.7	5.9	6.3	6.5	5.5	4.7
	6-19 times	1.1	1.6	2.8	3.0	3.2	2.2	2.1
	20-40 times	0.3	0.6	1.0	1.4	1.4	1.0	0.6
	40+ times	0.2	0.5	1.0	1.0	1.1	1.0	0.8

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cigarettes	No Answer	0.8	0.7	0.5	0.6	0.6	0.7	0.6
	None	96.6	92.9	88.1	84.9	80.7	78.0	77.5
	Few times	1.7	3.9	5.9	6.7	7.6	8.1	8.4
	1-5 cigarettes/day	0.6	1.4	3.0	4.0	5.0	6.1	6.2
	1/2 pack/day	0.1	0.5	1.2	1.9	3.2	3.7	4.1
	1 pack/day	0.1	0.3	0.8	1.2	2.1	2.6	2.5
	1 & 1/2 pack/day	0.0	0.1	0.3	0.4	0.4	0.5	0.3
	2+ pack/day	0.1	0.2	0.2	0.3	0.4	0.3	0.3
Cigars	No Answer	2.1	2.5	2.8	3.3	3.7	3.7	3.5
	Never	96.5	94.6	91.9	90.4	88.0	87.7	86.8
	1-5 times	1.0	1.8	3.3	3.5	4.9	5.3	6.4
	6-19 times	0.1	0.5	0.8	1.1	1.1	1.0	1.1
	20-40 times	0.1	0.2	0.5	0.6	0.7	0.7	0.6
Pipe	40+ times	0.1	0.4	0.7	1.2	1.6	1.6	1.7
	No Answer	2.3	2.7	3.0	3.7	4.0	3.9	3.7
	Never	97.4	96.2	94.6	93.1	91.0	90.5	90.0
	1-5 times	0.3	0.8	1.5	1.9	2.8	3.2	4.1
	6-19 times	0.0	0.2	0.5	0.7	1.0	0.9	1.1
Smokeless Tobacco	20-40 times	0.0	0.1	0.2	0.3	0.5	0.8	0.6
	40+ times	0.0	0.1	0.2	0.3	0.6	0.7	0.5
	No Answer	2.1	2.4	2.8	3.2	3.6	3.5	3.4
	Never	97.3	96.5	95.5	94.8	94.0	94.2	94.7
	1-5 times	0.4	0.7	1.1	1.4	1.4	1.3	1.0
Alcohol	6-19 times	0.1	0.3	0.2	0.2	0.4	0.3	0.2
	20-40 times	0.0	0.1	0.2	0.2	0.3	0.3	0.2
	40+ times	0.0	0.1	0.2	0.3	0.4	0.4	0.5
	No Answer	2.4	2.6	2.9	3.2	3.7	3.4	3.3
	Never	90.4	84.9	75.4	71.2	65.7	63.8	58.6
Marijuana	1-5 times	6.2	10.2	17.2	19.8	23.0	24.6	27.3
	6-19 times	0.7	1.6	3.1	3.9	5.4	5.8	7.9
	20-40 times	0.2	0.5	0.9	1.3	1.4	1.5	1.9
	40+ times	0.1	0.3	0.6	0.6	0.8	0.9	1.0
	No Answer	2.3	2.6	2.9	3.4	3.8	3.8	3.6
	Never	96.5	93.7	88.9	86.9	82.6	82.7	82.5
	1-5 times	0.9	2.4	4.9	5.9	7.3	7.1	7.8
	6-19 times	0.2	0.7	1.6	1.8	2.7	2.9	2.5
	20-40 times	0.1	0.4	0.8	1.0	1.7	1.7	1.6
	40+ times	0.0	0.3	0.9	1.0	1.9	1.8	2.0

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Cocaine	No Answer	2.4	2.6	2.9	3.5	3.9	3.8	3.7
	Never	97.5	96.7	96.2	95.4	94.5	94.6	94.7
	1-5 times	0.1	0.5	0.7	0.9	1.1	1.1	1.2
	6-19 times	0.0	0.1	0.1	0.1	0.3	0.3	0.3
	20-40 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	40+ times	0.0	0.0	0.0	0.1	0.1	0.1	0.1
Crack	No Answer	2.3	2.6	3.0	3.5	3.9	3.8	3.6
	Never	97.3	96.9	96.2	95.8	95.4	95.4	95.7
	1-5 times	0.3	0.3	0.6	0.6	0.4	0.5	0.4
	6-19 times	0.0	0.1	0.1	0.1	0.2	0.2	0.2
	20-40 times	0.0	0.1	0.1	0.0	0.0	0.0	0.0
	40+ times	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Inhalants	No Answer	2.9	2.7	3.0	3.6	3.9	3.8	3.7
	Never	94.7	93.9	92.3	93.2	93.5	94.2	95.2
	1-5 times	1.9	2.4	3.6	2.6	2.0	1.7	0.9
	6-19 times	0.4	0.6	0.6	0.4	0.4	0.3	0.2
	20-40 times	0.1	0.2	0.3	0.2	0.1	0.0	0.0
	40+ times	0.1	0.1	0.1	0.1	0.1	0.0	0.0
Amphetamines	No Answer	3.4	2.9	3.1	3.6	4.0	3.8	3.7
	Never	96.4	96.3	95.4	94.2	92.9	93.1	94.0
	1-5 times	0.1	0.5	1.1	1.5	2.1	2.3	1.5
	6-19 times	0.0	0.1	0.3	0.5	0.6	0.6	0.6
	20-40 times	0.0	0.0	0.1	0.1	0.2	0.2	0.1
	40+ times	0.0	0.0	0.0	0.1	0.1	0.1	0.0
Methamphetamines	No Answer	3.0	2.7	3.1	3.7	4.1	3.9	3.8
	Never	96.9	96.7	96.2	95.6	95.0	95.2	95.6
	1-5 times	0.1	0.4	0.4	0.6	0.6	0.5	0.4
	6-19 times	0.0	0.1	0.2	0.1	0.2	0.2	0.2
	20-40 times	0.0	0.1	0.0	0.0	0.1	0.1	0.0
	40+ times	0.0	0.0	0.0	0.0	0.1	0.0	0.0
Ritalin/Adderall	No Answer	3.5	3.0	3.1	3.6	4.0	3.9	3.7
	Never	96.0	96.0	94.7	93.2	92.3	92.9	93.8
	1-5 times	0.3	0.7	1.5	2.3	2.6	2.2	1.8
	6-19 times	0.1	0.2	0.4	0.5	0.7	0.6	0.5
	20-40 times	0.0	0.1	0.1	0.2	0.3	0.2	0.2
	40+ times	0.0	0.1	0.1	0.2	0.1	0.1	0.1
Tranquilizers	No Answer	2.8	2.8	3.1	3.6	4.1	3.9	3.8
	Never	95.3	93.8	91.6	90.9	90.0	91.1	92.2
	1-5 times	1.4	2.5	4.0	3.8	4.1	3.5	2.9
	6-19 times	0.3	0.7	0.9	1.0	1.1	0.9	0.7
	20-40 times	0.1	0.1	0.3	0.4	0.4	0.3	0.3
	40+ times	0.0	0.1	0.1	0.3	0.2	0.2	0.1

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Narcotics	No Answer	3.3	3.0	3.2	3.6	4.0	3.9	3.8
	Never	96.4	95.9	94.5	93.5	92.1	92.6	93.4
	1-5 times	0.2	0.8	1.5	2.0	2.6	2.3	1.9
	6-19 times	0.1	0.2	0.5	0.5	0.9	0.7	0.6
	20-40 times	0.0	0.0	0.2	0.2	0.3	0.3	0.1
	40+ times	0.0	0.0	0.1	0.1	0.2	0.2	0.2
Psychedelics (All)	No Answer	3.3	2.8	3.0	3.5	3.8	3.7	3.6
	Never	96.6	96.6	96.0	95.6	95.1	95.2	95.6
	1-5 times	0.1	0.4	0.7	0.7	0.8	0.8	0.5
	6-19 times	0.0	0.1	0.2	0.2	0.2	0.2	0.1
	20-40 times	0.0	0.1	0.1	0.0	0.1	0.1	0.0
	40+ times	0.0	0.0	0.0	0.0	0.1	0.1	0.1
LSD Only	No Answer	3.5	3.0	3.2	3.7	4.1	3.9	3.8
	Never	96.4	96.5	96.1	95.5	95.0	95.3	95.8
	1-5 times	0.1	0.3	0.5	0.6	0.6	0.6	0.3
	6-19 times	0.0	0.1	0.1	0.1	0.2	0.1	0.1
	20-40 times	0.0	0.0	0.1	0.0	0.1	0.0	0.0
	40+ times	0.0	0.0	0.0	0.0	0.0	0.1	0.1
Other Psychedelics	No Answer	3.7	3.1	3.3	3.8	4.0	3.9	3.8
	Never	96.2	96.6	96.2	95.8	95.3	95.5	95.7
	1-5 times	0.1	0.2	0.4	0.3	0.5	0.4	0.3
	6-19 times	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	20-40 times	0.0	0.1	0.0	0.0	0.0	0.1	0.0
	40+ times	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	No Answer	3.1	2.9	3.0	3.6	4.0	3.9	3.8
	Never	96.7	96.7	96.4	95.8	95.4	95.6	95.8
	1-5 times	0.1	0.2	0.4	0.4	0.3	0.3	0.2
	6-19 times	0.0	0.1	0.1	0.1	0.2	0.2	0.1
	20-40 times	0.0	0.1	0.0	0.0	0.1	0.0	0.0
	40+ times	0.0	0.0	0.0	0.0	0.0	0.0	0.1
Steroids	No Answer	2.7	2.8	3.2	3.7	4.2	3.9	3.8
	Never	97.0	96.7	96.4	95.9	95.5	95.9	95.9
	1-5 times	0.2	0.4	0.2	0.3	0.2	0.2	0.1
	6-19 times	0.0	0.1	0.1	0.1	0.1	0.0	0.1
	20-40 times	0.0	0.0	0.0	0.0	0.1	0.0	0.1
	40+ times	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Injected Drugs	No Answer	3.0	2.9	3.1	3.5	3.8	3.6	3.7
	Never	96.7	96.5	96.2	95.9	95.6	95.8	96.0
	1-5 times	0.2	0.3	0.4	0.4	0.3	0.2	0.2
	6-19 times	0.0	0.1	0.1	0.1	0.1	0.2	0.1
	20-40 times	0.0	0.1	0.0	0.0	0.1	0.1	0.0
	40+ times	0.1	0.1	0.1	0.0	0.1	0.1	0.0

**Monthly Use of Alcohol, Tobacco, and Other Drugs  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
MDMA	No Answer	3.8	3.1	3.2	3.6	4.1	3.9	3.7
	Never	96.1	96.3	95.9	95.5	94.7	95.0	95.4
	1-5 times	0.1	0.4	0.7	0.7	0.9	0.8	0.7
	6-19 times	0.0	0.1	0.2	0.1	0.2	0.2	0.1
	20-40 times	0.0	0.0	0.0	0.1	0.0	0.1	0.0
	40+ times	0.0	0.0	0.0	0.0	0.1	0.1	0.0
Rohypnol	No Answer	3.8	3.1	3.3	3.8	4.1	3.9	3.9
	Never	96.2	96.7	96.3	95.7	95.4	95.6	95.8
	1-5 times	0.0	0.1	0.3	0.4	0.4	0.3	0.2
	6-19 times	0.0	0.0	0.1	0.0	0.1	0.1	0.0
	20-40 times	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	40+ times	0.0	0.0	0.0	0.0	0.0	0.0	0.1
GHB	No Answer	4.0	3.2	3.3	3.7	4.2	4.0	3.9
	Never	96.0	96.7	96.4	96.0	95.5	95.7	95.9
	1-5 times	0.0	0.1	0.2	0.2	0.2	0.2	0.2
	6-19 times	0.1	0.1	0.0	0.1	0.0	0.0	0.1
	20-40 times	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	40+ times	0.0			0.0	0.0	0.0	0.0
Over the counter drugs	No Answer	3.3	3.2	3.5	4.1	4.5	4.2	4.0
	Never	93.7	92.4	89.5	88.7	88.3	90.7	91.6
	1-5 times	2.3	3.1	5.0	5.1	5.1	3.6	3.2
	6-19 times	0.5	0.8	1.3	1.4	1.3	0.9	0.7
	20-40 times	0.1	0.4	0.5	0.5	0.5	0.3	0.3
	40+ times	0.1	0.2	0.2	0.3	0.4	0.3	0.2

**Binge Drinking in the Past Two Weeks  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Binge drinking	No Answer	5.3	4.2	3.5	4.3	4.3	4.0	3.9
	None	89.9	87.5	82.7	79.4	76.3	76.4	71.9
	Once	2.6	4.2	6.6	7.7	8.8	8.6	10.6
	Twice	1.1	1.9	3.2	3.9	4.5	4.8	6.1
	3-5 times	0.7	1.3	2.4	3.0	4.0	3.9	5.4
	6-9 times	0.2	0.5	0.8	1.0	1.0	1.0	1.1
	10+ times	0.2	0.4	0.8	0.8	1.2	1.2	1.0

**Main Sources of Alcoholic Beverages in the Past Month  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Main sources of alcoholic beverages	No Answer	10.2	8.9	8.5	10.2	10.3	11.0	11.5
	No drink	82.8	78.9	71.3	66.9	62.3	60.6	55.0
	Liquor stores/ convenience stores/ supermarkets	0.1	0.1	0.1	0.1	0.3	0.3	0.6
	Restaurants/bars/ clubs	0.0	0.0	0.1	0.1	0.1	0.2	0.5
	Public events	0.0	0.1	0.1	0.1	0.1	0.1	0.1
	Had someone else buy it	0.2	0.8	2.3	3.8	7.2	8.9	11.4
	Received from person 21 or older	1.0	1.6	3.4	4.4	6.1	7.5	10.3
	Took it from a store	0.1	0.1	0.1	0.2	0.2	0.2	0.1
	Family members	3.1	4.2	6.1	5.4	4.8	4.0	3.9
	Other ways	2.5	5.3	8.0	8.9	8.6	7.2	6.6

**Most Important Reasons for Drinking  
by Indiana Female 6th - 12th Graders, 2007**  
(Values expressed as percentages, multiple responses allowed)

	Grade						
	6th	7th	8th	9th	10th	11th	12th
Did not drink alcoholic beverages	81.5	70.7	56.0	50.0	42.1	39.4	34.3
To experiment	8.4	10.1	11.2	9.7	8.4	7.0	6.4
To relax or relieve tension	0.3	0.4	0.7	0.8	1.1	1.3	2.1
To feel good or get high	0.3	0.4	0.6	0.7	1.0	0.9	1.0
To seek deeper insights or understanding	0.1	0.1	0.1	0.1	0.2	0.1	0.2
To have a good time with friends	0.9	2.4	4.7	7.7	11.6	14.6	18.9
To fit in with a group I like	0.4	0.5	0.6	0.7	0.9	0.9	0.9
To get away from my problems	0.7	1.1	2.2	2.7	3.1	3.3	3.7
Because of boredom	0.8	1.9	3.3	4.1	5.4	5.1	5.3
Because of anger	1.0	1.9	3.1	3.7	4.2	4.0	3.5
To get through the day	0.2	0.4	0.7	0.7	0.7	0.4	0.4
To increase the effects of other drugs	0.0	0.2	0.4	0.6	0.8	1.0	0.8
To decrease the effects of other drugs	0.1	0.0	0.1	0.1	0.1	0.1	0.0
To get to sleep	0.3	0.4	0.7	0.8	1.0	1.0	1.2
Because it tastes good	4.5	8.6	14.2	16.0	17.6	19.6	20.1
Because I am hooked	0.6	1.1	1.5	1.6	1.6	1.5	1.0

**Consequences of ATOD Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had a hangover	No Answer	3.8	2.9	2.9	3.0	3.2	3.2	3.1
	Never	92.4	87.5	77.8	72.7	65.1	62.1	58.4
	Once	1.9	4.7	8.4	10.3	12.2	13.3	13.9
	Twice	0.9	2.0	4.3	5.3	7.2	7.9	8.1
	3-5 times	0.5	1.6	3.6	4.6	6.9	7.3	8.9
	6-10 times	0.1	0.5	1.4	1.9	2.5	2.7	4.0
	11 or more times	0.3	0.7	1.7	2.3	3.0	3.5	3.6
Had nausea or vomited	No Answer	4.2	3.3	3.1	3.3	3.4	3.3	3.3
	Never	92.2	88.8	81.5	76.0	68.9	64.8	60.9
	Once	2.1	4.5	7.6	10.4	12.5	14.0	14.9
	Twice	0.8	1.7	3.7	4.7	5.9	6.7	7.8
	3-5 times	0.5	0.9	2.4	3.1	5.6	6.9	8.5
	6-10 times	0.1	0.4	0.9	1.3	2.1	2.4	2.9
	11 or more times	0.1	0.4	0.7	1.2	1.6	1.8	1.7
Had a memory loss	No Answer	4.3	3.5	3.5	3.7	3.8	4.0	3.7
	Never	93.7	91.3	85.1	81.1	75.1	73.5	72.6
	Once	1.1	2.7	5.4	6.7	8.4	9.0	9.2
	Twice	0.4	1.1	2.3	3.1	4.2	4.4	4.8
	3-5 times	0.3	0.8	1.8	2.9	4.4	5.1	5.5
	6-10 times	0.1	0.3	0.9	1.2	2.0	1.9	2.3
	11 or more times	0.2	0.4	1.0	1.4	2.1	2.2	1.9
Poor on school test	No Answer	4.5	3.5	3.5	3.6	3.9	3.7	3.7
	Never	93.2	91.9	89.0	88.9	88.1	89.3	90.6
	Once	1.1	2.1	3.1	2.8	2.4	2.4	2.0
	Twice	0.4	0.9	1.3	1.4	1.6	1.2	1.0
	3-5 times	0.4	0.8	1.4	1.5	1.9	1.6	1.3
	6-10 times	0.1	0.3	0.7	0.7	0.8	0.6	0.5
	11 or more times	0.3	0.5	1.1	1.2	1.4	1.2	0.8
Missed school	No Answer	4.4	3.6	3.4	3.8	3.8	3.8	3.7
	Never	93.5	92.7	90.6	89.7	88.3	89.3	89.5
	Once	0.8	1.5	2.3	2.4	2.8	2.3	2.6
	Twice	0.4	0.7	1.0	1.3	1.4	1.3	1.4
	3-5 times	0.4	0.7	1.1	1.2	1.7	1.5	1.3
	6-10 times	0.2	0.4	0.8	0.6	0.8	0.8	0.7
	11 or more times	0.3	0.4	1.0	0.9	1.2	1.0	0.9
Got into a fight	No Answer	4.4	3.5	3.5	3.8	3.8	3.8	3.7
	Never	91.4	88.9	83.8	82.5	79.2	79.8	80.2
	Once	2.1	3.7	5.6	6.0	7.3	7.2	7.0
	Twice	0.8	1.4	2.5	2.9	3.9	4.2	3.7
	3-5 times	0.5	1.2	2.1	2.4	3.0	2.7	3.4
	6-10 times	0.2	0.5	0.9	1.0	1.3	0.9	1.0
	11 or more times	0.6	0.8	1.5	1.4	1.5	1.3	1.0

**Consequences of ATOD Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Damaged property	No Answer	4.5	3.6	3.6	3.9	4.0	4.0	3.7
	Never	94.5	94.4	92.4	92.3	91.9	92.9	93.7
	Once	0.5	1.0	1.9	1.6	1.7	1.5	1.3
	Twice	0.2	0.4	0.7	0.9	1.0	0.6	0.5
	3-5 times	0.1	0.4	0.7	0.6	0.8	0.6	0.5
	6-10 times	0.0	0.1	0.4	0.2	0.2	0.2	0.1
	11 or more times	0.1	0.2	0.4	0.4	0.4	0.3	0.2
Had trouble with parents	No Answer	4.5	3.6	3.5	3.7	3.9	3.8	3.6
	Never	92.1	89.6	84.9	83.4	80.8	81.3	83.0
	Once	1.6	3.5	5.7	6.6	8.0	8.4	7.4
	Twice	0.6	1.1	1.9	2.4	2.8	2.9	2.8
	3-5 times	0.4	1.0	1.6	1.7	2.1	1.8	1.9
	6-10 times	0.1	0.3	0.7	0.6	0.9	0.7	0.6
	11 or more times	0.6	1.0	1.6	1.6	1.4	1.1	0.8
Had trouble with police	No Answer	4.5	3.6	3.6	3.8	4.0	4.1	3.8
	Never	94.2	93.7	91.7	90.8	90.0	90.3	90.9
	Once	0.7	1.4	2.2	2.9	3.5	3.5	3.9
	Twice	0.3	0.6	0.9	1.1	1.1	1.0	0.7
	3-5 times	0.2	0.3	0.9	0.7	0.8	0.6	0.3
	6-10 times	0.0	0.1	0.3	0.3	0.2	0.2	0.1
	11 or more times	0.1	0.3	0.4	0.4	0.4	0.4	0.2
Driven under the influence	No Answer	4.6	3.9	4.1	4.3	4.4	4.4	4.1
	Never	94.5	94.5	92.4	91.7	88.2	83.3	80.5
	Once	0.4	0.8	1.6	1.9	3.7	5.3	6.2
	Twice	0.1	0.2	0.7	0.8	1.6	2.4	3.1
	3-5 times	0.2	0.3	0.6	0.7	1.2	2.3	3.2
	6-10 times	0.0	0.1	0.3	0.3	0.4	1.1	1.2
	11 or more times	0.1	0.2	0.4	0.4	0.7	1.3	1.8
Rode with a drunk driver	No Answer	4.4	3.4	3.5	3.8	3.9	3.9	3.7
	Never	90.8	87.7	80.1	76.3	70.3	69.7	69.8
	Once	1.8	3.1	4.8	5.7	7.5	7.7	7.7
	Twice	0.8	1.6	2.9	3.7	4.7	4.7	5.0
	3-5 times	0.8	1.6	3.3	4.6	5.5	6.0	5.9
	6-10 times	0.3	0.8	1.6	1.7	2.6	2.5	2.7
	11 or more times	1.0	1.8	3.7	4.1	5.4	5.5	5.3
Forgot things you did while high	No Answer	4.6	3.7	3.9	3.9	4.2	4.3	3.8
	Never	92.8	89.2	81.4	76.7	70.3	69.1	69.0
	Once	1.5	3.6	6.4	8.1	9.4	9.7	9.7
	Twice	0.5	1.5	2.8	4.0	5.3	5.3	5.7
	3-5 times	0.3	1.1	2.8	3.6	5.3	6.0	6.1
	6-10 times	0.1	0.4	1.1	1.8	2.3	2.5	2.8
	11 or more times	0.2	0.5	1.5	1.9	3.1	3.2	2.9

**Consequences of ATOD Use  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Had been told to cut down	No Answer	4.5	3.7	3.7	3.9	4.0	3.9	3.8
	Never	94.0	93.0	90.3	88.6	86.7	87.3	87.4
	Once	0.6	1.4	2.2	2.6	3.2	2.9	2.9
	Twice	0.2	0.6	1.1	1.5	1.8	1.7	1.8
	3-5 times	0.3	0.5	1.0	1.5	1.9	1.7	2.3
	6-10 times	0.1	0.2	0.6	0.7	0.8	0.8	0.6
	11 or more times	0.3	0.5	1.2	1.2	1.5	1.6	1.2
Used alcohol or drugs to relax or fit in	No Answer	4.5	3.6	3.8	3.9	3.9	4.0	3.8
	Never	92.0	89.2	83.3	80.3	76.3	75.9	75.9
	Once	1.6	3.1	5.3	6.4	6.9	6.5	6.7
	Twice	0.7	1.4	2.4	2.8	3.9	3.7	3.7
	3-5 times	0.5	1.3	2.4	3.0	4.0	4.4	4.8
	6-10 times	0.2	0.5	1.0	1.2	1.8	1.9	2.1
	11 or more times	0.4	0.9	1.8	2.4	3.3	3.5	2.9
Used alcohol or drugs alone	No Answer	4.4	3.5	3.6	3.9	3.9	4.0	3.6
	Never	91.6	88.3	81.2	79.2	76.5	77.9	79.8
	Once	1.8	3.6	5.9	6.3	6.4	5.8	6.0
	Twice	0.9	1.6	3.3	3.3	4.3	3.8	3.3
	3-5 times	0.7	1.6	2.9	3.7	4.2	4.0	3.4
	6-10 times	0.2	0.6	1.3	1.5	1.8	1.6	1.5
	11 or more times	0.4	0.8	1.9	2.1	2.8	2.9	2.4

**Perceived Risk of Selected Substance Use Behavior  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	3.3	2.7	3.0	3.6	3.6	3.5	3.8
	No Risk	7.4	6.9	6.1	5.0	3.9	3.7	3.4
	Slight Risk	14.5	12.6	12.3	10.5	9.3	8.7	7.4
	Moderate Risk	27.5	27.9	28.2	26.8	25.2	24.9	23.3
	Great Risk	47.2	49.7	50.4	54.2	57.9	59.3	62.1
Smoke marijuana occasionally	No Answer	3.5	3.0	3.2	3.7	3.9	3.8	3.9
	No Risk	7.4	8.0	9.2	10.8	12.4	12.7	12.8
	Slight Risk	11.4	12.8	17.9	19.6	23.1	26.1	26.3
	Moderate Risk	33.0	34.0	33.1	31.4	29.6	28.5	29.8
	Great Risk	44.6	42.1	36.5	34.4	31.0	28.8	27.3
Smoke marijuana regularly	No Answer	3.6	3.1	3.3	3.7	3.9	3.9	4.1
	No Risk	7.2	6.7	6.4	6.6	6.8	6.2	5.8
	Slight Risk	3.9	4.3	6.9	8.1	9.8	11.3	11.8
	Moderate Risk	13.6	13.7	16.8	18.1	20.4	23.2	23.4
	Great Risk	71.7	72.2	66.6	63.4	59.1	55.4	54.9
Use cocaine occasionally	No Answer	3.9	3.3	3.4	4.0	4.0	3.9	4.1
	No Risk	7.2	6.1	4.9	4.1	3.2	2.9	2.8
	Slight Risk	9.4	8.5	8.3	7.3	6.2	5.1	4.5
	Moderate Risk	33.1	33.4	32.7	30.9	27.1	26.5	22.8
	Great Risk	46.4	48.7	50.6	53.7	59.6	61.5	65.8
Use cocaine regularly	No Answer	5.1	4.7	5.3	5.2	5.3	5.4	5.4
	No Risk	7.2	5.9	4.7	3.8	2.8	2.6	2.5
	Slight Risk	3.6	2.4	2.7	2.1	1.8	1.3	1.1
	Moderate Risk	11.8	10.6	9.3	9.2	7.5	6.7	5.6
	Great Risk	72.3	76.4	78.1	79.7	82.6	84.0	85.4
Occasionally consume 1-2 alcoholic drinks	No Answer	4.7	4.0	4.2	4.5	4.7	4.4	4.3
	No Risk	19.6	21.7	26.2	27.1	27.1	29.9	31.2
	Slight Risk	36.3	37.9	38.4	38.2	39.3	39.5	40.5
	Moderate Risk	21.1	20.2	17.6	16.8	15.9	14.3	13.0
	Great Risk	18.4	16.3	13.6	13.5	13.0	12.0	11.0
Binge drink weekly	No Answer	3.8	3.3	3.5	3.9	4.1	3.9	4.0
	No Risk	8.3	7.8	7.8	7.0	6.1	6.1	5.7
	Slight Risk	11.3	11.9	14.9	15.4	15.0	15.7	16.4
	Moderate Risk	27.4	29.0	29.9	30.2	30.7	32.4	33.0
	Great Risk	49.2	48.0	44.0	43.5	44.1	41.8	40.8

**Perceived Peer Approval of Using Selected Substances  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	5.4	4.4	4.1	4.4	4.4	4.5	4.7
	Strongly Approve	2.6	2.6	2.4	2.5	2.6	2.5	1.8
	Approve	1.3	2.5	4.4	5.3	6.3	7.2	7.6
	Do Not Know	10.9	12.0	14.5	14.5	15.7	15.5	15.9
	Disapprove	12.6	13.7	16.5	17.3	18.9	19.8	21.4
Smoke marijuana occasionally	Strongly Disapprove	67.2	64.8	58.1	56.1	52.1	50.4	48.6
	No Answer	5.7	4.6	4.3	4.5	4.6	4.8	4.9
	Strongly Approve	2.8	2.9	3.3	3.5	4.5	3.9	3.7
	Approve	1.3	3.1	7.3	9.2	12.6	14.0	14.7
	Do Not Know	8.2	9.5	12.2	12.9	14.0	14.5	15.3
Smoke marijuana regularly	Disapprove	9.6	10.1	12.6	13.5	14.9	15.7	17.0
	Strongly Disapprove	72.4	69.7	60.3	56.4	49.4	47.1	44.5
	No Answer	5.6	4.8	4.3	4.6	4.8	4.8	4.9
	Strongly Approve	2.7	2.8	3.2	3.5	4.2	3.6	3.2
	Approve	0.9	1.7	4.3	5.2	7.4	7.7	7.4
Use cocaine occasionally	Do Not Know	7.9	9.1	11.1	11.8	12.2	12.7	13.0
	Disapprove	6.5	7.1	10.0	10.9	12.7	14.0	14.8
	Strongly Disapprove	76.4	74.5	67.1	64.0	58.7	57.2	56.6
	No Answer	5.7	4.8	4.4	4.8	4.8	4.8	5.0
	Strongly Approve	2.5	2.1	1.7	1.8	1.8	1.4	1.3
Use cocaine regularly	Approve	0.8	1.0	1.4	1.4	1.6	1.5	1.4
	Do Not Know	7.7	8.4	9.5	9.3	8.7	8.8	8.2
	Disapprove	9.1	9.7	11.2	11.3	11.2	11.0	10.0
	Strongly Disapprove	74.2	74.0	71.7	71.5	71.9	72.6	74.1
	No Answer	6.1	5.2	5.1	5.6	5.5	5.8	6.0
Occasionally consume 1-2 alcoholic drinks	Strongly Approve	2.6	2.2	1.8	1.8	1.7	1.3	1.2
	Approve	0.5	0.8	1.1	0.9	1.1	0.8	0.6
	Do Not Know	7.7	8.1	8.9	8.8	8.2	8.5	7.6
	Disapprove	5.8	6.3	7.9	8.0	8.0	7.9	7.3
	Strongly Disapprove	77.3	77.5	75.2	74.8	75.5	75.7	77.3
Binge drink weekly	No Answer	6.0	4.8	4.6	4.9	4.9	5.0	5.0
	Strongly Approve	3.2	4.4	5.8	6.9	8.1	8.3	9.8
	Approve	5.8	10.4	18.2	22.9	29.2	32.3	35.5
	Do Not Know	15.0	17.2	20.7	21.2	21.2	20.6	20.2
	Disapprove	14.0	14.3	14.2	14.0	12.8	12.5	11.2
	Strongly Disapprove	55.9	49.0	36.5	30.2	23.9	21.2	18.3
	No Answer	5.9	4.7	4.4	4.8	4.7	4.8	5.1
	Strongly Approve	3.1	3.5	4.2	5.3	5.6	5.6	6.0
	Approve	1.6	4.1	8.0	10.7	14.6	16.3	18.2
	Do Not Know	10.2	12.2	16.1	18.2	19.3	19.5	19.8
	Disapprove	9.3	10.5	13.6	14.1	15.5	15.8	16.9
	Strongly Disapprove	70.0	64.9	53.5	46.9	40.4	37.9	34.0

**Perceived Parental Approval of Using Selected Substances  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Smoke 1+ pack cigarettes per day	No Answer	5.5	4.5	4.1	4.7	4.5	4.8	4.8
	Strongly Approve	2.4	2.2	1.7	1.5	1.5	1.3	1.2
	Approve	0.5	0.5	1.1	1.4	2.0	2.3	2.0
	Do Not Know	3.9	4.0	3.9	3.9	4.3	4.7	5.1
	Disapprove	3.8	3.6	5.2	5.9	7.1	8.3	9.7
Smoke marijuana occasionally	Strongly Disapprove	83.8	85.3	84.1	82.6	80.6	78.7	77.2
	No Answer	5.6	4.6	4.2	4.7	4.6	4.8	5.0
	Strongly Approve	2.4	2.1	1.7	1.5	1.5	1.2	1.2
	Approve	0.3	0.6	1.2	1.4	2.1	2.3	2.0
	Do Not Know	3.3	3.6	4.0	4.0	4.5	4.7	5.2
Smoke marijuana regularly	Disapprove	2.8	2.7	4.3	4.9	6.1	6.8	7.2
	Strongly Disapprove	85.6	86.5	84.7	83.5	81.2	80.2	79.4
	No Answer	5.6	4.6	4.2	4.8	4.6	4.9	5.0
	Strongly Approve	2.4	2.1	1.7	1.5	1.5	1.1	1.1
	Approve	0.2	0.3	0.7	0.8	1.3	1.3	1.1
Use cocaine occasionally	Do Not Know	3.3	3.5	3.7	3.7	4.0	4.3	4.5
	Disapprove	2.2	2.0	3.6	3.8	4.8	5.1	5.2
	Strongly Disapprove	86.4	87.4	86.0	85.4	83.8	83.2	83.0
	No Answer	5.7	4.7	4.3	4.9	4.6	4.9	5.1
	Strongly Approve	2.4	2.0	1.5	1.4	1.3	1.0	1.0
Use cocaine regularly	Approve	0.1	0.2	0.2	0.2	0.2	0.2	0.1
	Do Not Know	3.1	3.2	3.0	2.8	3.0	2.9	2.9
	Disapprove	2.5	2.1	2.9	3.0	3.2	3.3	3.1
	Strongly Disapprove	86.1	87.8	88.1	87.6	87.6	87.7	87.8
	No Answer	5.8	4.8	4.5	5.1	4.8	5.2	5.4
Occasionally consume 1-2 alcoholic drinks	Strongly Approve	2.4	2.1	1.5	1.4	1.3	1.0	1.0
	Approve	0.2	0.1	0.2	0.2	0.2	0.2	0.1
	Do Not Know	3.2	3.2	3.0	2.8	3.0	2.9	2.9
	Disapprove	1.9	1.7	2.6	2.7	2.9	3.0	2.9
	Strongly Disapprove	86.5	88.1	88.3	87.8	87.8	87.8	87.7
Binge drink weekly	No Answer	5.9	4.8	4.4	5.1	4.8	5.0	5.2
	Strongly Approve	2.5	2.2	1.8	1.8	1.7	1.6	1.5
	Approve	3.6	4.7	6.5	7.1	8.3	9.9	13.2
	Do Not Know	7.2	8.1	9.5	9.4	9.9	10.2	12.4
	Disapprove	6.4	7.4	9.9	11.8	13.0	14.1	15.6
	Strongly Disapprove	74.5	72.7	67.8	64.9	62.5	59.1	52.1
	No Answer	6.0	4.8	4.6	5.1	5.0	5.2	5.2
	Strongly Approve	2.4	2.1	1.7	1.6	1.4	1.2	1.2
	Approve	0.7	0.9	1.3	1.6	2.0	2.5	2.9
	Do Not Know	4.3	4.9	5.6	5.9	6.2	6.6	8.0
	Disapprove	3.6	4.0	6.0	7.0	8.0	8.9	11.3
	Strongly Disapprove	83.0	83.4	80.8	78.9	77.5	75.6	71.5

**Days being Absent from School in the Past Year  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No Answer	2.2	1.4	0.9	0.5	0.5	0.4	0.3
Never	12.9	12.3	10.8	12.8	10.4	9.3	7.3
Once	10.9	12.1	10.2	12.2	10.4	9.3	8.3
Twice	17.5	16.4	14.9	15.0	14.8	13.6	12.7
3-5 days	34.2	33.9	33.8	31.7	32.9	34.1	33.5
6-10 days	15.3	16.4	18.4	17.7	18.9	20.7	22.4
11-19 days	5.2	5.3	7.2	6.4	7.4	8.4	9.8
20 or more days	1.8	2.3	3.8	3.5	4.7	4.2	5.8

**Participation in Some Form of Camp  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No participation in camps	70.6	74.1	75.8	78.0	78.1	77.4	79.0
Afternoons R.O.C.K. in Indiana	7.9	5.6	4.4	2.0	1.0	0.7	0.5
Youth Leadership Programs	9.5	11.9	12.0	11.1	10.7	10.1	9.9
S.A.D.D., S.T.A.N.D. or other prevention programs	9.8	6.8	6.1	7.1	8.1	8.9	8.1
Participated in 2+ types of camps	2.1	1.7	1.7	1.8	2.2	2.9	2.3

**Participation in Organized Family Events  
by Indiana 6th - 12th Graders, 2007  
(Values expressed as percentages)**

	Grade						
	6th	7th	8th	9th	10th	11th	12th
No Answer	3.6	3.2	2.4	2.3	1.8	1.8	1.7
Never	16.2	17.7	19.7	18.4	18.8	18.6	17.9
Once per Week	38.1	40.7	42.4	44.0	45.3	47.2	48.7
Twice per Week	20.7	19.5	18.2	19.1	18.4	18.0	17.5
3+ Times per Week	21.5	18.9	17.4	16.2	15.7	14.4	14.2

**Number of Days per School Year Spent in Various Afterschool Activities  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
School sports team	No Answer	5.5	4.5	4.7	5.8	5.4	5.2	5.6
	None	33.2	34.3	37.3	41.1	46.8	51.6	56.2
	1-9 Days	17.3	12.4	10.5	7.4	5.7	4.4	4.0
	10-29 Days	9.8	6.8	5.9	3.2	2.2	2.1	1.8
	30-59 Days	10.8	11.2	10.2	6.8	5.4	5.6	5.0
	60-89 Days	8.1	10.7	10.4	10.4	9.9	8.9	8.6
	90-129 Days	8.1	10.7	11.1	12.9	12.7	11.6	10.0
	130-180 Days	7.2	9.3	9.9	12.4	12.0	10.5	8.6
School arts programs	No Answer	6.4	5.5	5.3	6.3	6.0	6.0	6.3
	None	48.1	49.9	47.9	50.7	53.1	54.7	55.4
	1-9 Days	21.1	20.7	20.1	17.1	14.8	12.6	11.0
	10-29 Days	8.5	8.1	8.4	7.1	6.5	6.3	6.2
	30-59 Days	5.6	5.6	6.4	5.3	5.4	5.5	5.6
	60-89 Days	3.3	3.3	4.0	4.6	4.8	5.0	4.9
	90-129 Days	2.7	2.5	3.1	3.8	4.4	4.4	4.8
	130-180 Days	4.2	4.5	4.7	5.0	4.9	5.5	5.9
School club/intramurals	No Answer	7.8	6.6	6.4	7.5	6.8	6.4	6.7
	None	50.4	53.0	52.5	49.8	48.2	46.6	44.8
	1-9 Days	17.7	17.8	16.8	17.1	16.7	17.0	16.1
	10-29 Days	9.8	9.2	9.0	10.0	11.3	12.0	13.4
	30-59 Days	5.9	5.6	6.5	6.6	7.1	8.1	8.5
	60-89 Days	3.0	3.4	3.7	4.0	4.5	4.9	4.8
	90-129 Days	2.5	2.2	2.5	2.7	3.0	3.0	3.1
	130-180 Days	2.8	2.3	2.5	2.2	2.4	2.0	2.5
Youth centers (YMCA, etc)	No Answer	8.0	6.3	6.0	7.2	6.7	6.3	6.7
	None	63.2	65.3	66.9	69.1	71.3	73.5	75.8
	1-9 Days	13.8	13.1	12.5	12.0	10.5	9.7	8.5
	10-29 Days	5.4	5.5	5.3	4.2	4.3	3.9	3.9
	30-59 Days	3.4	4.1	3.5	3.1	3.0	3.0	2.1
	60-89 Days	2.3	2.2	2.2	1.9	1.8	1.7	1.5
	90-129 Days	1.5	1.6	1.6	1.3	1.2	1.0	0.8
	130-180 Days	2.6	2.0	1.9	1.2	1.1	0.9	0.7
Extended day programs	No Answer	9.3	7.8	7.3	8.2	7.7	7.1	7.4
	None	56.8	61.9	63.6	68.3	72.1	75.6	79.0
	1-9 Days	19.1	18.4	17.2	14.8	13.0	10.9	7.9
	10-29 Days	6.5	5.4	5.4	4.1	3.2	2.9	2.6
	30-59 Days	3.3	2.7	2.7	2.0	1.7	1.6	1.3
	60-89 Days	1.7	1.5	1.4	1.2	1.0	0.8	0.8
	90-129 Days	1.4	1.0	1.0	0.8	0.6	0.5	0.5
	130-180 Days	2.0	1.3	1.3	0.7	0.7	0.6	0.5

**Number of Days per School Year Spent in Various Afterschool Activities  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Supervised youth activities (4H, scouts)	No Answer	10.5	9.5	8.9	10.0	9.1	8.5	8.4
	None	51.3	53.9	58.8	62.1	64.5	67.3	70.3
	1-9 Days	13.3	11.6	10.6	9.3	9.0	8.2	7.4
	10-29 Days	7.5	7.4	6.7	5.9	5.9	5.6	5.3
	30-59 Days	6.3	6.6	5.6	5.1	4.6	4.5	3.8
	60-89 Days	4.1	4.4	4.0	3.5	3.0	3.0	2.3
	90-129 Days	3.3	3.2	2.7	2.2	2.0	1.7	1.4
	130-180 Days	3.7	3.3	2.7	2.0	1.8	1.3	1.2
Afterschool classes (art, sport, etc)	No Answer	8.8	7.5	6.8	8.1	7.4	6.9	7.0
	None	29.0	32.1	35.1	38.6	42.9	47.2	52.6
	1-9 Days	18.7	16.3	16.0	13.9	12.9	12.3	10.0
	10-29 Days	10.0	9.7	8.9	8.2	7.6	7.0	7.4
	30-59 Days	9.3	8.9	8.6	8.0	7.4	7.3	6.7
	60-89 Days	6.9	7.7	7.0	6.8	6.6	6.3	4.8
	90-129 Days	6.9	7.1	6.9	6.6	6.3	5.1	4.3
	130-180 Days	10.4	10.7	10.6	9.7	8.9	7.9	7.3
Socialization without adult supervision	No Answer	6.6	5.5	4.9	6.0	5.5	5.3	6.0
	None	33.9	20.3	12.4	9.5	6.6	5.1	5.0
	1-9 Days	21.8	20.8	17.2	13.7	11.0	9.1	8.1
	10-29 Days	10.7	13.3	13.9	13.6	11.8	10.9	10.7
	30-59 Days	7.8	10.9	12.8	15.1	14.3	14.9	15.4
	60-89 Days	5.7	9.0	12.4	14.3	16.0	16.4	16.6
	90-129 Days	5.3	8.7	11.7	13.6	16.8	18.7	18.5
	130-180 Days	8.2	11.5	14.7	14.2	18.0	19.6	19.7
At home without adult supervision	No Answer	8.1	6.7	6.2	7.2	6.5	6.4	7.0
	None	28.1	18.1	12.6	9.6	7.9	6.9	6.2
	1-9 Days	29.0	25.0	20.4	17.4	14.4	13.6	12.4
	10-29 Days	12.3	15.2	16.1	15.6	15.6	15.6	14.6
	30-59 Days	7.4	11.3	13.6	15.9	16.2	16.8	17.3
	60-89 Days	5.2	8.2	10.3	12.8	14.5	15.5	15.3
	90-129 Days	4.1	7.2	9.8	11.5	13.1	13.4	13.5
	130-180 Days	5.7	8.3	11.1	10.1	11.9	11.8	13.8
At home with adult supervision	No Answer	8.7	7.3	7.0	7.6	7.0	6.8	7.3
	None	8.6	7.1	6.5	5.7	6.0	5.6	7.0
	1-9 Days	13.5	11.4	10.7	10.0	9.2	9.2	9.6
	10-29 Days	8.9	9.6	10.5	10.5	10.2	10.0	11.4
	30-59 Days	8.0	10.9	11.8	13.2	13.9	14.7	15.8
	60-89 Days	8.8	10.7	13.5	14.9	16.3	16.5	16.6
	90-129 Days	16.9	18.0	19.8	20.6	20.3	20.9	18.7
	130-180 Days	26.5	24.8	20.1	17.5	17.1	16.2	13.7

**Number of Days per School Year Spent in Various Afterschool Activities  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Work for pay outside home	No Answer	8.4	6.7	6.4	7.3	6.5	6.0	6.4
	None	54.8	50.3	47.9	48.2	40.0	29.1	21.4
	1-9 Days	18.3	19.0	17.8	14.3	10.3	6.5	5.3
	10-29 Days	7.9	10.0	11.4	10.5	8.5	6.7	6.4
	30-59 Days	4.1	6.1	7.7	8.4	9.8	10.1	9.8
	60-89 Days	2.3	3.2	3.8	5.2	9.8	12.9	14.0
	90-129 Days	1.5	2.2	2.4	3.7	8.3	15.7	18.0
	130-180 Days	2.6	2.4	2.6	2.5	6.7	13.2	18.7
Work for pay inside a home	No Answer	7.5	6.3	6.1	7.2	6.8	6.7	7.0
	None	34.5	33.6	33.6	37.4	41.4	47.0	52.0
	1-9 Days	22.3	21.2	20.0	17.5	16.5	16.1	15.7
	10-29 Days	13.4	14.4	14.4	13.6	12.7	11.1	10.1
	30-59 Days	8.1	9.1	10.4	9.9	8.5	7.8	6.5
	60-89 Days	4.8	5.5	5.6	5.4	5.3	4.5	3.4
	90-129 Days	3.4	3.9	3.8	3.9	3.7	2.9	2.2
	130-180 Days	6.1	6.1	6.1	5.0	5.0	3.8	3.2
Volunteer outside the home	No Answer	8.4	6.6	6.2	7.2	6.8	6.7	6.8
	None	48.0	46.9	46.9	46.6	45.3	43.1	40.3
	1-9 Days	21.1	21.9	20.9	20.1	19.0	20.0	19.0
	10-29 Days	9.8	10.3	10.9	11.6	12.9	13.9	15.8
	30-59 Days	5.3	6.2	6.6	7.0	7.5	8.3	9.2
	60-89 Days	2.6	3.4	3.8	3.5	4.0	4.3	4.6
	90-129 Days	2.1	2.3	2.4	2.3	2.6	2.2	2.5
	130-180 Days	2.7	2.3	2.2	1.6	2.0	1.5	1.9
Volunteer inside a home	No Answer	7.4	6.0	5.9	6.8	6.4	6.3	6.6
	None	34.3	33.7	34.7	36.1	37.4	38.8	40.7
	1-9 Days	23.6	22.3	20.9	19.1	17.0	17.6	16.6
	10-29 Days	11.7	13.0	12.2	12.0	11.6	10.9	11.6
	30-59 Days	7.8	8.4	9.1	9.3	9.2	9.8	9.0
	60-89 Days	4.8	5.2	5.5	5.6	6.1	5.2	5.4
	90-129 Days	3.9	4.2	4.5	4.5	4.7	4.2	3.9
	130-180 Days	6.5	7.1	7.2	6.6	7.6	7.2	6.3

Note: The number of school days in a year is estimated as 180 days.

**Perceived Safety in Neighborhood and in School  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Out alone during the day	No Answer	4.8	4.5	4.2	5.1	5.0	4.3	4.8
	Very Safe	45.4	51.6	55.6	60.5	63.5	69.4	71.8
	Somewhat Safe	35.3	34.2	32.1	28.4	26.4	22.6	20.2
	Somewhat Unsafe	9.8	7.0	6.0	4.2	3.5	2.5	2.2
	Very Unsafe	4.7	2.7	2.1	1.7	1.6	1.2	1.1
Out alone after dark	No Answer	4.9	4.7	4.4	5.3	5.2	4.4	5.1
	Very Safe	12.4	16.4	18.5	22.4	24.5	28.6	29.5
	Somewhat Safe	27.4	32.4	36.3	37.0	37.8	39.0	40.1
	Somewhat Unsafe	25.3	25.8	24.4	23.0	21.9	19.8	17.6
	Very Unsafe	30.0	20.7	16.3	12.3	10.7	8.3	7.7
Inside home alone during the day	No Answer	5.1	5.1	4.7	5.7	5.4	4.6	5.1
	Very Safe	61.2	68.3	73.0	75.0	76.8	79.8	80.1
	Somewhat Safe	25.1	21.2	18.3	16.1	14.7	13.3	12.5
	Somewhat Unsafe	5.7	3.6	2.6	1.9	2.0	1.4	1.3
	Very Unsafe	2.9	1.7	1.4	1.3	1.1	0.8	0.9
Inside home alone after dark	No Answer	5.4	5.1	4.8	5.6	5.3	4.6	5.0
	Very Safe	32.1	38.9	43.2	46.4	48.5	51.7	53.4
	Somewhat Safe	29.6	30.6	31.7	30.8	30.4	29.8	29.0
	Somewhat Unsafe	18.5	16.3	14.0	12.0	11.0	10.3	9.3
	Very Unsafe	14.3	9.1	6.5	5.1	4.7	3.7	3.3
Traveling to/from school	No Answer	5.2	4.9	4.5	5.6	5.3	4.6	5.1
	Very Safe	60.0	63.0	63.8	66.2	66.6	69.6	71.3
	Somewhat Safe	24.2	24.2	24.4	23.2	23.4	22.3	20.7
	Somewhat Unsafe	6.4	5.2	4.8	3.5	3.1	2.3	1.9
	Very Unsafe	4.1	2.6	2.5	1.5	1.6	1.2	1.0
Staying in class	No Answer	4.9	4.7	4.5	5.4	5.2	4.5	4.9
	Very Safe	83.0	80.8	76.3	73.8	70.4	72.5	72.0
	Somewhat Safe	9.5	11.9	15.7	17.2	20.1	19.3	19.7
	Somewhat Unsafe	1.4	1.4	2.1	2.0	2.5	2.3	2.0
	Very Unsafe	1.2	1.3	1.5	1.5	1.8	1.4	1.3
Alone in school hallways	No Answer	5.0	4.9	4.6	5.6	5.3	4.6	5.0
	Very Safe	70.1	68.2	68.9	64.8	64.2	67.2	67.6
	Somewhat Safe	19.0	20.4	20.2	22.9	23.4	22.3	22.0
	Somewhat Unsafe	4.0	4.6	4.2	4.7	4.7	4.1	3.5
	Very Unsafe	2.0	1.9	2.1	2.0	2.3	1.8	1.8
Staying for afterschool activities	No Answer	5.4	5.3	4.9	5.8	5.5	4.9	5.3
	Very Safe	69.8	70.1	69.3	67.1	65.6	68.6	68.5
	Somewhat Safe	18.8	18.8	20.2	21.4	23.0	21.2	21.4
	Somewhat Unsafe	3.9	3.7	3.5	3.4	3.5	3.4	3.0
	Very Unsafe	2.0	2.1	2.1	2.2	2.4	1.9	1.8

**Annual Prevalence of Weapon Use and Violent Behavior  
by Indiana Female 6th - 12th Graders, 2007**  
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Physical fight	No Answer	5.1	5.0	4.4	5.6	5.3	4.8	5.0
	Never	73.0	71.9	67.0	71.1	72.2	76.8	80.4
	1-2 times	14.8	15.8	19.6	16.8	16.6	13.9	11.6
	3-9 times	4.4	5.0	6.3	4.7	4.4	3.4	2.2
	10+ times	2.7	2.3	2.7	1.7	1.5	1.1	0.7
Shouting argument	No Answer	5.4	5.3	4.7	5.9	5.5	4.8	5.3
	Never	49.0	42.5	35.3	35.5	34.8	36.8	39.8
	1-2 times	24.5	24.7	25.1	25.7	25.4	26.1	25.7
	3-9 times	11.6	14.3	18.3	18.3	18.9	18.9	18.3
	10+ times	9.5	13.2	16.6	14.7	15.3	13.4	10.8
Carried gun for hunting	No Answer	5.3	5.2	4.8	6.0	5.6	4.8	5.2
	Never	86.7	86.4	85.5	84.8	85.6	87.2	87.5
	1-2 times	4.3	4.7	5.5	4.7	4.7	4.3	4.0
	3-9 times	2.4	2.1	2.6	2.7	2.4	2.2	2.2
	10+ times	1.3	1.6	1.7	1.8	1.8	1.5	1.1
Carried gun for other reasons	No Answer	5.2	5.1	4.8	5.9	5.5	4.9	5.3
	Never	93.2	92.7	92.4	91.7	92.1	93.0	93.2
	1-2 times	0.9	1.3	1.8	1.4	1.3	1.2	0.9
	3-9 times	0.3	0.4	0.5	0.5	0.5	0.5	0.3
	10+ times	0.4	0.5	0.5	0.5	0.6	0.4	0.4
Carried weapon for hunting	No Answer	5.3	5.2	4.8	6.1	5.6	4.9	5.3
	Never	90.0	89.8	89.4	89.0	89.6	90.8	91.2
	1-2 times	3.1	3.0	3.5	2.7	2.6	2.4	1.7
	3-9 times	0.9	1.1	1.3	1.4	1.1	1.1	1.1
	10+ times	0.7	0.9	1.0	0.9	1.2	0.8	0.7
Carried weapon for other reasons	No Answer	5.4	5.2	4.9	6.1	5.6	4.9	5.2
	Never	91.2	90.5	88.3	88.2	88.3	89.6	89.8
	1-2 times	2.1	2.6	3.9	3.1	3.1	2.9	2.5
	3-9 times	0.7	1.0	1.5	1.3	1.4	1.1	0.9
	10+ times	0.6	0.7	1.3	1.3	1.6	1.4	1.6
Carried gun to school	No Answer	5.3	5.2	4.8	6.1	5.5	4.8	5.2
	Never	94.3	94.2	94.4	93.3	93.7	94.5	94.4
	1-2 times	0.2	0.3	0.4	0.3	0.3	0.3	0.2
	3-9 times	0.0	0.1	0.2	0.1	0.2	0.1	0.0
	10+ times	0.1	0.2	0.3	0.2	0.3	0.2	0.2
Carried weapon to school	No Answer	5.4	5.3	4.9	6.0	5.6	4.9	5.3
	Never	93.3	92.8	91.9	91.2	91.3	92.6	92.5
	1-2 times	1.0	1.3	2.1	1.6	1.6	1.3	1.0
	3-9 times	0.2	0.3	0.6	0.6	0.6	0.4	0.4
	10+ times	0.2	0.3	0.6	0.5	0.9	0.8	0.7

**Annual Prevalence of Weapon Use and Violent Behavior  
by Indiana Female 6th - 12th Graders, 2007**  
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Threatened by weapon on school property	No Answer	5.5	5.4	5.1	6.2	5.8	5.1	5.4
	Never	89.7	89.7	88.5	88.7	89.5	91.9	92.3
	1-2 times	3.8	3.8	5.1	4.0	3.5	2.3	1.8
	3-9 times	0.6	0.7	0.8	0.6	0.7	0.5	0.3
	10+ times	0.4	0.5	0.4	0.4	0.5	0.3	0.3
Property stolen while at school	No Answer	5.7	5.8	5.2	6.4	5.8	5.1	5.4
	Never	70.4	66.2	62.2	63.4	65.2	69.9	72.8
	1-2 times	18.1	21.6	24.6	24.0	23.1	20.7	18.1
	3-9 times	4.3	4.8	6.2	5.1	4.7	3.5	3.2
	10+ times	1.6	1.6	1.8	1.1	1.2	0.7	0.5
Missed school: Felt unsafe on school property	No Answer	5.5	5.5	5.0	6.2	5.7	5.0	5.4
	Never	90.1	89.4	88.8	88.7	87.3	90.5	89.1
	1-2 times	3.4	4.1	4.7	3.9	5.4	3.4	4.6
	3-9 times	0.6	0.6	0.9	0.7	1.0	0.7	0.6
	10+ times	0.4	0.4	0.5	0.5	0.5	0.4	0.3
Missed school: Felt unsafe on way to/from school	No Answer	5.3	5.3	4.9	6.1	5.7	5.1	5.4
	Never	91.9	92.2	91.8	91.4	91.3	92.7	92.2
	1-2 times	2.0	1.9	2.3	1.8	2.1	1.6	1.7
	3-9 times	0.4	0.4	0.6	0.3	0.5	0.3	0.4
	10+ times	0.4	0.3	0.5	0.4	0.5	0.3	0.3

**Gambling during the Last Twelve Months  
by Indiana Female 6th - 12th Graders, 2007**  
(Values expressed as percentages)

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Played cards	No Answer	4.3	4.0	4.2	5.5	5.3	5.0	5.2
	Not at all	86.2	84.1	79.9	79.6	79.2	80.8	79.8
	Less than once a month	6.1	8.0	10.5	10.4	10.7	9.9	10.9
	About once a month	2.1	2.4	3.6	3.2	3.2	3.0	2.9
	About once a week	0.9	1.0	1.2	1.1	1.0	1.0	0.9
	Daily	0.4	0.5	0.6	0.3	0.5	0.4	0.2
Bet on games	No Answer	4.5	4.0	4.2	5.7	5.5	5.2	5.2
	Not at all	84.9	82.8	79.4	79.1	80.6	82.9	83.8
	Less than once a month	6.9	8.7	10.7	10.3	9.6	8.5	7.9
	About once a month	2.1	2.7	3.6	3.3	2.8	2.4	2.3
	About once a week	1.0	1.1	1.3	1.1	1.0	0.7	0.5
	Daily	0.6	0.7	0.7	0.5	0.5	0.4	0.2
Bet on sports	No Answer	4.8	4.3	4.6	5.9	5.8	5.4	5.7
	Not at all	84.8	82.8	79.8	79.4	81.3	83.3	83.6
	Less than once a month	7.3	9.1	10.6	10.4	9.1	8.6	8.2
	About once a month	1.9	2.3	3.1	2.9	2.4	1.7	1.8
	About once a week	0.8	0.9	1.2	1.0	1.0	0.6	0.6
	Daily	0.5	0.6	0.7	0.3	0.5	0.4	0.2
Bought lottery tickets	No Answer	4.9	4.3	4.4	6.0	5.7	5.5	5.5
	Not at all	82.6	81.9	81.1	81.6	82.4	82.6	65.8
	Less than once a month	7.9	9.0	9.2	8.2	7.9	8.1	20.1
	About once a month	2.8	2.8	3.1	2.8	2.5	2.2	5.7
	About once a week	1.3	1.4	1.5	1.0	1.0	1.1	2.4
	Daily	0.5	0.6	0.7	0.4	0.5	0.5	0.6
Gambled in a casino	No Answer	4.8	4.4	4.4	5.8	5.7	5.4	5.4
	Not at all	94.5	94.7	94.4	93.2	93.1	93.3	92.9
	Less than once a month	0.5	0.5	0.7	0.5	0.6	0.9	1.2
	About once a month	0.1	0.1	0.2	0.2	0.1	0.1	0.2
	About once a week	0.1	0.1	0.1	0.1	0.1	0.1	0.1
	Daily	0.1	0.2	0.2	0.2	0.3	0.2	0.1
Played online	No Answer	4.7	4.3	4.5	5.9	5.8	5.4	5.4
	Not at all	93.3	93.5	93.0	92.2	92.4	93.3	93.3
	Less than once a month	1.0	1.2	1.3	1.1	0.9	0.7	0.6
	About once a month	0.4	0.4	0.5	0.4	0.4	0.3	0.2
	About once a week	0.3	0.3	0.2	0.2	0.2	0.1	0.2
	Daily	0.3	0.3	0.5	0.2	0.4	0.3	0.2

**Gambling during the Last Twelve Months  
by Indiana Female 6th - 12th Graders, 2007  
(Values expressed as percentages)**

		Grade						
		6th	7th	8th	9th	10th	11th	12th
Felt bad about the amount bet	No Answer	5.6	4.7	4.8	6.3	6.0	5.7	5.7
	Yes	3.4	3.2	2.8	2.2	1.4	1.3	1.5
	No	12.1	16.0	19.5	18.9	19.1	17.4	20.1
	I do not bet for money	78.8	76.1	72.9	72.6	73.5	75.7	72.8
Would like to stop betting but could not	No Answer	5.4	4.7	4.8	6.3	6.0	5.8	5.8
	Yes	2.9	2.6	2.5	1.5	1.1	0.7	0.8
	No	12.4	15.7	19.3	18.9	18.8	17.6	20.0
	I do not bet for money	79.3	76.9	73.5	73.4	74.2	75.9	73.5

## APPENDIX: INSTRUMENT



# Indiana Prevention Resource Center

# ALCOHOL, TOBACCO AND OTHER DRUG USE SURVEY

© COPYRIGHT 2007, 2005, 2002, 2001, 1995, 1993, 1991 The Trustees of Indiana University. All rights reserved.  
Questions # 26 - 28 used with permission from the Minnesota Department of Education.

This survey is funded, in part, by a contract with the Indiana Family and Social Services Administration, Division of Mental Health and Addiction, HHS / SAMHSA, CSAP, SAPT Block Grant.

## PRIVACY STATEMENT

This survey is completely confidential. Complete the survey in private and deposit your completed survey in the privacy envelope, as instructed by your teacher or supervisor. Do not place your name or any identifying marks on the survey form or envelope. There is no way for anyone to identify your individual responses. The survey envelopes will not be opened until they are received at the statewide processing center.

## YOUR PARTICIPATION IS VOLUNTARY

Your participation in this survey is voluntary. If you do not wish to answer a question, just leave it blank. If you do not wish to participate in this survey, you may:

- Return the entire survey form blank
- Answer questions with random answers and then answer "not truthfully at all" to the last question. Your form will be discarded and no one will know that you did not participate.
- Inform the teacher or supervisor that you choose not to participate in the survey.


**You will not be penalized in any way for deciding not to participate in this survey.** We do value your participation, however. Accurate results are dependent upon getting as many students as possible to volunteer to participate in this confidential survey. The results are used by many state and federal agencies to help determine how much money will be given to youth programs and to evaluate the effectiveness of drug programs in your community. Your help is important to this effort.

Thank you for your assistance.

SCHOOL CODE									
A	B	C	D	E	F	G	H	I	J
<input type="text"/>									
<input type="radio"/> 0									
<input type="radio"/> 1									
<input type="radio"/> 2									
<input type="radio"/> 3									
<input type="radio"/> 4									
<input type="radio"/> 5									
<input type="radio"/> 6									
<input type="radio"/> 7									
<input type="radio"/> 8									
<input type="radio"/> 9									

Please enter the unique ten-digit school code provided by your teacher or supervisor.

- Use a No. 2 pencil only.
- Do not use ink, ballpoint, or felt tip pens.
- Make solid marks that fill the circle completely.
- Erase cleanly any marks you wish to change.
- Make no stray marks on this form.
- Any mention of "drug" refers to *non-prescription drugs*.



CORRECT: ●

INCORRECT: ✓✗○

1. SEX:  Male  Female2. AGE:  7 yrs or younger  8 yrs  9 yrs  10 yrs  11 yrs  12 yrs  
 13 yrs  14 yrs  15 yrs  16 yrs  17 yrs  18 yrs or older3. GRADE:  6th  7th  8th  9th  10th  11th  12th4. ETHNICITY:  Hispanic  Non-Hispanic5. RACE:  White or Caucasian  Black or African-American  American Indian or Alaskan Native  
 Asian  Native Hawaiian or Other  Other  
Pacific Islander

## 6. IN THE PAST YEAR, HOW OFTEN HAVE YOU BEEN ABSENT FROM SCHOOL?

 Never  Once  Twice  3-5 days  6-10 days  11-19 days  20 or more days

## 7. DURING THE PAST YEAR, DID YOU ATTEND OR PARTICIPATE IN...(mark all that apply)?

 An "Afternoons R.O.C.K. in Indiana" after-school prevention program  
 Youth Leadership Programs (L.E.A.D. Initiative, Teen Institute, 4-H Junior Leaders, etc.)  
 S.A.D.D., S.T.A.N.D., or other prevention group at your school

## 8. HOW OFTEN IN THE PAST YEAR HAVE YOU PARTICIPATED IN ORGANIZED FAMILY ACTIVITIES (e.g., picnics, watching movies, trips, games, etc.)?

 Never  Once per week  Twice per week  3 or more times per week

## TOBACCO CIGARETTE USE

## 9. HAVE YOU EVER SMOKED CIGARETTES?

 Never Smoked  Occasionally but not regularly  Regularly now  
 Once or Twice  Regularly in the past

## 10. IF YOU HAVE EVER SMOKED CIGARETTES, AT WHAT AGE DID YOU FIRST SMOKE THEM?

 Never Smoked  7 yrs or younger  8 yrs  9 yrs  
 10 yrs  11 yrs  12 yrs  13 yrs  
 14 yrs  15 yrs  16 yrs  17 yrs  18 yrs or older

## 11. HOW OFTEN IN THE PAST YEAR HAVE YOU SMOKED CIGARETTES?

 None  A few times  1 to 5 cigarettes per day  
 About one half pack per day  About 1 pack per day  
 About 1 and a half packs per day  Two or more packs per day

## 12. HOW OFTEN IN THE PAST MONTH HAVE YOU SMOKED CIGARETTES?


 None  A few times  1 to 5 cigarettes per day  
 About one half pack per day  About 1 pack per day  
 About 1 and a half packs per day  Two or more packs per day

Continue on to the next page

PLEASE DO NOT WRITE IN THIS AREA



Serial

**Serial****DO NOT REPORT USE UNDER DOCTOR'S ORDERS FOR QUESTIONS ON THIS PAGE****USE OF ALCOHOL AND OTHER DRUGS****LIFETIME USE:**

13. HAVE YOU EVER USED...?

	Never	1-5 Times	6-19 Times	20-40 Times	More Than 40 Times
Snuff/Smokeless tobacco	<input type="radio"/>				
Cigars (tobacco)	<input type="radio"/>				
Pipe (tobacco, hookah, water-pipe)	<input type="radio"/>				
Alcohol (beer, wine, wine coolers, liquor)	<input type="radio"/>				
Marijuana (pot, weed, hash)	<input type="radio"/>				
Cocaine	<input type="radio"/>				
Crack	<input type="radio"/>				
Inhalants (glue, fumes, amyls)	<input type="radio"/>				
Amphetamines (uppers)	<input type="radio"/>				
Methamphetamines (meth, crank, crystal)	<input type="radio"/>				
Ritalin, Adderall (non-prescribed use only)	<input type="radio"/>				
Tranquilizers or Sleeping Pills (downers) (non-prescribed)	<input type="radio"/>				
Narcotics (opium, morphine, codeine, oxycontin) (non-prescribed)	<input type="radio"/>				
Heroin	<input type="radio"/>				
LSD (acid)	<input type="radio"/>				
MDMA (ecstacy, XTC, X)	<input type="radio"/>				
Other Psychedelics (psilocybin, mescaline, etc.)	<input type="radio"/>				
Rohypnol (roofies)	<input type="radio"/>				
GHB	<input type="radio"/>				
Steroids (non-prescribed)	<input type="radio"/>				
Over-the-counter drugs (non-medical use)	<input type="radio"/>				

**AGE OF FIRST USE:**

14. IF YOU HAVE EVER USED THESE DRUGS, AT WHAT AGE DID YOU FIRST USE THEM?

Never Used	7 or Younger	8 Yrs	9 Yrs	10 Yrs	11 Yrs	12 Yrs	13 Yrs	14 Yrs	15 Yrs	16 Yrs	17 Yrs	18 or Older
Snuff/Smokeless tobacco	<input type="radio"/>											
Cigars (tobacco)	<input type="radio"/>											
Pipe (tobacco, hookah, water-pipe)	<input type="radio"/>											
Alcohol (beer, wine, wine coolers, liquor)	<input type="radio"/>											
Marijuana (pot, weed, hash)	<input type="radio"/>											
Cocaine	<input type="radio"/>											
Crack	<input type="radio"/>											
Inhalants (glue, fumes, amyls)	<input type="radio"/>											
Amphetamines (uppers)	<input type="radio"/>											
Methamphetamines (meth, crank, crystal)	<input type="radio"/>											
Ritalin, Adderall (non-prescribed use only)	<input type="radio"/>											
Tranquilizers or Sleeping Pills (downers) (non-prescribed)	<input type="radio"/>											
Narcotics (opium, morphine, codeine, oxycontin) (non-prescribed)	<input type="radio"/>											
Heroin	<input type="radio"/>											
LSD (acid)	<input type="radio"/>											
MDMA (ecstacy, XTC, X)	<input type="radio"/>											
Other Psychedelics (psilocybin, mescaline, etc.)	<input type="radio"/>											
Rohypnol (roofies)	<input type="radio"/>											
GHB	<input type="radio"/>											
Steroids (non-prescribed)	<input type="radio"/>											
Over-the-counter drugs (non-medical use)	<input type="radio"/>											

Continue on to the next page

# DO NOT REPORT USE UNDER DOCTOR'S ORDERS FOR QUESTIONS ON THIS PAGE

## ANNUAL USE:

15. HOW MANY TIMES IN THE LAST YEAR HAVE YOU USED...

Never	1-5 Times	6-19 Times	20-40 Times	More Than 40 Times
-------	-----------	------------	-------------	--------------------

- Snuff/Smokeless tobacco
- Cigars (tobacco)
- Pipe (tobacco, hookah, water-pipe)
- Alcohol (beer, wine, wine coolers, liquor)
- Marijuana (pot, weed, hash)
- Cocaine
- Crack
- Inhalants (glue, fumes, amyls)
- Amphetamines (uppers)
- Methamphetamines (meth, crank, crystal)
- Ritalin, Adderall (non-prescribed use only)
- Tranquilizers or Sleeping Pills (downers) (non-prescribed)
- Narcotics (opium, morphine, codeine, oxycontin) (non-prescribed)
- Heroin
- LSD (acid)
- MDMA (ecstacy, XTC, X)
- Other Psychedelics (psilocybin, mescaline, etc.)
- Rohypnol (roofies)
- GHB
- Steroids (non-prescribed)
- Over-the-counter drugs (non-medical use)

## USE IN PAST MONTH:

16. HOW MANY TIMES IN THE LAST MONTH (30 DAYS) HAVE YOU USED...

Never	1-5 Times	6-19 Times	20-40 Times	More Than 40 Times
-------	-----------	------------	-------------	--------------------

- Snuff/Smokeless tobacco
- Cigars (tobacco)
- Pipe (tobacco, hookah, water-pipe)
- Alcohol (beer, wine, wine coolers, liquor)
- Marijuana (pot, weed, hash)
- Cocaine
- Crack
- Inhalants (glue, fumes, amyls)
- Amphetamines (uppers)
- Methamphetamines (meth, crank, crystal)
- Ritalin, Adderall (non-prescribed use only)
- Tranquilizers or Sleeping Pills (downers) (non-prescribed)
- Narcotics (opium, morphine, codeine, oxycontin) (non-prescribed)
- Heroin
- LSD (acid)
- MDMA (ecstacy, XTC, X)
- Other Psychedelics (psilocybin, mescaline, etc.)
- Rohypnol (roofies)
- GHB
- Steroids (non-prescribed)
- Over-the-counter drugs (non-medical use)

**Continue on to the next page**

## ALCOHOL USE

17. HOW MANY TIMES IN THE LAST TWO WEEKS HAVE YOU HAD FIVE OR MORE ALCOHOLIC DRINKS  
(beer, wine, liquor) AT A SITTING?

- None     Once     Twice     3 to 5 times     6 to 9 times     10 or more times

18. DURING THE PAST MONTH (30 days), HOW DID YOU USUALLY GET YOUR ALCOHOL?

- I did not drink alcohol during the past month (30 days).
- I bought it at a store such as a liquor store, convenience store, supermarket, discount store, or gas station.
- I bought it at a restaurant, bar, or club.
- I bought it at a public event such as a concert or sporting event.
- I gave someone else money to buy it for me.
- A person 21 years old or older gave it to me.
- I took it from a store.
- I got it from a family member.
- I got it some other way.

19. WHAT HAVE BEEN THE MOST IMPORTANT REASONS FOR YOUR DRINKING ALCOHOLIC BEVERAGES? (Mark all that apply.)

- |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <input type="radio"/> I did not drink alcoholic beverages.<br><input type="radio"/> To experiment (to see what it's like)<br><input type="radio"/> To relax or relieve tension<br><input type="radio"/> To feel good or get high<br><input type="radio"/> To seek deeper insights and understanding<br><input type="radio"/> To have a good time with my friends<br><input type="radio"/> To fit in with a group I like<br><input type="radio"/> To get away from my problems or troubles | <input type="radio"/> Because of boredom, nothing else to do<br><input type="radio"/> Because of anger or frustration<br><input type="radio"/> To get through the day<br><input type="radio"/> To increase the effects of some other drug(s)<br><input type="radio"/> To decrease (offset) the effects of some other drug(s)<br><input type="radio"/> To get to sleep<br><input type="radio"/> Because it tastes good<br><input type="radio"/> Because I am "hooked" (I feel I have to drink.) |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|


### CONSEQUENCES OF USE:

20. HOW MANY TIMES HAVE YOU EXPERIENCED THE FOLLOWING DUE TO YOUR DRINKING OR DRUG USE DURING THE PAST YEAR?

	Never	Once	Twice	3-5 Times	6-10 Times	11 or More Times
--	-------	------	-------	-----------	------------	------------------

- |                                                                                            |                       |                       |                       |                       |                       |                       |
|--------------------------------------------------------------------------------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Had a hangover                                                                             | <input type="radio"/> |
| Got nauseated or vomited                                                                   | <input type="radio"/> |
| Had a memory loss                                                                          | <input type="radio"/> |
| Performed poorly on a test or project                                                      | <input type="radio"/> |
| Missed school                                                                              | <input type="radio"/> |
| Got into a fight or argument                                                               | <input type="radio"/> |
| Damaged property, pulled fire alarms, etc.                                                 | <input type="radio"/> |
| Been in trouble with your parents                                                          | <input type="radio"/> |
| Been in trouble with the police                                                            | <input type="radio"/> |
| Driven a car while under the influence                                                     | <input type="radio"/> |
| Ridden in a car driven by someone who was "high" or had been using alcohol or drugs        | <input type="radio"/> |
| Forgot things you did while using alcohol or drugs                                         | <input type="radio"/> |
| Your family or friends ever told you that you should cut down on your drinking or drug use | <input type="radio"/> |
| Used alcohol or drugs to relax, feel better about yourself, or fit in                      | <input type="radio"/> |
| Used alcohol or drugs while you were by yourself or alone                                  | <input type="radio"/> |

**Continue on to the next page**

**Serial**

## INJECTION DRUG USE

21. HOW MANY TIMES HAVE YOU USED A NEEDLE OR SYRINGE TO INJECT A DRUG (non-medical use)?

Never	1-5 Times	6-19 Times	20-40 Times	More than 40 Times
-------	-----------	------------	-------------	--------------------

During your lifetime

During the last year

During the last month (30 days)

## BELIEFS AND PERCEPTIONS

### BELIEFS ABOUT DRUGS:

22. HOW MUCH DO YOU THINK PEOPLE RISK HARMING THEMSELVES (physically or in other ways) IF THEY...

No Risk	Slight Risk	Moderate Risk	Great Risk
---------	-------------	---------------	------------

- Smoke one or more packs of cigarettes per day
- Smoke marijuana (pot) occasionally
- Smoke marijuana regularly
- Use cocaine occasionally
- Use cocaine regularly
- Take one or two drinks of alcohol (beer, wine, liquor) occasionally
- Have five or more drinks once or twice each weekend

### PEER PERCEPTIONS:

23. HOW DO YOU THINK YOUR CLOSE FRIENDS FEEL (or would feel) ABOUT YOU DOING EACH OF THE FOLLOWING THINGS?

Strongly Approve	Approve	Don't Know	Disapprove	Strongly Disapprove
------------------	---------	------------	------------	---------------------

- Smoke one or more packs of cigarettes per day
- Smoke marijuana (pot) occasionally
- Smoke marijuana regularly
- Use cocaine occasionally
- Use cocaine regularly
- Take one or two drinks of alcohol (beer, wine, liquor) occasionally
- Have five or more drinks once or twice each weekend

### PARENTS/GUARDIANS' PERCEPTIONS:

24. HOW DO YOU THINK YOUR PARENTS/GUARDIANS FEEL (or would feel) ABOUT YOU DOING EACH OF THE FOLLOWING THINGS?

Strongly Approve	Approve	Don't Know	Disapprove	Strongly Disapprove
------------------	---------	------------	------------	---------------------

- Smoke one or more packs of cigarettes per day
- Smoke marijuana (pot) occasionally
- Smoke marijuana regularly
- Use cocaine occasionally
- Use cocaine regularly
- Take one or two drinks of alcohol (beer, wine, liquor) occasionally
- Have five or more drinks once or twice each weekend

**Continue on to the next page**

## AFTER SCHOOL HOURS (approximately 3 p.m. to 6 p.m.)

25. THERE ARE ABOUT 180 SCHOOL DAYS PER YEAR.

ON HOW MANY OF THOSE DAYS THIS YEAR WILL YOU SPEND MOST OF THE AFTER SCHOOL HOURS...

(mark all that apply)

### **Participating in...**

	None	1 to 9	10 to 29	30 to 59	60 to 89	90 to 129	130 to 180
or playing on official school sports teams	<input type="radio"/>						
official school arts, music, or drama programs	<input type="radio"/>						
other official school activities such as clubs and intramural	<input type="radio"/>						
supervised activities at youth centers (Boys & Girls Clubs, Parks Department, etc.)	<input type="radio"/>						
extended day programs at school	<input type="radio"/>						
other supervised youth activities (4-H, scouts, little league, etc.)	<input type="radio"/>						
classes such as music, art, computer, dance, sport lessons, etc.	<input type="radio"/>						

### **Socializing...**

with friends without adults supervision (e.g., going to the mall, hanging out in private homes, on streets, in parks, etc.)	<input type="radio"/>						
at home without adult supervision	<input type="radio"/>						
at home with adult supervision	<input type="radio"/>						

### **Working....**

for pay outside your home	<input type="radio"/>						
for pay inside a home (housekeeping, child care, etc.)	<input type="radio"/>						
as a volunteer outside your home	<input type="radio"/>						
as a volunteer inside a home (housekeeping, child care, etc.)	<input type="radio"/>						

## GAMBLING

26. DURING THE LAST 12 MONTHS, HOW OFTEN HAVE YOU DONE THESE ACTIVITIES?

	Not at all	Less than once a month	About once a month	About once a week	Daily
Played cards for money	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Bet money on games of personal skill like pool, golf, or bowling	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Bet money on sports teams	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Bought lottery tickets or scratch offs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Gambled in a casino	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Played online for money	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

27. DURING THE LAST 12 MONTHS, HAVE YOU EVER FELT BAD ABOUT THE AMOUNT YOU BET, OR ABOUT WHAT HAPPENS WHEN YOU BET MONEY?


Yes       No       I don't bet for money

28. DURING THE LAST 12 MONTHS, HAVE YOU EVER FELT THAT YOU WOULD LIKE TO STOP BETTING MONEY BUT DIDN'T THINK YOU COULD?

Yes       No       I don't bet for money

**Continue on to the next page**

PLEASE DO NOT WRITE IN THIS AREA



**Serial**

FORM SS-02

## Indiana Prevention Resource Center

**ALCOHOL, TOBACCO AND OTHER DRUG USE SURVEY**

© COPYRIGHT 2007, 2005, 2002, 2001, 1995, 1993, 1991 The Trustees of Indiana University. All rights reserved.  
Questions # 26 - 28 used with permission from the Minnesota Department of Education.

**SAFETY AND SECURITY****NEIGHBORHOOD ENVIRONMENT:**

29. IN YOUR NEIGHBORHOOD, HOW SAFE DO YOU FEEL...

	Very Safe	Somewhat Safe	Somewhat Unsafe	Very Unsafe
--	-----------	---------------	-----------------	-------------

Being out alone during the day

Being out alone after dark

Being inside your home alone during the day

Being inside your home alone after dark

While traveling to or from school

**SCHOOL ENVIRONMENT:**

30. IN YOUR SCHOOL, HOW SAFE DO YOU FEEL...

	Very Safe	Somewhat Safe	Somewhat Unsafe	Very Unsafe
--	-----------	---------------	-----------------	-------------

While in class

Alone in the hallways

Staying for after-school activities

**PERSONAL SAFETY ISSUES:**

31. IN THE PAST YEAR, HOW OFTEN HAVE YOU...

	Never	1-2 Times	3-9 Times	10 or More Times
--	-------	-----------	-----------	------------------

Been in a physical fight

Been in a serious argument involving shouting

Carried a gun for hunting or target practice

Carried a gun for other reasons

Carried some other weapon for hunting

Carried some other weapon for other reasons

Carried a gun to school

Carried some other weapon to school

Been threatened by a weapon on school property

Had your property stolen or damaged while on school property

Missed school because you felt unsafe on school property

Missed school because you felt unsafe on your way to or from school

32. HOW TRUTHFULLY HAVE YOU ANSWERED THESE QUESTIONS?

 Not truthfully at all Somewhat truthfully Completely truthfully

**THANK YOU  
FOR YOUR PARTICIPATION**



THE INDIANA PREVENTION RESOURCE CENTER  
2007 PREVALENCE STATISTICS  
MAIN FINDINGS

**ALCOHOL, TOBACCO AND OTHER DRUG USE BY INDIANA CHILDREN AND ADOLESCENTS**

Survey Conducted March & April 2007  
Report Dated August 2007

Conducted by:  
Indiana Prevention Resource Center

Published by:  
Institute for Drug Abuse Prevention

Department of Applied Health Science  
School of Health, Physical Education, and Recreation  
Indiana University

Funded, in part by a contract with the  
Indiana Family and Social Services Administration  
Division of Mental Health and Addiction, and the  
School of Health, Physical Education, and Recreation

Contact:  
**Indiana Prevention Resource Center**  
501 North Morton Street Suite 110  
Bloomington, IN 47404

<http://www.drugs.indiana.edu>  
[drugprc@indiana.edu](mailto:drugprc@indiana.edu)  
Toll Free: 1-800-346-3077  
Fax: 812-855-4940

